

PAPER_440 — Bubble Nebula NGC 7635: Per-System MUGE with E(t) GROWING Expansion and Low-Mass Central Star

Source: grok_share_68eb34022.txt — Document 12: “Master Universal Gravity Equation_The Bubble Nebula Evolution_03May2025.docx” (lines 3788–4125) **Session:** 119 **CP4 Class:** BubbleNebulaPerSystemMUGE_GrowingExpansion_Calculator (#95)

1. Overview

PAPER_440 delivers the **complete per-system MUGE** for the Bubble Nebula (NGC 7635) — a stellar wind nebula in Cassiopeia driven by the O-star SAO 20575 (BD+60°2522), $M_\star = 46 M_\odot$, at $d \approx 11$ kly. The bubble radius is $r \approx 5$ ly = 4.731×10^{16} m, with expansion age $\tau_{\text{exp}} = 4$ Myr.

Novel claim (Q1): First UQFF MUGE for NGC 7635 with a **GROWING expansion factor** $E(t) = E_0(1 - e^{-t/\tau_{\text{exp}}})$ — in contrast to the decaying erosion in Pillars of Creation (PAPER_435). Here E INCREASES from 0 to $E_0 = 0.1$ as the stellar wind bubble inflates, meaning the $(1 - E(t))$ suppression of self-gravity GROWS over time — physically representing the process where wind energy excavates an increasingly larger volume, reducing the effective gravitational restoring force at the bubble wall. Wind velocity $v_w = 1800$ km/s is also unique to this system (vs Wd2’s 2000 km/s).

2. System Parameters

Parameter	Symbol	Value
Central star mass	$M_\star$	$46 M_\odot = 9.149 \times 10^{31}$ kg
Bubble radius	r	5 ly = 4.731×10^{16} m
Expansion timescale	τ_{exp}	4 Myr = 1.262×10^{14} s
Max expansion factor	E_0	0.1
Magnetic field	B	10^{-6} T
Wind density	ρ_w	10^{-21} kg/m ³
Wind velocity	v_w	1.8×10^6 m/s (1800 km/s)
Fluid density	ρ_f	10^{-21} kg/m ³
Hubble constant	H_0	2.184×10^{-18} s ⁻¹

3. Growing Expansion Function

$$E(t) = 0.1 \left(1 - e^{-t/\tau_{\text{exp}}} \right)$$

Time	$E(t)$	$(1 - E(t))$	Physical meaning
$t = 0$	0	1.000	No wind yet, full self-gravity

Time	$E(t)$	$(1 - E(t))$	Physical meaning
$t = \tau = 4 \text{ Myr}$	0.0632	0.937	6.3% suppression
$t = \infty$	0.100	0.900	10% max suppression

Contrast with Pillars (PAPER_435): $E_{\text{PoC}}(t) = E_0 e^{-t/\tau}$ DECREASES (starts high, decays to 0). Here E GROWS — fundamentally different topology.

4. Complete 10-Term MUGE

$$g_{\text{Bub}}(r, t) = T_1(1 - E(t)) + T_2(1 - E(t)) + T_3 + T_4 + T_5 + T_6 + T_7 + T_8 + T_9 + T_{10}$$

T1 — Newtonian + $\mathbf{H} \mathbf{t} + \mathbf{B} \times (\mathbf{1} - \mathbf{E}(\mathbf{t}))$:

$$T_1 = \frac{GM_\star}{r^2} (1 + H_0 t) (1 - B/B_{\text{crit}}) (1 - E(t))$$

$$\frac{GM_\star}{r^2} = \frac{6.674 \times 10^{-11} \times 9.149 \times 10^{31}}{(4.731 \times 10^{16})^2} = \frac{6.104 \times 10^{21}}{2.238 \times 10^{33}} \approx 2.73 \times 10^{-12} \text{ m/s}^2$$

At $t = \tau_{\text{exp}} = 4 \text{ Myr}$: $T_1 \approx 2.73 \times 10^{-12} \times 0.937 \approx 2.55 \times 10^{-12} \text{ m/s}^2$

T2 — UQFF $\mathbf{Ug} \times (\mathbf{1} - \mathbf{E}(\mathbf{t}))$: $\approx 2 \times 2.73 \times 10^{-12} \times 1.1 \times 0.937 \approx 5.62 \times 10^{-12} \text{ m/s}^2$

T3-T8: All negligible or minor

T9 — Wind ram pressure:

$$T_9 = \frac{\rho_w v_w^2}{\rho_f} = \frac{10^{-21} \times (1.8 \times 10^6)^2}{10^{-21}} = 3.24 \times 10^{12} \text{ m}^2/\text{s}^2 \Rightarrow a_w = \frac{3.24 \times 10^{12}}{r} \approx 6.85 \times 10^{-5} \text{ m/s}^2$$

5. Canonical Numerical Result

$$g_{\text{Bub}}(t = 4 \text{ Myr}) \approx 6.85 \times 10^{-5} \text{ m/s}^2 \quad [\text{wind dominant by } 10^7 \times g_{\text{self}}]$$

Wind/gravity after expansion at $t = \tau_{\text{exp}}$:

$$\frac{a_w}{g_{\text{self}} \times (1 - E)} = \frac{6.85 \times 10^{-5}}{2.55 \times 10^{-12}} \approx 2.7 \times 10^7$$

The growing $E(t)$ means the bubble is progressively more dynamically unbound as it expands.

6. Uniqueness vs Prior Papers

Prior Paper	Overlap	New in PAPER_440
PAPER_435 (PoC)	Same E form but DECAYING	GROWING $E(t)$ — unique topology
PAPER_383	NGC 7635 brief	Full 10-term evaluation
None	$v_w = 1800$ km/s specific	Only system with this exact v_w

7. Comparison to Standard Model

Standard Weaver et al. (1977) stellar wind bubble model: $R \propto (L_w t^3 / \rho_0)^{1/5}$, purely kinematic. UQFF adds the $(1 - E(t))$ gravitational channel where the expanding bubble reduces the effective self-gravity — absent from Weaver’s model, in principle testable by comparing bubble shell deceleration profiles.

8. Testable Predictions

Q5 Prediction 1: $E_0 = 0.1$ with $\tau_{\text{exp}} = 4$ Myr predicts that at current age ~ 4 Myr, the self-gravity at the bubble wall is suppressed by 6.3% relative to if the star had never blown a wind. UQFF predicts this appears as a 6.3% deficit in the gas column density at the bubble wall vs predictions from simple r^{-2} falloff — measurable in Herschel dust emission maps.

Q5 Prediction 2: Wind velocity $v_w = 1800$ km/s (vs Wd2’s 2000 km/s) predicts a lower shock temperature $T_{\text{bub}} = m_p v_w^2 / (3k_B) \approx 2.4 \times 10^8$ K — softer X-ray spectrum ($kT \approx 2$ keV vs 3.5 keV for Wd2), testable with XMM-Newton or Chandra.

Q5 Prediction 3: At $t \rightarrow \infty$, $E \rightarrow E_0 = 0.1$ — UQFF predicts the bubble expansion velocity asymptotically decreases by 10% from the initial value as the full $(1 - E_0) = 0.9$ factor is reached. This predicts a $\sim 10\%$ deceleration in the observed bubble expansion rate at ages $\gg 4\tau = 16$ Myr, testable against very old WR nebulae around evolved massive stars.