

PAPER_276: UQFF

Date: March 2026 \$n = [int]# UQFF PAPER_276 ## Andromeda Friedmann-UQFF Gravity Coupling: $H(z) \cdot t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[Om_m*(1+z)^3+Om_L]/M
 $H_{\text{UQFF}}=H(z)*t_{\text{H}} \sim 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (G \cdot M / r) \cdot H(z) \cdot t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_{\text{H}} = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_{\text{H}} \sim 0.987$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273-275 established the Andromeda-specific physics: blueshift approach amplifier ?, HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z) \cdot t \sim 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \end{aligned}$$

$$= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc}$$

$$\text{In SI units: } H_{\text{SI}} = 69.969 \times 10^3 / 3.086 \times 10^{22} = \mathbf{2.269 \times 10^{-8} \text{ s}^{-1}}$$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = G M / r = 6.674 \times 10^{-11} \cdot 1.989 \times 10^{30} / (1.04 \times 10^{17})^2 = \mathbf{1.227 \times 10^{-8} \text{ m/s}^2}$ - At $t = 1 \text{ Gyr}$ ($3.156 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \cdot 2.269 \times 10^{-8} \times 3.156 \times 10^8 = \mathbf{8.79 \times 10^{-8} \text{ m/s}^2}$ - At $t = t_{\text{Hubble}}$ ($4.352 \times 10^9 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \approx 0.987 = \mathbf{1.211 \times 10^{-8} \text{ m/s}^2}$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^9 \text{ s}$ (Hubble time $\approx 13.8 \text{ Gyr}$).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987 \approx \mathbf{1}$ is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($\Omega_m + \Omega_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 \cdot t_H$ is the dimensionless Hubble number $\approx 0.96 \approx 1.0$ for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H \approx \mathbf{\text{Friedmann-UQFF Near-Unity Resonance Coefficient}}$

Parameter	Value
H_0	70.0 km/s/Mpc

Parameter	Value
O_m	0.3
O_?	0.7
z (Andromeda)	-0.001
H(z)	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^{17} \text{ s}$
H_UQFF	0.987 (~1)
g_expansion(t_H)	$1.211 \times 10^{-10} \text{ m/s}^2$
g_base	$1.227 \times 10^{-10} \text{ m/s}^2$
Ratio g_exp/g_base	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 10^6 = 1.989 \times 10^{-2} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned}
 a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
 &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
 &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2
 \end{aligned}$$

At ~9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. M_visible / M_DM_mass Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$M_{\text{visible}} = (1 - f_{\text{DM}}) \times M$$

$$M_{\text{DM, mass}} = f_{\text{DM}} \times M$$

For Andromeda defaults: $M = 1.989 \times 10^{14} \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^{14} = \mathbf{3.978 \times 10^{14} \text{ kg}}$ (20% visible baryons) - $M_{\text{DM, mass}} = 0.80 \times 1.989 \times 10^{14} = \mathbf{1.591 \times 10^{14} \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 26-32), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_total

The full UQFF 2.0 equation for Andromeda with all PAPER_273-276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at t=0 (current epoch):

Term	Value (m/s)	Paper
$g_{\text{grav}} = G M / r$	1.227×10^6	baseline
$U_{g, \text{sum}}$ (26 layers)	6.380×10^6	26-layer Triadic
?-term	2.998×10^6	cosmological
g_{quantum}	$\sim 2 \times 10^6$	HUP
g_{Lorentz}	$\sim 2 \times 10^6$	IGM EM
g_{fluid}	$\sim 8 \times 10^6$	IGM buoyancy
$F_{\text{res}}(?_{\text{HI}})$ at t=0	1×10^6	PAPER_274
g_{DM}	1.356×10^6	PAPER_275
$g_{\text{expansion}}$ at t=0	0	PAPER_276
a_{dust}	4.29×10^6	PAPER_276
?_approach	1.001001	PAPER_273

Note: $g_{\text{expansion}}$ grows linearly with t, dominating over g_{grav} at $t = t_{\text{H}}$.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to `exportState()`: 1. $H_0 = 70.0 \text{ km/s/Mpc}$ 2. $\Omega_{\text{m}} = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $\text{Mpc_to_m} = 3.086 \times 10^{22}$ 5. $\rho_{\text{dust}} = 1 \times 10^6 \text{ kg/m}^3$ 6. $M_{\text{visible}} = (1 - f_{\text{DM}}) M$ 7. $M_{\text{DM, mass}} = f_{\text{DM}} M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) \cdot t$. At $t = t_H$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
2. **$H_{\text{UQFF}} = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 \cdot t_H \approx 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value a blueshift Friedmann suppression.
3. **$M_{\text{visible}}/M_{\text{DM_mass}}$ cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{\text{dust}} \approx 4 \times 10^{-12}$ m/s²).

Watermark: ©2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value

Andromeda Friedmann-UQFF Gravity Coupling: $H(z) \cdot t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[0m_m*(1+z)^3+0m_L]/M
 $H_{\text{UQFF}}=H(z) \cdot t_H \approx 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (G \cdot M / r) \cdot H(z) \cdot t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_H \approx 0.987$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273-275 established the Andromeda-specific physics: blueshift approach amplifier λ , HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z) \cdot t \approx 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann H(z) Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0$ km/s/Mpc, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10^3 / 3.086 \times 10^{22} = 2.269 \times 10^{-18} \text{ s}^{-1}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = GM/r^2 = 6.674 \times 10^{-11} \times 1.989 \times 10^{30} / (1.04 \times 10^{22})^2 = 1.227 \times 10^{-18} \text{ m/s}^2$ - At $t = 1$ Gyr (3.156×10^8 s): $g_{\text{expansion}} = 1.227 \times 10^{-18} \times 2.269 \times 10^{-18} \times 3.156 \times 10^8 = 8.79 \times 10^{-28} \text{ m/s}^2$ - At $t = t_{\text{Hubble}}$ (4.352×10^{17} s): $g_{\text{expansion}} = 1.227 \times 10^{-18} \times 0.987 = 1.211 \times 10^{-27} \text{ m/s}^2$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^{17}$ s (Hubble time ≈ 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = 0.987$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987 \approx 1$ is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 \times t_H$ is the dimensionless Hubble number $\approx 0.96 \approx 1.0$ for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H \approx$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^7 \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{expansion}}(t_H)$	$1.211 \times 10^{-10} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-10} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0 \Rightarrow H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 106 = 1.989 \times 10^8 \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned}
a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
&= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
&= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2
\end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. $M_{\text{visible}} / M_{\text{DM_mass}}$ Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned}
M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\
M_{\text{DM, mass}} &= f_{\text{DM}} \times M
\end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM_mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 26-32), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_{total}

The full UQFF 2.0 equation for Andromeda with all PAPER_273-276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at $t=0$ (current epoch):

Term	Value (m/s ²)	Paper
$g_{\text{grav}} = G M / r$	$1.227 \times 10^{??}$	baseline
$U_{g, \text{sum}}$ (26 layers)	$6.380 \times 10^{??}$	26-layer Triadic
?-term	$2.998 \times 10^{??6}$	cosmological
g_{quantum}	$\sim 2 \times 10^{??6}$	HUP
g_{Lorentz}	$\sim 2 \times 10^{??6}$	IGM EM
g_{fluid}	$\sim 8 \times 10^{??6}$	IGM buoyancy
$F_{\text{res}}(?_{\text{HI}})$ at $t=0$	$1 \times 10^{??6}$	PAPER_274
g_{DM}	$1.356 \times 10^{??6}$	PAPER_275
$g_{\text{expansion}}$ at $t=0$	0	PAPER_276

Term	Value (m/s $\diamond$)	Paper
a_dust	4.29×10^{-10}	PAPER_276
γ_{approach}	1.001001	PAPER_273

Note: $g_{\text{expansion}}$ grows linearly with t , dominating over g_{grav} at $t = t_H$.

7. exportState Update (25 $\rightarrow$ 32 parameters)

Seven new parameters added to `exportState()`: 1. $H_0 = 70.0$ km/s/Mpc 2. $\Omega_m = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $\text{Mpc_to_m} = 3.086 \times 10^{22}$ 5. $\rho_{\text{dust}} = 1 \times 10^{-27}$ kg/m $\diamond$ 6. $M_{\text{visible}} = (1 - f_{\text{DM}}) M$ 7. $M_{\text{DM_mass}} = f_{\text{DM}} M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) \diamond t$. At $t = t_H$, adds 98.7% of g_{base} $\diamond$ near-gravitational doubling over cosmological time.
2. **$H_{\text{UQFF}} = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 \diamond t_H \diamond 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value $\diamond$ a blueshift Friedmann suppression.
3. **$M_{\text{visible}}/M_{\text{DM_mass}}$ cascade + dust drag:** Explicit DM split bookkeeping (`updateCache`) and ISM dust ram-pressure ($a_{\text{dust}} \diamond 4 \times 10^{-10}$ m/s $\diamond$).

Watermark: $\diamond$ 2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value ## Andromeda Friedmann-UQFF Gravity Coupling: $H(z) \diamond t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: `AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[0m_m*(1+z)^3+0m_L]/M`
 $H_{\text{UQFF}} = H(z) * t_H \sim 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (G \diamond M / r \diamond) \diamond H(z) \diamond t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals 69.969 km/s/Mpc $= 2.269 \times 10^{-8}$ s $^{-1}$. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7$ s, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_H \diamond$ **0.987** $\diamond$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273 275 established the Andromeda-specific physics: blueshift approach amplifier ?, HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaitre cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z) \propto t^{-1}$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10^3 / 3.086 \times 10^{22} = 2.269 \times 10^{-8} \text{ s}^{-1}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = G M / r = 6.674 \times 10^{-11} \times 1.989 \times 10^{30} / (1.04 \times 10^{22})^2 = 1.227 \times 10^{-8} \text{ m/s}^2$ - At $t = 1 \text{ Gyr}$ ($3.156 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 2.269 \times 10^{-8} \times 3.156 \times 10^8 = 8.79 \times 10^{-9} \text{ m/s}^2$ - At $t = t_{\text{Hubble}}$ ($4.352 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 0.987 = 1.211 \times 10^{-8} \text{ m/s}^2$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^{17}$ s (Hubble time ≈ 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987 \approx 1$ is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 \times t_H$ is the dimensionless Hubble number $\approx 0.96 \approx 1.0$ for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H \approx \mathbf{\text{Friedmann-UQFF Near-Unity Resonance Coefficient}}$

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^{17} \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{exp}}(t_H)$	$1.211 \times 10^{-8} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-8} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 \text{ kg/m}^3 / 106 = 1.989 \times 10^{-18} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned} a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2 \end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. M_visible / M_DM_mass Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned} M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\ M_{\text{DM, mass}} &= f_{\text{DM}} \times M \end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM, mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 26-32), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_total

The full UQFF 2.0 equation for Andromeda with all PAPER_273-276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at t=0 (current epoch):

Term	Value (m/s ²)	Paper
$g_{\text{grav}} = G M/r$	1.227×10^{-6}	baseline
Ug_{sum} (26 layers)	6.380×10^{-6}	26-layer Triadic
γ -term	2.998×10^{-6}	cosmological
g_{quantum}	$\sim 2 \times 10^{-6}$	HUP
g_{Lorentz}	$\sim 2 \times 10^{-6}$	IGM EM
g_{fluid}	$\sim 8 \times 10^{-6}$	IGM buoyancy
$F_{\text{res}}(\gamma_{\text{HI}})$ at $t=0$	1×10^{-6}	PAPER_274
g_{DM}	1.356×10^{-6}	PAPER_275
$g_{\text{expansion}}$ at $t=0$	0	PAPER_276
a_{dust}	4.29×10^{-6}	PAPER_276
γ_{approach}	1.001001	PAPER_273

Note: $g_{\text{expansion}}$ grows linearly with t , dominating over g_{grav} at $t = t_H$.

7. exportState Update (25 → 32 parameters)

Seven new parameters added to `exportState()`: 1. $H_0 = 70.0$ km/s/Mpc 2. $\Omega_{\text{m}} = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $\text{Mpc_to_m} = 3.086 \times 10^{22}$ 5. $\rho_{\text{dust}} = 1 \times 10^{-27}$ kg/m³ 6. $M_{\text{visible}} = (1 - f_{\text{DM}}) M$ 7. $M_{\text{DM_mass}} = f_{\text{DM}} M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

- Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) t$. At $t = t_H$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
- $H_{\text{UQFF}} = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 t_H = 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value a blueshift Friedmann suppression.
- $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{\text{dust}} \sim 4 \times 10^{-6}$ m/s²).

Watermark: ©2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value

Andromeda Friedmann-UQFF Gravity Coupling: $H(z)t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF: $g_{\text{exp}} = G M/r^2 H(z) t$; $H(z) = H_0 \sqrt{\Omega_{\text{m}}(1+z)^3 + \Omega_{\text{L}}}$ / M
 $H_{\text{UQFF}} = H(z) t_{H=0.987}$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (G \cdot M / r) \cdot H(z) \cdot t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_H = 0.987$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273-275 established the Andromeda-specific physics: blueshift approach amplifier, HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z) \cdot t_H \approx 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m (1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10^3 / 3.086 \times 10^{22} = 2.269 \times 10^{-8} \text{ s}^{-1}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = G M / r = 6.674 \times 10^{-11} \times 1.989 \times 10^{30} / (1.04 \times 10^{17}) = 1.227 \times 10^{-8} \text{ m/s}^2$ - At $t = 1 \text{ Gyr}$ ($3.156 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 2.269 \times 10^8 \times 3.156 \times 10^8 = 8.79 \times 10^{-8} \text{ m/s}^2$ - At $t = t_{\text{Hubble}}$ ($4.352 \times 10^9 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 4.352 \times 10^9 \approx 0.987 = 1.211 \times 10^{-8} \text{ m/s}^2$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^9 \text{ s}$ (Hubble time $\approx 13.8 \text{ Gyr}$).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987 \approx 1$ is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 \times t_H$ is the dimensionless Hubble number $\approx 0.96 \approx 1.0$ for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H \approx \mathbf{0.987}$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^9 \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{expansion}}(t_H)$	$1.211 \times 10^{-8} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-8} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 10^6 = 1.989 \times 10^{-2} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned} a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2 \end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. M_{visible} / $M_{\text{DM_mass}}$ Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned} M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\ M_{\text{DM_mass}} &= f_{\text{DM}} \times M \end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM_mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 26-32), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_total

The full UQFF 2.0 equation for Andromeda with all PAPER_273 276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at t=0 (current epoch):

Term	Value (m/s)	Paper
g_grav = G M/r	1.227×10^{-6}	baseline
Ug_sum (26 layers)	6.380×10^{-6}	26-layer Triadic
?-term	2.998×10^{-6}	cosmological
g_quantum	$\sim 2 \times 10^{-6}$	HUP
g_Lorentz	$\sim 2 \times 10^{-6}$	IGM EM
g_fluid	$\sim 8 \times 10^{-6}$	IGM buoyancy
F_res(?_HI) at t=0	1×10^{-6}	PAPER_274
g_DM	1.356×10^{-6}	PAPER_275
g_expansion at t=0	0	PAPER_276
a_dust	4.29×10^{-6}	PAPER_276
?_approach	1.001001	PAPER_273

Note: g_expansion grows linearly with t, dominating over g_grav at $t = t_H$.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to exportState(): 1. $H_0 = 70.0$ km/s/Mpc 2. $\Omega_m = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $M_{\text{pc_to_m}} = 3.086 \times 10^6$ 5. $\rho_{\text{dust}} = 1 \times 10^{-6}$ kg/m³ 6. $M_{\text{visible}} = (1 + f_{\text{DM}}) M$ 7. $M_{\text{DM_mass}} = f_{\text{DM}} M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

- Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) \cdot t$. At $t = t_H$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
- H_UQFF = 0.987:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 \cdot t_H \approx 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value a blueshift Friedmann suppression.
- M_visible/M_DM_mass cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{\text{dust}} \approx 4 \times 10^{-6}$ m/s²).

Watermark: ♦2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved

UQFF computed: MUGE buoyancy ratio $U_{bi}/F_U = [SSq] \diamond r \diamond /GM = 5.7e-1 \diamond 5.0e-4 = 2.85e-4$; compressed MUGE baseline $g = 5.4e-7 \text{ m/s} \diamond$ at r_{ISCO} .

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_\mu \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} =$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.064 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 17, \quad n_{\text{channel}} = 17/26$$

Since $p_{\text{DVP}} = 17$ is **sub-threshold** (threshold at $p > 26$), the system's vacuum topology inherits sub-threshold damping from the DVP lattice, producing smooth rather than resonant UQFF coupling profiles. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10^4 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.064	✓ Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 17$	✓ Sub-threshold
BSH layers	26 harmonic terms	$j = 1\dots 26,$ $\cos(2\pi j/26)$	✓ Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	✓ Canonical
[SSq]	0.57	Applied in BSH saturation	✓ Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.