

PAPER_319: Compact HII SFR Gravitational Binding Phase Transition

Author: Daniel T. Murphy **Date:** March 2026 ## $t_{\text{cross}} = 67,730 \text{ yr}$ | $s\text{SFR} = 5 \times 10^{-4} \text{ yr}^{-1}$ (50× Lagoon) | $m_{\text{factor}}(t_{\text{age}}) = 151$ ### FIRST UQFF Compact HII Region SFR Runaway Gravitational Binding Phase Transition

Session: 91

Module: ORION_UQFF_MODULE.cpp (33rd C++ module)

System: Orion Nebula M42/NGC 1976 — compact HII region with active star formation

Watermark: Copyright — Daniel T. Murphy, Session 91, March 2026

Abstract

This paper derives the SFR-driven gravitational binding phase transition for the Orion Nebula. With $\text{SFR} = 1 \text{ M}_{\text{sun}}/\text{yr}$ acting on an initial mass $M = 2000 \text{ M}_{\text{sun}}$, the specific star formation rate **$s\text{SFR} = 5 \times 10^{-4} \text{ yr}^{-1}$ is 50× that of the Lagoon Nebula** ($1 \times 10^{-5} \text{ yr}^{-1}$, PAPER_305). The system is born wind-dominated (unbound), but as SFR continuously adds mass, the effective gravitational acceleration amplified by the SFR mass factor $m_{\text{factor}}(t) = 1 + \text{SFR} \times t_{\text{yr}} / M_{\text{sun_count}}$ crosses the growing wind ram pressure at **$t_{\text{cross}} = 67,730 \text{ yr}$** , transitioning the nebula from unbound to gravitationally bound. By $t_{\text{age}} = 300 \text{ kyr}$, $\text{binding_ratio} = g_{\text{SFR}}/a_{\text{wind}} = 2.654$. This is the FIRST UQFF compact HII SFR runaway gravitational binding phase transition.

System Parameters

Parameter	Value	Description
M	$2000 \text{ M}_{\text{sun}} = 3.978 \times 10^{33} \text{ kg}$	Initial mass
SFR	$1 \text{ M}_{\text{sun}}/\text{yr} = 6.303 \times 10^{22} \text{ kg/s}$	Star formation rate
sSFR	$5 \times 10^{-4} \text{ yr}^{-1}$	Specific SFR = SFR/M
t_{age}	$3 \times 10^5 \text{ yr}$	Nebula age
v_{wind}	$8 \times 10^3 \text{ m/s}$	HII ionization front
g_{base}	$1.907 \times 10^{-11} \text{ m/s}^2$	Base gravity (PAPER_317)
$a_{\text{wind}}(t=0)$	$5.424 \times 10^{-10} \text{ m/s}^2$	Initial wind acceleration

Key Equations

SFR Mass Factor

$$M_{\text{sf}}(t) = \frac{\text{SFR}_{\text{yr}} \times t_{\text{yr}}}{M_{\text{sun_count}}}, \quad m_{\text{factor}}(t) = 1 + M_{\text{sf}}(t)$$

SFR-Amplified Gravitational Acceleration

$$g_{\text{SFR}}(t) = g_{\text{base}} \times m_{\text{factor}}(t) = g_{\text{base}} \left(1 + \frac{\text{SFR}_{\text{yr}} \cdot t_{\text{yr}}}{M_{\text{sun_count}}} \right)$$

Wind Acceleration (time-evolving)

$$a_{\text{wind}}(t) = \frac{v_{\text{wind}}^2}{r} \left(1 + \frac{t}{t_{\text{age}}} \right)$$

Crossover Time t_{cross} (PAPER_319 Key Result)

Setting $g_{\text{SFR}}(t) = a_{\text{wind}}(t)$:

$$g_{\text{base}} (1 + \text{sSFR} \cdot t) = a_{\text{wind},0} \left(1 + \frac{t}{t_{\text{age,yr}}} \right)$$

$$t_{\text{cross}} = \frac{a_{\text{wind},0} - g_{\text{base}}}{g_{\text{base}} \cdot \text{sSFR} - a_{\text{wind},0}/t_{\text{age,yr}}} = \boxed{67,730 \text{ yr}}$$

Specific SFR

$$\text{sSFR} = \frac{\text{SFR}_{\text{yr}}}{M_{\text{sun,count}}} = \frac{1}{2000} = 5 \times 10^{-4} \text{ yr}^{-1} = 50 \times \text{Lagoon}$$

Gas Depletion Timescale

$$t_{\text{consume}} = \frac{M_{\text{sun,count}}}{\text{SFR}_{\text{yr}}} = \frac{2000}{1} = 2000 \text{ yr} \quad (\text{without OMC-1 replenishment})$$

Results Summary

Quantity	Value	Significance
sSFR	$5 \times 10^{-4} \text{ yr}^{-1}$	50× Lagoon Nebula
t_{cross}	67,730 yr	Unbound → bound transition
$m_{\text{factor}}(t_{\text{age}}=300 \text{ kyr})$	151	SFR amplification
$g_{\text{SFR}}(t_{\text{age}})$	$2.878 \times 10^{-9} \text{ m/s}^2$	$151 \times g_{\text{base}}$
$a_{\text{wind}}(t_{\text{age}})$	$1.085 \times 10^{-9} \text{ m/s}^2$	
$\text{binding_ratio}(t_{\text{age}})$	2.654	Gravitationally bound
$m_{\text{factor}}(1 \text{ Myr})$	501	
$\text{binding_ratio}(1 \text{ Myr})$	4.069	Increasingly bound
t_{consume}	2000 yr	Gas depletion (no replenishment)

Phase Transition Diagram

$t=0$	$t_{\text{cross}}=67.7 \text{ kyr}$	$t_{\text{age}}=300 \text{ kyr}$	$t=1 \text{ Myr}$
UNBOUND	TRANSITION	BOUND $2.65 \times$	BOUND $4.07 \times$
$a_{\text{wind}} > g_{\text{SFR}}$	$a_{\text{wind}} = g_{\text{SFR}}$	$g_{\text{SFR}} > a_{\text{wind}}$	$g_{\text{SFR}} \gg a_{\text{wind}}$
$\eta=28.47 \text{ wind}$	— crossover —	$\eta_{\text{B}}=2.654$	$\eta_{\text{B}}=4.069$

Comparison: UQFF SFR Class

Module	SFR (M_{sun}/yr)	M (M_{sun})	sSFR (yr^{-1})	PAPER
Lagoon	0.1	10,000	1×10^{-5}	PAPER_305
Orion	1.0	2000	5×10^{-4}	PAPER_319
M16 (Eagle)	$\sim 0.01 \times$	1200	—	PAPER_284

Orion sSFR is $50 \times$ Lagoon — UQFF identifies this as the “ultra-compact HII” class with rapid gravitational binding crossover. Lagoon has $t_{\text{consume}} = 100 \text{ kyr}$; Orion has $t_{\text{consume}} = \mathbf{2000 \text{ yr}}$, the shortest gas depletion time in the UQFF series, sustained only by continuous OMC-1 giant molecular cloud inflow.

UQFF Physical Interpretation

The phase transition at $t_{\text{cross}} = 67,730 \text{ yr}$ is a structural boundary in Orion’s evolution:

- **Before t_{cross} :** The system is wind-dominated ($\eta_{\text{wind}} > 1$). UV photoionization and ram pressure drive a champagne flow outward; newly formed stars are subject to wind erosion.
- **After t_{cross} :** SFR has added sufficient mass that $g_{\text{SFR}} > a_{\text{wind}}$. The system becomes self-gravitating with respect to its own stellar formation. The cluster proceeds to form stars under its own gravity — consistent with the Orion OB1 association framework where the cluster is now 300 kyr old and gravitationally bound.

Within ORION_UQFF_MODULE.cpp, the SFR mass factor $m_{\text{factor}}(t)$ enters the base gravity term via $G * M * (1 + M_{\text{sf}}(t)) / r^2$, registered as WOLFRAM_TERM_ORION_BASE and WOLFRAM_TERM_ORION_SFR_BINDING.

WOLFRAM_TERM Registration

```
#define WOLFRAM_TERM_ORION_BASE(val) (val)
// g_base = (G*M*(1+M_sf(t))/r^2)*(1+Hz*t)*(1-B/B_crit)*(1+f_TRZ)
// M_sf(t) = SFR_yr*t_yr / M_sun_count [PAPER_319 SFR amplification]

#define WOLFRAM_TERM_ORION_SFR_BINDING(val) (val)
// wind = v_wind^2/r * (1+t/t_age) [crosses SFR gravity at t_cross=67730 yr]
```

Series first: *FIRST UQFF compact HII SFR runaway gravitational binding phase transition. Establishes sSFR as a new UQFF classification axis for HII regions: compact (Orion, $s\text{SFR} = 5 \times 10^{-4} \text{ yr}^{-1}$) vs extended (Lagoon, $s\text{SFR} = 1 \times 10^{-5} \text{ yr}^{-1}$).*

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_\mu \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} =$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.157 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 71, \quad n_{\text{channel}} = 8/26$$

Since $p_{\text{DVP}} = 71$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii

where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10^4 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.157	✓ Threshold-consistent
DVP prime BSH layers	$p_k \in \{2,3,\dots,113\}$ 26 harmonic terms	$p_{\text{DVP}} = 71$ $j = 1\dots 26,$ $\cos(2\pi j/26)$	✓ Resonant ✓ Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	✓ Canonical
[SSq]	0.57	Applied in BSH saturation	✓ Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.