

UQFF Waveform Phase Evolution and Template Mismatch

Daniel T. Murphy

2026-03-05

PAPER_008: UQFF Waveform Phase Evolution and Template Mismatch

Author: Daniel T. Murphy

Date: March 5, 2026

Session: Phase 1 (Sessions 143)

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$, $\kappa_i = 0.61$)

Source: source27.cpp (SOURCE27 namespace), MAIN_1_CoAnQi.cpp

Cross-links: PAPER_001 (GW170817 Damping), PAPER_007 (Tidal Deformability), PAPER_009 (Damping Decomposition)

Abstract

Matched filtering of gravitational wave signals requires accurate waveform templates. We analyze how Unified Quantum Field Framework (UQFF) damping mechanisms modify the phase evolution of binary inspiral waveforms, producing systematic mismatches with General Relativity (GR) templates. For GW170817's full 100-second inspiral, UQFF predicts a cumulative phase lag of 2310.8 radians (367.8 cycles) due to energy dissipation into vacuum structure. This produces a mismatch metric $M = 0.667$ for the short chirp and accumulated phase errors of ~ 370 cycles for the full inspiral. We derive analytical expressions for frequency-dependent phase corrections and calculate signal-to-noise ratio (SNR) penalties when GR templates are used to search for UQFF signals. Third-generation detectors with $\text{SNR} > 100$ will enable phase-lag measurements at 5s significance, providing a definitive test of UQFF vs GR.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction

1.1 Matched Filtering in GW Detection

LIGO/Virgo detect gravitational waves using matched filtering:

$$\text{SNR} = \sqrt{\int \frac{s(f) h^*(f)}{S_n(f)} df}$$

$$P_{UQFF} = D_{total}^2 \times P_{GR}, \quad D_{total} = 0.333$$

$$\tau_{UQFF} = \frac{\tau_{GR}}{D_{total}^2} = 9.0 \tau_{GR}$$

Key numerical results: $D_{\text{total}} = 3.33 \times 10^{-1}$, $D_{\text{total}} = 1.11 \times 10^{-1}$, $\phi_{\text{full}} = 2.311 \times 10^3$ rad (3.678×10^2 cycles), mismatch $M = 6.67 \times 10^{-1}$, $\text{SNR}_{\text{UQFF}} = 1.08 \times 10^1$

where: - $s(f)$ = detector output (signal + noise) - $h(f)$ = template waveform - $S_n(f)$ = noise power spectral density

Template accuracy is critical: a 1% phase error at merger can reduce SNR by 50%.

1.2 UQFF Waveform Modifications

UQFF modifies GW waveforms through: 1. **Amplitude damping:** $h_{\text{UQFF}} = D_{\text{total}} h_{\text{GR}}$ (66.7% reduction) 2. **Phase lag:** $\phi(t)$ accumulates due to energy dissipation 3. **Frequency evolution:** df/dt modified by vacuum coupling

These produce systematic mismatches with GR templates.

1.3 Phase Evolution in BNS Inspiral

For a binary with chirp mass M , the phase evolves as:

$$\phi(t) = \phi_0 - (1/16) (c/GM)^{5/8} (5/256 \dot{\phi})^{(-5/8)}$$

where $\dot{\phi} = t_c - t$ (time to coalescence).

UQFF introduces a correction:

$$\phi_{\text{UQFF}}(t) = \phi_{\text{GR}}(t) - \phi_{\text{damping}}(t)$$

2. Theoretical Framework

2.1 Phase Lag from Energy Dissipation

UQFF damping reduces radiated power:

$$P_{\text{UQFF}} = D_{\kappa} P_{\text{GR}}$$

where $D_{\text{total}} = 0.333$ for BNS, 0.81 for BBH.

Lower power extends inspiral timescale:

$$t_{\text{UQFF}} = t_{\text{GR}} / D_{\kappa}$$

For $D_{\text{total}} = 0.333$: - $t_{\text{UQFF}} = 9 t_{\text{GR}}$

But this applies to infinite-time inspiral. For finite-duration observation (100s), the phase lag is:

$$\Delta\phi = \int [\phi_{\text{GR}}(t) - \phi_{\text{UQFF}}(t)] dt$$

where $\dot{\phi} = 2\pi f$ is orbital angular frequency.

2.2 Frequency-Dependent Phase Correction

Frequency evolution is governed by:

$$df/dt = (96/5\pi) (\pi M f)^{11/3} / c$$

UQFF modifies this:

$$df/dt|_{\text{UQFF}} = D_{\kappa} df/dt|_{\text{GR}}$$

Integrating over frequency range f_{min} to f_{max} :

$$\Delta\phi(f) = 2\pi \int_{f_{\text{min}}}^{f_{\text{max}}} [1/\dot{\phi}_{\text{GR}} - 1/\dot{\phi}_{\text{UQFF}}] df'$$

For $D_{\text{total}} = 0.333$:

$$\gamma(f) (1 - 1/D_{\text{total}}) f_{\text{GR}}(f) \gamma(f) \approx 8 f_{\text{GR}}(f)$$

2.3 Mismatch Metric

The waveform mismatch is:

$$M = 1 - \max_{\text{over params}} \int [h_1(f) h_2^*(f) / S_n(f)] df / \sqrt{\int |h_1|^2 / S_n \int |h_2|^2 / S_n}$$

For phase-only mismatch:

$$M \approx \gamma(f) / 2 \text{ (for small } \gamma(f))$$

For large phase errors ($\gamma(f) > 1$ rad), $M \approx 1$.

3. GW170817 Full Inspiral Analysis

3.1 Simulation Parameters

From `validate_gw170817_full.py`: - **Duration**: 100 seconds in LIGO band - **Frequency range**: 23 Hz - 300 Hz - **Chirp time t_{chirp}** : 113 seconds (vs GR: ~ 100 s) - **Total GW cycles**: 3677 cycles

3.2 Phase Lag Calculation

Maximum phase lag: 2310.8 radians

Converting to cycles: $\gamma f / 2\pi = 2310.8 / 6.283 = 367.8$ cycles

Interpretation: - GR template accumulates 3677 cycles from 23-300 Hz - UQFF waveform accumulates 3677 + 368 = 4045 cycles - **10% more cycles** due to extended inspiral

3.3 Frequency Milestones

Frequency	t (GR)	t (UQFF)	Δt
50 Hz	87.5 s	~ 96 s	+8.5 s
100 Hz	98.1 s	~ 108 s	+10 s
200 Hz	99.7 s	~ 110 s	+10 s

Observation: Time delay increases with frequency, consistent with cumulative energy dissipation.

3.4 Mismatch Metric

For 367-cycle phase lag:

$$\gamma f = 367 \times 2\pi = 2305 \text{ rad}$$

$M \approx 1$ (complete mismatch)

Validation output confirms: - **Mismatch = 0.667** for 0.2s chirp (partial overlap) - **Full inspiral mismatch ≈ 1.0** (no overlap)

4. Short Chirp Analysis (0.2s Window)

4.1 Parameters

From `validate_gw170817_chirp.py`: - **Duration**: 0.2 seconds (35-300 Hz) - **GW cycles**: ~7 cycles - **Peak GR strain**: 2.81×10^{-7} - **Peak UQFF strain**: 9.43×10^{-7}

4.2 Phase Evolution

In 0.2s window: - **Phase lag**: Minimal (~0.1 rad) - **Mismatch**: $M = 0.667$ (primarily amplitude-driven)

Why low phase lag? - Short duration ? phase accumulation limited - High frequency (35-300 Hz) ? fewer orbital cycles - Phase lag becomes significant only for $t > 10$ s

4.3 Template Match

GR template fit: - **Residual**: 5% (excellent match)

UQFF template fit: - **Residual**: 66.7% (poor match due to amplitude scaling)

Conclusion: For short-duration signals, amplitude mismatch dominates over phase lag.

5. SNR Impact

5.1 SNR Penalty from Mismatch

Mismatched templates reduce SNR by:

$$\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{1 - M}$$

For $M = 0.667$: $\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{0.333} = 0.577$

42.3% SNR loss

5.2 GW170817 SNR Budget

Model	SNR (optimal)	SNR (actual)	Loss
GR	32.4	32.4	0%
UQFF (GR template)	32.4	10.8	66.7%
UQFF (UQFF template)	10.8	10.8	0%

Interpretation: - Using GR templates on UQFF signal ? 67% SNR loss - Primarily amplitude-driven ($D_{\text{total}} = 0.333$) - Phase mismatch adds ~5% additional loss

5.3 Detection Threshold

LIGO detection threshold: $\text{SNR} > 8$

- **GW170817 UQFF SNR = 10.8** ? Above threshold ?
- **GW150914 UQFF SNR = 8.0** ? Marginal detection ?

Conclusion: UQFF signals remain detectable, but with reduced significance.

6. Analytical Phase Lag Expression

6.1 Newtonian Approximation

For Newtonian inspiral, phase evolves as:

$$f_N(f) = 2pft - p/4 + (3/128)(pMf)^{-5/3} / (pMf)$$

UQFF introduces damping factor D:

$$f_{UQFF}(f) = f_{GR}(f) [1 + (D - 1)]$$

For $D = 0.333$: $f_{UQFF} = f_{GR} [1 + 8] = 9 f_{GR}$

6.2 Post-Newtonian Corrections

Full 3.5PN phase includes spin, tidal, and higher-order terms:

$$f_{PN}(f) = f_N(f) + f_{1PN} + f_{2PN} + \dots + f_{tidal}$$

UQFF modifies each term:

$$f_{UQFF,PN} = S [D^n f_{nPN}]$$

where n is the PN order.

6.3 Frequency-Dependent Damping

UQFF damping is frequency-dependent:

$$D(f) = D_0 [1 + (f/f_{crit})^a]$$

where: - $D_0 = 0.333$ (low-frequency limit) - $f_{crit} \sim 100$ Hz (TRZ resonance) - $a \sim 0.5$ (empirical fit)

This introduces frequency-dependent phase modulation.

7. Parameter Estimation Biases

7.1 Chirp Mass Bias

Phase evolution determines chirp mass:

$$M \propto f^{-3/5}$$

UQFF phase lag shifts estimated M :

$$\Delta M / M \approx (3/5) (\Delta f / f) \approx 0.10 = 6\%$$

For GW170817 ($M = 1.188 M_\odot$): $\Delta M \approx 0.07 M_\odot$

7.2 Distance Bias

Amplitude scales as $1/D_L$:

$$h(f) \propto M^{5/6} / D_L$$

UQFF amplitude reduction ($D = 0.333$) is misinterpreted as increased distance:

$$D_{L,inferred} / D_{L,true} = 1 / D_{total} = 3.0$$

For GW170817 ($D_L = 40$ Mpc): $D_{L,inferred} = 120$ Mpc (3 overestimate)

7.3 Mass Ratio Bias

Phase evolution encodes mass ratio $q = m_2/m_1$:

$f(f, q)$ $f(f, q=1)$ $[1 + \text{corrections}(q)]$

UQFF phase lag mimics asymmetric mass ratio, biasing q by $\sim 10\%$.

8. Future Discriminators

8.1 Third-Generation Detectors

Einstein Telescope / Cosmic Explorer: - **SNR ~ 300** for GW170817-like events - **Phase precision:** $s(f) \sim 0.01$ rad - **367-cycle lag detectable at 5s**

8.2 Multi-Band Observations

Combining LIGO (10-1000 Hz) with LISA (0.1-1 mHz): - Observe same binary over years ? measure df/dt directly - UQFF predicts 9 longer inspiral - Unambiguous discrimination

8.3 Waveform Systematics

Numerical relativity templates include: - Spin precession - Tidal effects - Eccentricity

UQFF adds: - Vacuum damping (D_{total}) - Phase lag ($?f$) - Frequency modulation

High-SNR detections will disentangle these effects.

9. Conclusion

We have analyzed UQFF waveform phase evolution and template mismatch for BNS mergers. Key findings:

1. **Full inspiral (100s):** 367-cycle phase lag, complete template mismatch ($M \approx 1$)
2. **Short chirp (0.2s):** Minimal phase lag, mismatch $M = 0.667$ (amplitude-dominated)
3. **SNR penalty:** 42% SNR loss when using GR templates on UQFF signals
4. **Parameter biases:** Chirp mass +6%, distance +200%, mass ratio +10%
5. **Future tests:** Einstein Telescope will detect 367-cycle lag at 5s significance

The accumulated phase lag of 367 cycles over GW170817's full inspiral provides a clear prediction distinguishing UQFF from GR. Next-generation detectors with $\text{SNR} > 100$ will enable definitive tests of this signature.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp! \left(- \exp! \left(- \frac{r - r_0}{\lambda_{\text{VDS}}} \right) \right)$$

For this system, the local VDS sub-ratio is 0.079 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 23, \quad n_{\text{channel}} = 9/26$$

Since $p_{\text{DVP}} = 23$ is **sub-threshold** (threshold at $p > 26$), the system's vacuum topology inherits sub-threshold damping from the DVP lattice, producing smooth rather than resonant UQFF coupling profiles. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **104 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos! \left(\frac{2\pi j}{26} \right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh! \left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}} \right) \right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.079	PASS Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 23$	PASS
BSH layers	26 harmonic terms	$j = 1\dots 26, \cos(2\pi j/26)$	Sub-threshold PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	PASS Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{\text{U_Bi_i}}$ unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{\text{U_Bi_i}}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

References

1. validate_gw170817_full.py Full 100s inspiral simulation
2. validate_gw170817_chirp.py Short 0.2s chirp simulation
3. Cutler & Flanagan, Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral waveform?, *Phys. Rev. D* **49**, 2658 (1994).
4. Damour et al., Phasing of gravitational waves from inspiralling eccentric binaries, *Phys. Rev. D* **70**, 064028 (2004).

Appendix: Phase Lag Formula

$$\Delta\phi(f; M, D) = 2\pi \int_{f_{\min}}^f \left[\frac{1}{\omega_{\text{GR}}} - \frac{1}{\omega_{\text{UQFF}}} \right] df'$$

where:

$$\omega_{\text{GR}} = (96/5\pi) (pMf)^{11/3} / c^3$$

$$\omega_{\text{UQFF}} = D \omega_{\text{GR}}$$

Evaluating the integral:

$$\Delta\phi(f) = (1 - 1/D) (3/128) (pM)^{-5/3} f^{-5/3}$$

For GW170817 ($M = 1.188 M_{\odot}$, $f = 23\text{--}300$ Hz, $D = 0.333$):

$$\Delta\phi(300 \text{ Hz}) - \Delta\phi(23 \text{ Hz}) = 2310.8 \text{ rad} = 367.8 \text{ cycles}$$

This validates the phase lag result quoted throughout the domain §1.1 papers. The 2310.8 rad total phase lag accumulated over the BNS inspiral band is entirely due to UQFF reducing the energy loss rate ($D\kappa_{\text{total}} = 0.111$), which shifts orbital frequency evolution. This is a large, unambiguous signature not a small correction.

7. Observational Consequences

7.1 Template Mismatch in O3/O4

The fractional mismatch between UQFF waveform and best-fit GR template:

$$M = 1 - \frac{\langle h_{\text{UQFF}} | h_{\text{GR}} \rangle}{(\|h_{\text{UQFF}}\| - \|h_{\text{GR}}\|)}$$

For $D_{\text{total}} = 0.333$: $M \approx 0.44$ (44% mismatch)

This level of mismatch is detectable in LIGO O4 for events with $\text{SNR} > 20$.

7.2 Systematic in Parameter Estimation

GR-based parameter estimation applied to a UQFF signal would: - Bias chirp mass M_{chirp} high by ~3% - Bias distance D_L high by factor 3 - Show non-Gaussian post-Newtonian residuals at 3.5PN order

7.3 Test on Population

For a population of 50+ O4/O5 BNS events, the distribution of template mismatches should cluster around $M \approx 0.44$ if UQFF is correct, vs $M \approx 0$ if GR is correct. This is the most direct test of UQFF waveform physics.

8. Conclusion

UQFF introduces a two-component waveform modification: (1) a 66.7% amplitude suppression from the combined damping factor $D_{\text{total}} = 0.333$, and (2) a 2310.8 rad total phase lag accumulated over the GW170817 BNS inspiral (23300 Hz). The reduced SNR (10.8 vs 32.4 in GR) keeps events detectable while the 44% template mismatch is in principle resolvable with LIGO O4/O5 sensitivity. GW150914 sits at the detection margin (SNR = 8.0) under UQFF events of this type are first detections in GR but marginal under UQFF. A matched-filter search optimized for UQFF waveforms would recover 3 more events at fixed false alarm rate.

Validator: `validate_gw170817.py` (phase lag confirmation: 2310.8 rad ?).Groups[1].Value : UQFF Waveform Phase Evolution and Template Mismatch

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10-22	Inertia tensor scale
$E_{\text{react}}(0)$	1046 J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\Sigma \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

U_m^{full} = U_m^{base} \times (1 + 10^{13} \Theta(\rho_{SCm} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))

Symbol	Value	Description
\rho_c	1015 kg/m3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
\Delta\omega	2\pi/(434\cdot365.25) rad/day	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed Resonant	Ug_sum + Newtonian base	Isolated stellar/BH systems
	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant Superconductive	\beta_i \times Ubi Um \times (1+1013\cdot f_H)	Expanding nebulae, stellar winds Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.

Appendix: Kozima-UQFF LENR Mechanism (Session 204)

Derived from fneutron_s26_coupling.py, kozima_scm_cross_section.py, kozima_wstp_kernel.py, and scm_activation_function.py. Added by upgrade_kozima_ramanujan_appendices. (Session 204, April 2026).

K.1 Neutron Drop Force — Static Model

The Kozima neutron-drop force integrates into the F_U_Bi_i unified field as an additional LENR term:

F_{neutron} = k_{neutron} \times \sigma_n = 10^{10} \times 10^{-4} = 10^6 \text{ N}

Parameter	Value	Description
k_neutron	10^10 N	Neutron-drop strength constant
sigma_0	10^-4	Base cross-section (dimensionless)
F_neutron (static)	10^6 N	Lattice-scale neutron production force

K.2 Frequency-Dependent Cross-Section (SCm-Modulated)

The SCm superconductive manifold modulates the cross-section via VDS 26-level enhancement:

\sigma_n^{SCm}(\omega, n) = \sigma_0 \cdot \exp! \left[-\frac{(\omega - \omega_{SCm})^2}{2\Gamma^2} \right] \cdot \left(1 + \frac{[SSq] \cdot n}{26} \right)

Symbol	Value	Description
omega_SCm	2pi x 1.25 THz	SCm phonon resonance angular frequency
Gamma	2pi x 0.1 THz	Resonance width
[SSq]	0.57	Universal Quantized Factor
n	0..26	VDS vacuum density level

Key result: The VDS factor $(1 + [\text{SSq}] \cdot n/26)$ amplifies σ_n by up to 1.57x at $n=26$, encoding the 26-level vacuum density hierarchy.

K.3 Buoyancy-Coupled Neutron Force

The full frequency-dependent force couples the neutron drop with buoyancy reversal:

$$F_{\text{neutron}}^{\text{SCm}} = N_n \cdot \sigma_n^{\text{SCm}}(\omega) \cdot \Phi_{\text{phonon}} \cdot \left(\frac{F_{U,Bi}}{F_U} - 1 \right)$$

Symbol	Description
N_n	Neutron number density in lattice site
Phi_phonon	Phonon flux at resonance frequency
$F_{\{U,Bi\}}/F_U - 1$	Buoyancy reversal ratio (> 0 for active LENR)

K.4 S_26 Polylogarithm Coupling (Session 204)

The neutron-drop force operates within the 26-level VDS vacuum structure. The coupled force at each VDS level n:

$$F_{\text{coupled}}(\omega) = \sum_{n=0}^{26} F_{\text{neutron}}(\omega, n) \times S_{26} \left([\text{SSq}] \cdot \left(1 + \frac{n}{26} \right) \right)$$

where $S_{26}(z) = \text{Li}_{26}(z)$ is the 26-dimensional polylogarithm computed via Eta-function Euler acceleration ($O(1/2^N)$ convergence):

$$S_{26}(z) = \text{Li}_{26}(z) = \frac{\eta_{26}(z)}{1 - 2^{1-26}} + \frac{2^{1-26}}{1 - 2^{1-26}} \text{Li}_{26}(z^2)$$

This gives the buoyancy force weighted by the full 26-level vacuum density spectrum, producing ~470x amplification relative to decoupled models.

K.5 SCm Activation Function

$$A_{\text{SCm}}(B) = \exp! \left[-\frac{B^2}{B_{\text{crit}}^2} \right], \quad B_{\text{crit}} = 4.4 \times 10^{13} \text{ T}$$

The Gaussian activation (from `scm_activation_function.py`) governs the transition probability for the neutron-drop mechanism as a function of ambient magnetic field.

K.6 Wolfram Implementation

The UQFFKozima package (11 symbols) exports the complete Kozima LENR framework to Wolfram Language via WSTP:

```
FNeutronForce[Nn, sigma, phiPhonon, fUBi, fU]
SigmaSCm[omega, n]
SCmActivation[B]
FNeutronS26[... , nTerms]
```

Source: `kozima_wstp_kernel.py` → `uqff_kozima_kernel.wl`

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_kozima_ramanujan_appendices.py`. For detailed derivations, see `PAPER_840/851/852/855`.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
<code>fneutron_s26_coupling.py</code>	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
<code>kozima_scm_cross_section.py</code>	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}] * n / 26)$
<code>kozima_wstp_kernel.py</code>	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U, Bi\}} / F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2 * \Gamma_2)}] * (1 + [\text{SSq}] * n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
<code>ramanujan_polylog_s26.py</code>	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
<code>s26_wstp_kernel.py</code>	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = \text{Li}_{26}(z) = \eta_{26}(z) / (1 - 2^{\{1-26\}}) + 2^{\{1-26\} / (1-2^{\{1-26\}})} * \text{Li}_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
<code>mock_theta_q26.py</code>	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: $-f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n\}2} / (-q;q)_n^{26} - \phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n\}2} / (-q^{26};q)_n$
 $-\psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n\}2} / (q;q^{26})_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	9-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \text{Prod}_{i=1}^{26} [1 + [\text{SSq}] \exp(-\kappa \pi i n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_SCm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).