

Sombrero SMBH Dominance Ratio γ_{BH} and UQFF Sphere of Influence r_{SOI}

Daniel T. Murphy

2026-03-01

PAPER_279 — Sombrero SMBH Dominance Ratio γ_{BH} and UQFF Sphere of Influence r_{SOI}

Date: March 2026

Author: Daniel T. Murphy **Module:** SOMBRERO_UQFF_MODULE.cpp (UQFF 2.0) **Session:** 77 — March 2026 **Framework:** Unified Quantum Field Framework (UQFF 2.0) **Status:** Complete — embedded in SOMBRERO_UQFF_MODULE.cpp **Whitepaper Series:** 279/1000 **DOI (Provisional):** UQFF-2026-279-BH

Abstract

The Sombrero Galaxy (M104) harbours one of the most massive black holes relative to its host galaxy mass of any object in the Local Universe: $M_{\text{BH}} = 109 M_{\text{sun}}$ in a galaxy of total mass $M = 1011 M_{\text{sun}}$, giving a **SMBH Dominance Ratio** $\gamma_{\text{BH}} = M_{\text{BH}}/M = 0.01$ (1%). For comparison, the Milky Way's Sgr A* has $\gamma_{\text{BH}} \approx 4 \times 10^{-5}$ (~0.004%); Sombrero's SMBH is **250× more dominant relative to its host**. Within the UQFF framework, we define the **UQFF Sphere of Influence** $r_{\text{SOI}} = r \times \sqrt{\gamma_{\text{BH}}}$, the radius at which the central BH's direct gravitational contribution equals the total galaxy contribution at the reference radius. For Sombrero, $r_{\text{SOI}} = 2.36 \times 10^{19} \text{ m}$ — a precise UQFF prediction setting the boundary inside which BH gravity exceeds galaxy-mean gravity in the UQFF model.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Observational Motivation

1.1 The Sombrero SMBH

The Sombrero Galaxy's central black hole mass has been measured through stellar and gas kinematics:

- Ford et al. (1996): $M_{\text{BH}} = (1.0 \pm 0.75) \times 109 M_{\text{sun}}$ (gas kinematics, HST)
- Kormendy et al. (1996): confirmation from stellar dynamics
- Jardel et al. (2011): $M_{\text{BH}} \approx 6.6 \times 10^8 M_{\text{sun}}$ (JAM modelling, consistent with ~109 range)

We adopt $M_{\text{BH}} = 1.0 \times 109 M_{\text{sun}} = 1.989 \times 10^{39} \text{ kg}$.

The total galaxy mass (stars + gas + DM within the reference radius) is $M = 1011 M_{\text{sun}} = 1.989 \times 10^{41} \text{ kg}$.

1.2 Why γ_{BH} Matters

In standard astrophysical models, BH sphere-of-influence calculations use the velocity dispersion ($r_{\text{SOI}} = GM_{\text{BH}}/\sigma^2$). The UQFF framework generalises this to a mass-ratio-based definition consistent with the 26-layer Triadic gravity structure, yielding a dimensionless parameter γ_{BH} that naturally encodes the BH's dominance within the UQFF vacuum field hierarchy.

2. Theoretical Derivation

2.1 SMBH Dominance Ratio

We define the dimensionless **SMBH Dominance Ratio**:

$$\gamma_{\text{BH}} = \frac{M_{\text{BH}}}{M}$$

For the Sombrero Galaxy:

$$\gamma_{\text{BH}}^{\text{Sombrero}} = \frac{10^9 M_{\odot}}{10^{11} M_{\odot}} = 0.01 = 1\%$$

2.2 SMBH Direct Gravitational Contribution

The BH contribution to g_{total} at the reference radius r :

$$g_{\text{BH}} = \frac{GM_{\text{BH}}}{r^2} = \gamma_{\text{BH}} \cdot \frac{GM}{r^2} = \gamma_{\text{BH}} \cdot g_{\text{base}}$$

Substituting $\gamma_{\text{BH}} = 0.01$ and $g_{\text{base}} = 2.382 \times 10^{-10} \text{ m/s}^2$:

$$g_{\text{BH}} = 0.01 \times 2.382 \times 10^{-10} = 2.382 \times 10^{-12} \text{ m/s}^2$$

2.3 UQFF Sphere of Influence

The **UQFF Sphere of Influence** r_{SOI} is defined as the radius $r' < r$ where the BH gravitational contribution equals the total galaxy contribution at r :

$$g_{\text{BH}}(r') = g_{\text{base}}(r)$$

$$\frac{GM_{\text{BH}}}{r'^2} = \frac{GM}{r^2}$$

Solving:

$$r'^2 = r^2 \cdot \frac{M_{\text{BH}}}{M} = r^2 \cdot \gamma_{\text{BH}}$$

$$r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$$

For Sombrero:

$$r_{\text{SOI}} = 2.36 \times 10^{20} \cdot \sqrt{0.01} = 2.36 \times 10^{20} \times 0.1 = 2.36 \times 10^{19} \text{ m}$$

Physical interpretation: Within $r_{\text{SOI}} = 2.36 \times 10^{19} \text{ m}$ ($\sim 2.49 \text{ kly}$), the direct BH gravitational acceleration exceeds the galaxy-mean g_{base} . This is the UQFF-predicted boundary of BH gravitational dominance.

2.4 Verification

At $r' = r_{\text{SOI}}$:

$$g_{\text{BH}}(r_{\text{SOI}}) = \frac{GM_{\text{BH}}}{r_{\text{SOI}}^2} = \frac{G \cdot 0.01M}{(0.1r)^2} = \frac{0.01 \cdot GM}{0.01 \cdot r^2} = \frac{GM}{r^2} = g_{\text{base}} \text{ PASS}$$

3. Comparative SMBH Dominance Table

Galaxy	M _{BH} (M _{sun})	M _{total} (M _{sun})	γ_{BH}	$r_{\text{SOI}} / r_{\text{ref}}$
Milky Way (Sgr A*)	$\sim 4 \times 10^6$	$\sim 10^{11}$	$\sim 4 \times 10^{-5}$	$\sim 6.3 \times 10^{-3}$
Andromeda (M31)	$\sim 1.4 \times 10^8$	$\sim 10^{12}$	$\sim 1.4 \times 10^{-4}$	$\sim 1.2 \times 10^{-2}$
M87	$\sim 6.5 \times 10^9$	$\sim 6 \times 10^{12}$	$\sim 1.1 \times 10^{-3}$	$\sim 3.3 \times 10^{-2}$
Sombrero (M104)	$\sim 10^9$	$\sim 10^{11}$	0.01	0.1

Sombrero's $\gamma_{\text{BH}} = 0.01$ is the highest of any nearby well-measured galaxy in the UQFF catalogue, making it the dominant test-case for SMBH-galaxy UQFF coupling.

Key comparison ratios: - Sombrero/Sgr A*: γ_{BH} ratio = $0.01/4 \times 10^{-5} = \mathbf{250\times}$ - Sombrero/M87: γ_{BH} ratio = $0.01/1.1 \times 10^{-3} \approx \mathbf{9\times}$ - Sombrero/Andromeda: γ_{BH} ratio = $0.01/1.4 \times 10^{-4} \approx \mathbf{71\times}$

4. UQFF BH Contribution in the Master Equation

The BH term enters computeG() as an additive contribution alongside the 26-layer Triadic sum:

$$g_{\text{total}} = [\dots + g_{\text{BH}} + \dots] \cdot \kappa_{\text{recession}} \cdot \sigma_{\text{SC}}$$

$$= [\dots + 2.382 \times 10^{-12} \text{ m/s}^2 + \dots] \times 0.99374 \times (1 - 10^{-20})$$

BH fractional contribution to g_{total} (estimated):

$$\frac{g_{\text{BH}}}{g_{\text{total}}} \approx \frac{2.382 \times 10^{-12}}{1.238 \times 10^{-8} + 2.382 \times 10^{-12} + \dots} \approx \frac{2.382 \times 10^{-12}}{1.24 \times 10^{-8}} \approx 1.9 \times 10^{-4}$$

While the BH's direct gravitational contribution at the reference radius r is a small fraction of the 26-layer Triadic sum ($\sim 0.019\%$), the r_{SOI} formula reveals that inside $2.36 \times 10^{19} \text{ m}$, BH gravity dominates the reference-point baseline — a qualitative distinction for UQFF predictions of inner galactic structure.

5. Module Implementation

```
// PAPER_279 – SOMBRERO_UQFF_MODULE.cpp, updateCache()
gamma_BH = M_BH / M; // 0.01 = 1%
r_SOI = r * std::sqrt(gamma_BH); // 2.36e20 × 0.1 = 2.36e19 m

// Applied in computeG():
double g_BH = G_grav * M_BH / (r * r); // = gamma_BH * g_base = 2.382e-12 m/s2
```

Computed values for Sombrero M104:

Quantity	Value	Units
M_BH = 109 M_sun	1.989×10 ³⁹	kg
M = 1011 M_sun	1.989×10 ⁴¹	kg
$\gamma_{\text{BH}} = M_{\text{BH}}/M$	0.01	dimensionless
$g_{\text{BH}} = G \cdot M_{\text{BH}}/r^2$	2.382×10 ⁻¹²	m/s ²
$r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$	2.36×10 ¹⁹	m
r_{SOI} in kly	~2.49	kly

6. Key Constants Introduced

Symbol	Value	Units	Description
γ_{BH}	0.01	dimensionless	SMBH Dominance Ratio M_{BH}/M
r_{SOI}	2.36×10 ¹⁹	m	UQFF Sphere of Influence radius
g_{BH}	2.382×10 ⁻¹²	m/s ²	BH direct gravitational contribution at r

7. Physical Significance

- SMBH Dominance Ratio as a universal UQFF parameter:** $\gamma_{\text{BH}} = M_{\text{BH}}/M$ provides a single dimensionless number characterising how BH-dominated a galaxy is. It ranges from ~10⁻⁵ (Sgr A*) to ~0.01 (Sombrero), spanning three orders of magnitude in the current UQFF catalogue.
- UQFF Sphere of Influence formula:** $r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$ is derived directly from setting $g_{\text{BH}}(r) = g_{\text{base}}(r)$. This provides a clean, parameter-free prediction for the BH dominance scale in any UQFF module with a known M_{BH}/M ratio.
- Sombrero as extreme BH-galaxy system:** With γ_{BH} 250× larger than the Milky Way's, Sombrero tests UQFF behaviour in the BH-dominated regime. The large γ_{BH} implies that inner BH effects begin influencing the 26-layer Triadic structure at radii as large as $r_{\text{SOI}} = 2.36 \times 10^{19}$ m.
- AGN feedback implications:** The large M_{BH} implies strong AGN feedback potential. UQFF predicts that AGN activity modifies the vacuum energy density locally, which would appear in the UQFF framework as a time-varying component of $U_{\text{g1_vec}}[i]$ for the innermost layers — a direction for future UQFF module development.
- Generalisation:** The formula $\gamma_{\text{BH}} = M_{\text{BH}}/M$ and $r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$ define a universal UQFF BH dominance prescription applicable to any galaxy module. Future UQFF modules for BH-dominated systems (e.g., NGC 1277, M87) should include this pair of parameters as standard.

8. References

- Ford, H. C. et al. (1996). ApJ, 458, 132. (Sombrero M_{BH} from gas kinematics, HST)
- Kormendy, J. et al. (1996). ApJ, 459, L57. (Sombrero BH mass confirmation)
- Jardel, J. R. et al. (2011). ApJ, 739, 21. (Sombrero DM halo and BH mass)
- Gültekin, K. et al. (2009). ApJ, 698, 198. (M_{BH}–σ correlation)
- PAPER_277: UQFF Gravitational Recession Damping κ_{recession} for z = +0.0063
- PAPER_278: Sombrero Dust Ring UQFF Gravitational Ring Resonator ω_{ring}
- SOMBRERO_UQFF_MODULE.cpp (UQFF 2.0, Session 77)

UQFF 2.0 — The SMBH Dominance Ratio $\gamma_{BH} = M_{BH}/M$ and UQFF Sphere of Influence $r_{SOI} = r \cdot \sqrt{\gamma_{BH}}$ are new universal parameters for UQFF galaxy modules, first derived and tested on the Sombrero Galaxy. — Daniel T. Murphy, Session 77, March 2026.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **magnetar-field** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_B)(\partial^\mu \phi_B) - V(\phi_B) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_B) = \frac{1}{2}m^2\phi_B^2 + \frac{\lambda}{4!}\phi_B^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_B$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_B} = \nabla \times (\rho_{\text{SCm}} \mathbf{v} \times \mathbf{B}) + \kappa B_{\text{crit}} \partial_t \phi_B = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_B = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp! \left(- \exp! \left(- \frac{r - r_0}{\lambda_{\text{VDS}}} \right) \right)$$

For this system, the local VDS sub-ratio is 0.135 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 29, \quad n_{\text{channel}} = 20/26$$

Since $p_{\text{DVP}} = 29$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **103 yr** (field decay quiescence):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos! \left(\frac{2\pi j}{26} \right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh! \left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}} \right) \right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}} / \rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.135	PASS Threshold-consistent
DVP prime	$p_k \in \{2, 3, \dots, 113\}$	$p_{\text{DVP}} = 29$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1 \dots 26, \cos(2\pi j/26)$	PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical

Framework	Canonical Value	This Paper	Status
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	PASS Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_p$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{U_Bi_i}$ unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_Bi_i}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [SSq] \cdot n/26)$
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{SCm} = N_n \cdot \sigma_n^{SCm}(\omega) \cdot \Phi_{\text{phonon}} \cdot (F_{\{U,Bi\}}/F_U - 1)$ where $\sigma_n^{SCm}(\omega, n) = \sigma_0 \cdot \exp[-(\omega - \omega_{SCm})^{2/(2 \cdot \Gamma_{\text{Gamma}} 2)}] \cdot (1 + [SSq] \cdot n/26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: $-f_{26}(q) = \sum_{n=0}^{25} q^{\{n\}2} / (-q;q)_{n^2} - \phi_{26}(q) = \sum_{n=0}^{25} q^{\{n\}2} / (-q^{2;q^2})_n$
 $- \psi_{26}(q) = \sum_{n=1}^{26} q^{\{n\}2} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	9-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq] \exp(-\kappa_i n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} s^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_Scm	0.99	SCm manifold completeness
rho_Scm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_Scm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).