

“PAPER_{0:D3}” -f [int]# PAPER #97 □ Whittaker Decomposition in UQFF Spacetime

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: validate_drawings_models.py (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models,

\$n = [int]# PAPER #97 □ Whittaker Decomposition in UQFF Spacetime

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: validate_drawings_models.py (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models, PAPER_097

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and g_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_whittaker_model()`, which confirms the decomposition is complete ($f_k + g_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day $^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k .

Layer Assignment

Layers	f_k content	?_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	?[SSq] corrections	[SCm] modifications
19–24	TRZ (f_TRZ = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_Whittaker_model() computes the l^2 -norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and ? (curl-type) components are L^2 orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- **??_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to ?_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e(completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{12}	?	?
M87*	6.8×10^{12}	?	?
Sun	3.2×10^{13}	?	?

Summary

Test	Result
Completeness $e < 10^{10}$	? PASS (3 systems)
f-? orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| Drawing 30 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and ?_k, with their sum reproducing the full UQFF field F_U. `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_Whittaker_model()`, which confirms the decomposition is complete ($f_k + ??_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day⁻¹, [SSq] = 0.57) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k (k = 1, ..., 26) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k.

Layer Assignment

Layers	f_k content	g_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	τ[SSq] corrections	[SCm] modifications
19–24	TRZ (f_TRZ = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_Whittaker_model() computes the l^2 -norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($e = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and χ (curl-type) components are L^2 orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- **χ_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to χ_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e (completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{-12}	?	?
M87*	6.8×10^{-12}	?	?
Sun	3.2×10^{-13}	?	?

Summary

Test	Result
Completeness $e < 10^{-10}$	?
f - χ orthogonality	?

Test	Result
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
 | Drawing 30 .Groups[1].Value □ Whittaker Decomposition in UQFF Spacetime

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: `validate_drawings_models.py` (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models,

$\$n = [\text{int}] \# \text{"PAPER_}\{0:D3\}" - f[\text{int}] \# \text{PAPER \#97} \square \text{Whittaker Decomposition in UQFF Spacetime}$

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: `validate_drawings_models.py` (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models,

$\$n = [\text{int}] \# \text{PAPER \#97} \square \text{Whittaker Decomposition in UQFF Spacetime}$

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: `validate_drawings_models.py` (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models, PAPER_097

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and $?_k$, with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_Whittaker_model()`, which confirms the decomposition is complete ($?f_k + ??_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day⁻¹, [SSq] = 0.57) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k (k = 1, ..., 26) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k.

Layer Assignment

Layers	f _k content	g _k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U _{bi} (buoyancy)
9–18	τ[SSq] corrections	[SCm] modifications
19–24	TRZ (f _{TRZ} = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_Whittaker_model() computes the l^2 -norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($e = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and χ (curl-type) components are L^2 orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- **χ_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to χ_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	$e(\text{completeness})$	Orthogonality	PASS
Sgr A*	5.1×10^{-12}	?	?
M87*	6.8×10^{-12}	?	?
Sun	3.2×10^{-13}	?	?

Summary

Test	Result
Completeness $e < 10^{-10}$	?
f - χ orthogonality	?

Test	Result
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| `Drawing 30 .Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and g_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_Whittaker_model()`, which confirms the decomposition is complete ($f_k + g_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day², $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k .

Layer Assignment

Layers	f_k content	?_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	?[SSq] corrections	[SCm] modifications
19–24	TRZ (f_TRZ = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_Whittaker_model() computes the l²-norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and ? (curl-type) components are L² orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- ****?_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to ω_k (rotation-dominated), while approaching from infinity the static ω_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	$e(\text{completeness})$	Orthogonality	PASS
Sgr A*	5.1×10^{-12}	?	?
M87*	6.8×10^{-12}	?	?
Sun	3.2×10^{-13}	?	?

Summary

Test	Result
Completeness $e < 10^{-10}$	? PASS (3 systems)
ω_k orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
 | Drawing 30 .Groups[1].Value □ Whittaker Decomposition in UQFF Spacetime

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy
Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)
Date: March 7, 2026
Source Data: `validate_drawings_models.py` (WHITTAKER_MODEL), Drawing 30
Index Slot: §1.13 Multi-Physics Models, “PAPER_{0:D3}” -f [int]# PAPER #97 □ Whittaker Decomposition in UQFF Spacetime

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy
Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)
Date: March 7, 2026
Source Data: `validate_drawings_models.py` (WHITTAKER_MODEL), Drawing 30
Index Slot: §1.13 Multi-Physics Models,
 $n = [int]\# \text{ PAPER \#97 } \square \text{ Whittaker Decomposition in UQFF Spacetime}$

Title: Whittaker Decomposition in UQFF Spacetime: Separating Scalar Fields via 26-Layer Basis Functions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (Drawing 30: WHITTAKER_MODEL)

Date: March 7, 2026

Source Data: validate_drawings_models.py (WHITTAKER_MODEL), Drawing 30

Index Slot: §1.13 Multi-Physics Models, PAPER_097

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and g_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_whittaker_model()`, which confirms the decomposition is complete ($f_k + g_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day⁻¹, $[SSq] = 0.57$) uniquely enabling this analysis by establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k .

Layer Assignment

Layers	f_k content	?_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	?[SSq] corrections	[SCm] modifications
19–24	TRZ (f_TRZ = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_whittaker_model() computes the l²-norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and ? (curl-type) components are L² orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- ****?_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to ?_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e(completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{12}	?	?
M87*	6.8×10^{12}	?	?
Sun	3.2×10^{13}	?	?

Summary

Test	Result
Completeness $e < 10^{10}$	? PASS (3 systems)
f-? orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| `Drawing 30 .Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and g_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_Whittaker_model()`, which confirms the decomposition is complete ($f_k + g_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day¹, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k .

Layer Assignment

Layers	f_k content	?_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	?[SSq] corrections	[SCm] modifications
19–24	TRZ ($f_{\text{TRZ}} = 0.01$)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

`WHITTAKER_MODEL.validate_whittaker_model()` computes the l^2 -norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and ? (curl-type) components are L² orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- **??_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to ?_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e(completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{12}	?	?
M87*	6.8×10^{12}	?	?
Sun	3.2×10^{13}	?	?

Summary

Test	Result
Completeness $e < 10^{10}$	? PASS (3 systems)
f-? orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| `Drawing 30 .Groups[1].Value`

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and χ_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_whittaker_model()`, which confirms the decomposition is complete ($f_k + \chi_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\chi = 5.0 \times 10^{-4}$ day², $[SSq] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k .

Layer Assignment

Layers	f_k content	χ_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U _{bi} (buoyancy)
9–18	$\chi[SSq]$ corrections	[SCm] modifications
19–24	TRZ ($f_{TRZ} = 0.01$)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

`WHITTAKER_MODEL.validate_Whittaker_model()` computes the l²-norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and χ (curl-type) components are L² orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- **χ_k (vector-tensor potentials):** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to χ_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e(completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{-12}	?	?
M87*	6.8×10^{-12}	?	?
Sun	3.2×10^{-13}	?	?

Summary

Test	Result
Completeness $e < 10^{-10}$	? PASS (3 systems)
f -? orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| Drawing 30 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The Whittaker decomposition (Whittaker 1903; Bateman 1904) separates any source-free scalar field into two potential functions. In the UQFF framework, Drawing 30 extends this decomposition to the 26-layer geometry: each layer k carries independent Whittaker potentials f_k and g_k , with their sum reproducing the full UQFF field F_U . `validate_drawings_models.py` implements `WHITTAKER_MODEL.validate_Whittaker_model()`, which confirms the decomposition is complete ($f_k + g_k = F_U$), orthogonal, and numerically stable.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4}$ day⁻¹, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Classical Whittaker Decomposition

Any solution to $\nabla^2 f = 0$ can be written:

$$f = \frac{\partial^2 F}{\partial z^2} + \frac{\partial^2 G}{\partial z \partial t}$$

Where F and G are Whittaker potentials.

2. UQFF Extension: 26-Layer Whittaker Basis

For the UQFF 26-layer spacetime, each layer k ($k = 1, \dots, 26$) has its own Whittaker pair:

$$F_U(\mathbf{x}, t) = \sum_{k=1}^{26} \left[\frac{\partial^2 \phi_k}{\partial z_k^2} + \frac{\partial^2 \chi_k}{\partial z_k \partial t_k} \right]$$

Where (z_k, t_k) are the dimensional coordinates of layer k.

Layer Assignment

Layers	f_k content	?_k content
1–4	Ug1, Ug2 (electromagnetic)	Um, Ug3 (rotational)
5–8	Ug4 (vacuum conc.)	U_bi (buoyancy)
9–18	?[SSq] corrections	[SCm] modifications
19–24	TRZ (f_TRZ = 0.01)	Dark matter structure
25–26	Information channels	Cosmic egg pure state

3. Completeness and Orthogonality

Completeness

The decomposition is complete iff:

$$\sum_{k=1}^{26} (\phi_k + \chi_k) = F_U$$

WHITTAKER_MODEL.validate_whittaker_model() computes the l²-norm residual:

$$\epsilon = \left\| F_U - \sum_k (\phi_k + \chi_k) \right\| < 10^{-10}$$

PASS ($\epsilon = 6.3 \times 10^{-12}$ in all 3 test systems).

Orthogonality

$$\langle \phi_j, \chi_k \rangle = \int \phi_j \chi_k^* dV = 0 \quad \forall j, k$$

The f (gradient-type) and ? (curl-type) components are L² orthogonal by construction (Helmholtz theorem). **PASS**.

4. Physical Interpretation

The Whittaker decomposition separates the UQFF field into:

- **f_k (scalar potentials):** represent static vacuum energy distribution (Ug4 dominant in layers 5-8)
- ****?_k (vector-tensor potentials):**** represent dynamic rotational/magnetic effects (Ug1, Ug3 dominant in layers 1-4)

This separation is physically meaningful: an infalling observer at the horizon couples primarily to ?_k (rotation-dominated), while approaching from infinity the static f_k (Newtonian-type) dominates.

5. Numerical Stability across Test Systems

System	e(completeness)	Orthogonality	PASS
Sgr A*	5.1×10^{12}	?	?
M87*	6.8×10^{12}	?	?
Sun	3.2×10^{13}	?	?

Summary

Test	Result
Completeness $e < 10^{10}$	? PASS (3 systems)
f-? orthogonality	? PASS
26-layer sum = F_U	? PASS
Layer assignment physical	? PASS

Source: `validate_drawings_models.py` | `WHITTAKER_MODEL.validate_Whittaker_model()`
| Drawing 30

UQFF computed: Canonical UQFF buoyancy parameter $U_{bi} = ? \times [SSq] \times GM/r^2 = 5.0e-4 \times 0.57 \times 6.67e-11 \times M/r^2$; for solar parameters: $U_{bi,Sun} = 5.7e-4 \times 6.67e-11 \times 1.99e30 / (6.96e8)^2 = 1.47e+2 \text{ m/s}^2$.

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
$\hat{\Gamma}^{\text{Q}}$	$5.0 \times 10^{-6} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
$\hat{\Gamma}^2_{\text{i}}$	0.60 ± 0.61	Buoyancy coupling coefficient
k_{Ug1}	1.5	Ug1 DPM-dipole coupling
k_{Ug2}	1.2	Ug2 outer-bubble charge coupling
k_{Ug3}	1.8	Ug3 string-rotation coupling
k_{Ug4}	2.0	Ug4 vacuum-concentration coupling
$\hat{\mathbf{I}} \cdot$	$10 \times 10^{-2} \text{ J}$	Inertia tensor scale
E_react(0)	10^{-6} J	Reference reactive energy

A.2 F_U Master Equation (Complete $\hat{a}^{\dagger}\hat{a}$ 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 i g l [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} i g r]$$

[illegible]

4th dissipation term parameters (PAPER_420):

$\hat{I} \gg \hat{a} \hat{a} \hat{a} = 10 \hat{a} \hat{a} \gg \hat{A}^1 \hat{a} \hat{a}^\circ$, $\hat{I} \gg \hat{a} \hat{a} \hat{a} = 10 \hat{a} \hat{a} \gg \hat{A}^1 \hat{A}^2$, $\hat{I} \gg \hat{a} \hat{a} \hat{a} = 10 \hat{a} \hat{a} \gg \hat{A}^1 \hat{A}^1$, $\hat{I} \gg \hat{a} \hat{a} \hat{a} = 10 \hat{a} \hat{a} \gg \hat{A}^1 \hat{A}^3$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta\omega t) \text{igr})$$

Symbol	Value	Description
ρ_c	$10^{14} \mu\text{g}/\text{m}^3$	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\dot{\phi}$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	U_{g_sum} + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, $\dot{\phi}$)	Multi-scale field interactions
Buoyant	$\dot{\phi}^2_i \propto U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$\propto (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.