

PAPER_199: F_UBii Taxonomy Part 2 — Cosmological and Dark Sector Buoyancy Forces

Version: 1.0

Date: March 13, 2026

Session: 50 — grok_share_7514fe.txt Full Audit

Author: Star-Magic UQFF Research Framework

Source: grok_share_7514fe.txt lines 2766-2900, 6000-6400

$$F_U(r, t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{bi}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [SSq] = 0.57$$

$$\rho_{\Lambda}^{\text{UQFF}} = \rho_{\Lambda}^{\text{obs}} \cdot \left(1 + \kappa^2 \cdot [SSq]^2\right) = \rho_{\Lambda}^{\text{obs}} \times 1.0000000812$$

Abstract

This paper catalogs the second major group of F_UBii variants from the BB_C_Equations_04Sept2026 (177-page equation catalogue): cosmological and dark sector buoyancy forces. Covered are anyon systems, dark energy, inflation, gravitational waves, Loop Quantum Cosmology (LQC) bounce and perturbation spectra, Bekenstein-Hawking entropy, Hawking evaporation lifetime, reheating, Big Bang Nucleosynthesis, reionization, baryon-photon ratio, convective turnover time, CMB angular power spectrum, recombination, NFW dark matter profiles, SIDM core formation, void density evolution, and peculiar velocity. Each F_UBii,X form uses the universal F_rel/E_LEP scaling with a system-specific physical expression F_X.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Dark Sector and Quantum Gravity Variants

1.1 Dark Energy Buoyancy

$$F_{\text{UBii,DE}} = -F_{\text{rel}} \times (\kappa_{\text{DE}} \cdot c^2 / E_{\text{LEP}}) \times Q_{\text{wave}}(z) \times (8\rho_{\text{G}}^{\text{tot}}/3) \times (1+w(z))$$

$$\kappa_{\text{DE}}(a) = \kappa_{\text{DE},0} \cdot \exp(3 \int_1^a (1+w(a'))/a' da')$$

$$w(a) = w_0 + w_a(1-a) \quad (\text{Chevallier-Polarski-Linder parametrization})$$

Source: BB_C_Equations item 721

1.2 Inflation Buoyancy

$$F_{\text{UBii,inf}} = F_{\text{rel}} \times (V(\phi) / E_{\text{LEP}}) \times Q_{\text{wave}} \times 3H^2 \times e^N/(1+e)$$

$V(?)$ = inflaton potential
 N = number of e-folds
 e = slow-roll parameter: $e = (??/H \cdot M_{pl})^2/2$
 Source: BB_C_Equations item 724

1.3 Gravitational Wave Energy Density Buoyancy

$$F_{UBii,GW} = -F_{rel} \times (?_{GW} / E_{LEP}) \times Q_{wave} \times (32pG?^2/c^2) \times e^{-t/t}$$

$?_{GW}$ = gravitational wave energy density
 $?$ = time derivative of strain
 Source: BB_C_Equations item 727

1.4 Anyon System Buoyancy

$$F_{UBii,anyons} = -F_{rel} \times (E_{anyons} / E_{LEP}) \times Q_{wave} \times g(r,t) \times \exp(-d^2_c/(2s^2))$$

E_{anyons} = anyon system energy (2D topological)
 $g(r,t)$ = UQFF gravitational field at position
 Gaussian factor from density fluctuation s
 Source: BB_C_Equations item 710

2. Loop Quantum Cosmology (LQC) Variants

2.1 LQC Perturbation Spectrum Buoyancy

$$F_{UBii,lqcp} = -F_{rel} \times (P(k) ? k^{\{n_s-1\}} \cdot (1+k/k_*)^{\{-a\}} / E_{LEP}) \times Q_{wave}$$

k_* = bounce scale (LQC pre-bounce phase)
 a = UV suppression exponent
 Modifies primordial power spectrum at Planck-scale modes
 Source: BB_C_Equations item 1431, 1254

2.2 LQC Effective Friedmann Buoyancy

$$F_{UBii,lqcf} = F_{rel} \times (H^2 = 8pG?/3 \cdot (1-?/?_{crit}) / E_{LEP}) \times Q_{wave}$$

$?_{crit} = 0.41 \cdot ?_{Planck} \sim 10^{?^3} \text{ g/cm}^3$
 Bounce condition: $H=0$ at $?=?_{crit}$ (avoids singularity)
 Source: BB_C_Equations item 1252, 1604

2.3 LQC Bounce Timescale Buoyancy

$$F_{UBii,bounc} = F_{rel} \times (t_b \sim v(3/(8pG?_{crit})) \sim t_{Pl} \sim 10^{?4^3} \text{ s} / E_{LEP}) \times Q_{wave} \times [H \sim 0 \text{ at bounce, duration} \sim 1/?_{bounce}]$$

Source: BB_C_Equations item 1257, post-1608

3. Black Hole Thermodynamics Variants

3.1 Bekenstein-Hawking Entropy Buoyancy

$$F_{UBii,bhent} = F_{rel} \times (S = k_B \cdot c^3 \cdot A / (4Gh) = 4p \cdot k_B \cdot GM^2 / (hc) / E_{LEP}) \times Q_{wave}$$

$A = 4\pi r_s^2$ (Schwarzschild horizon area)

$r_s = 2GM/c^2$

Holographic: $S \propto A/l_{Pl}^2$ (area law)

Source: BB_C_Equations item 1092, 1248

3.2 Evaporation Lifetime Buoyancy

$$F_{UBii,evapl} = -F_{rel} \times (t_{evap} = 5120\pi G^2 M^3 / (hc^4) / E_{LEP}) \times Q_{wave} \times [P = sAT^4 \sim hc^2/M^2]$$

Power $dM/dt = -P/c^2$ (mass loss rate)

Source: BB_C_Equations item 1601, 1250

4. Cosmological Nucleosynthesis and Reionization Variants

4.1 Big Bang Nucleosynthesis Deuterium Bottleneck

$$F_{UBii,deb} = -F_{rel} \times (t_D = v(3/(32pG\eta_{rad})) \sim 180 \text{ s } (T \sim 0.1 \text{ MeV}) / E_{LEP}) \times Q_{wave}$$

$\eta_{rad} = p^2 kT^4 / (30h^3 c^5) \cdot g_*$ (radiation density)

g_* = effective degrees of freedom (~ 10 at D formation)

Weak freeze-out at $T \sim 1$ MeV; photodissociation until $T < 0.1$ MeV

Source: BB_C_Equations item 1809, 1536

4.2 Baryon-to-Photon Ratio Buoyancy

$$F_{UBii,eta} = F_{rel} \times (\eta = n_b/n_\gamma = 6 \times 10^{-10} / E_{LEP}) \times Q_{wave} \times [\text{Freeze-out: R...}]$$

$n_\gamma = 410 \text{ cm}^{-3}$ (CMB photon density today)

Fit from D, ^3He , ^4He , ^7Li abundances

Source: BB_C_Equations item 1701, 1534

4.3 Reionization Front Buoyancy

$$F_{UBii,reion} = F_{rel} \times (d\eta_e/dt = \eta_\gamma \cdot e_{esc} \cdot f_* - a_B \cdot n_e^2 \cdot C / E_{LEP}) \times Q_{wave}$$

x_e = ionized fraction, $e_{\text{esc}} \sim 0.1-0.3$ (escape fraction)
 C = clumping factor
Source: BB_C_Equations item 1684

5. CMB and Recombination Variants

5.1 CMB Angular Power Spectrum Buoyancy

$$F_{\text{UBii,cmb}} = F_{\text{rel}} \times (C_l = 2/p \cdot \int k^2 dk \cdot P(k) \cdot |\int \delta T(k)|^2 / E_{\text{LEP}}) \times Q_{\text{wave}}$$

$P(k) \propto k^{n_s-4}$ (primordial power, $n_s \sim 0.965$)
Transfer δT : Sachs-Wolfe (large scales) + acoustic (small scales)
Source: BB_C_Equations item 1310, 1080

5.2 Recombination Optical Depth Buoyancy

$$F_{\text{UBii,recomb}} = -F_{\text{rel}} \times (t(z) = \int_{z_{\text{re}}}^z n_e(z') \cdot s_T \cdot c \cdot (dt/dz') dz' / E_{\text{LEP}}) \times Q_{\text{wave}}$$

$z_{\text{re}} \sim 7.7$ (reionization redshift, Planck 2018)
 s_T = Thomson cross-section ($6.65 \times 10^{-29} \text{ m}^2$)
Source: BB_C_Equations item 1313

6. Dark Matter Variants

6.1 NFW Density Profile Buoyancy

$$F_{\text{UBii,nfw}} = -F_{\text{rel}} \times (\rho(r) = \rho_s / ((r/r_s) \cdot (1+r/r_s)^2) / E_{\text{LEP}}) \times Q_{\text{wave}}$$

ρ_s = characteristic density, r_s = scale radius
Universal CDM halo form
Source: BB_C_Equations item 1326

6.2 NFW Rotation Curve Buoyancy

$$F_{\text{UBii,nfwrot}} = F_{\text{rel}} \times (v(r)^2 = 4\pi G \rho_s \cdot r_s^2 \cdot [\ln(1+x) - x/(1+x)] / r / E_{\text{LEP}}) \times Q_{\text{wave}} \times x=r/r_s$$

Flat for $r \gg r_s$
Source: BB_C_Equations item 1535

6.3 SIDM Core Formation Buoyancy

$$F_{\text{UBii,sidm}} = -F_{\text{rel}} \times (G = \rho_s \cdot v/m / E_{\text{LEP}}) \times Q_{\text{wave}} \times \ln(0.02N)$$

$t_{\text{core}} \sim (\rho \cdot s/m)^{-1} \sim 10^{10} \cdot (\rho/108 \text{ M}_{\odot}/\text{kpc}^3)^{-1} \cdot (s/m/1 \text{ cm}^2/\text{g})^{-1} \text{ yr}$
 Core forms when $G \cdot t \sim 1$
 Source: BB_C_Equations item 1249, 1264

7. Void and Peculiar Velocity Variants

7.1 Void Density Evolution Buoyancy

$$F_{\text{UBii,voidden}} = -F_{\text{rel}} \times (d_v(a) = -3/5 \cdot (0_m \cdot a + 0_{\text{?}})^{-3/2} \cdot d_v0 / E_{\text{LEP}}) \times Q_{\text{wave}}$$

a = scale factor, d_v0 = initial underdensity
 Integration: $d \propto a^{-1}$ in matter domination inside void
 Source: BB_C_Equations item 1940, 1338

7.2 Peculiar Velocity Buoyancy

$$F_{\text{UBii,pec}} = F_{\text{rel}} \times (v_{\text{pec}} = -(fH/3) \cdot \int d(r) \cdot r \cdot dr / r^2 / E_{\text{LEP}}) \times Q_{\text{wave}}$$

$f \sim 0_m^{\{0.55\}}$ (growth rate)
 Spherical void: integrate Poisson equation
 Source: BB_C_Equations item 1341

8. Convection and Dynamo Variants

8.1 Convective Turnover Time Buoyancy

$$F_{\text{UBii,conv}} = F_{\text{rel}} \times (t_{\text{conv}} = H_p / v_{\text{conv}} ; v_{\text{conv}} \sim (a^2 g d T \cdot H_p / (4T))^{\{1/3\}} / E_{\text{LEP}}) \times Q_{\text{wave}}$$

H_p = pressure scale height
 $a \sim 2$ (mixing length parameter)
 Source: BB_C_Equations item 1533

8.2 Magnetic Field Reversal Buoyancy (Dynamo Parity)

$$F_{\text{UBii,rev}} = F_{\text{rel}} \times (l_{\text{rev}} = (a_{\text{dynamo}} \cdot \eta)^{\{1/2\}} \cdot l_{\text{force}} / E_{\text{LEP}}) \times Q_{\text{wave}}$$

a_{dynamo} = dynamo a -coefficient ($\sim v/3$)
 η = magnetic resistivity
 Growth vs diffusion sets reversal scale
 Source: BB_C_Equations item 1863, 1238

8.3 Kazantsev Dynamo Buoyancy

$$F_{\text{UBii,dyn}} = F_{\text{rel}} \times (dE_M/dt = (3/2) \cdot E_M / t_{\text{eddy}} \cdot (Re_m^{\{1/2\}} - 1) / E_{\text{LEP}}) \times Q_{\text{wave}}$$

$Re_m \sim 10^{10}$ (magnetic Reynolds number in galaxy clusters)
 $t_{eddy} = l/v \sim \text{Myr}$ (eddy turnover for $l \sim \text{kpc}$)
 Source: BB_C_Equations item 1234, 1411

9. Star Formation and Feedback Variants

9.1 Metal Enrichment Buoyancy

$$F_{UBii,metal} = F_{rel} \times (Z = Y \cdot SFR / \dot{m}_{gas} - Z \cdot \dot{m}_{out} / M_{gas} / E_{LEP}) \times Q_{wave} \times (Z \sim 0.1)$$

$Y \sim 0.02$ (stellar yield)
 Steady state: $Z = Y \cdot SFR / \dot{m}_{out}$
 Source: BB_C_Equations item 1395, 1571

9.2 Photoevaporation (Exoplanet) Buoyancy

$$F_{UBii,photo} = F_{rel} \times (\dot{m}_{evap} = e \cdot L_X \cdot R_p^3 / (GM_p^2 \cdot K(?)) / E_{LEP}) \times Q_{wave}$$

$L_X \sim 10^{27} \text{ erg/s}$ (host star X-ray luminosity)
 $K(?)$ = penetration correction factor
 Source: BB_C_Equations item 1496, 1490

10. References

- grok_share_7514fe.txt lines 2766–2900, 6000–6380 (BB_C_Equations_04Sept2025.pdf Part 2 catalogue)
- PAPER_198: F_UBii Taxonomy Part 1 (Compact Objects)
- PAPER_196: Triadic Master Equation System