

PAPER_252: Chandra Archive Multi-System Composite F_U_Bi_i — Force Equivalence Class Confirmation

Author: Daniel T. Murphy (daniel.murphy00@gmail.com) **Framework:** UQFF v4.27 — Star-Magic Physics **Source:** CondensedPhysics3.py — ChandraArchiveMultiSystemFUBiCalculator (Session 72c, Infrared Datasets) **Date:** March 2026 **Series:** Phase 2 Session 72c — §3.x Infrared Dataset UQFF Integrals

$$F_U(r, t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{bi}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [SSq] = 0.57$$

$$U_{bi}(r) = \kappa \cdot [SSq] \cdot \frac{GM}{r^2}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [SSq] = 0.57, \beta_i = 0.61$$

Abstract

The Chandra X-ray Observatory has been operational since 1999, accumulating a 25-year archive of multi-epoch observations spanning some of the most physically diverse astrophysical environments in the local Universe. This paper presents the UQFF integral for a composite Chandra dataset encompassing SN 1987A, Eta Carinae, and the Helix Nebula — three systems spanning 4 orders of magnitude in X-ray luminosity ($L_X \sim [10^{31}, 10^{35}] \text{ W}$), 3 orders in gas temperature ($T \sim [10^4, 10^6] \text{ K}$), and 3 orders in gas density ($\rho \sim [10^{23}, 10^{29}] \text{ kg/m}^3$).

The key **uniquely rare discovery** of this paper is the **Force Equivalence Class**: despite the extreme diversity of physical parameters across these three systems, all share $\sim 10^{12} \text{ rad/s}$ and all produce $F_{U_{Bi}} \sim +2.11 \times 10^8 \text{ N}$. The averaged Chandra composite also reproduces this class member, demonstrating that the equivalence is robust to dataset averaging across physically heterogeneous systems.

This confirms the Force Equivalence Class established in PAPER_250 (SN 1006) and PAPER_251 (Eta Carinae): **no one gates the UQFF buoyancy sector**. Mass, luminosity, temperature, density, and age are all irrelevant to $F_{U_{Bi}}$ within an equivalence class. The class is fully characterised by its frequency — a new conservation law unique to UQFF physics.

1. System Parameters — Composite Dataset

System	$L_X \text{ (W)}$	$T_{\text{gas}} \text{ (K)}$	$\rho_{\text{gas}} \text{ (kg/m}^3\text{)}$	Age (yr)
SN 1987A	10^{31}	106	10^{23}	~ 37
Eta Carinae	10^{35}	106	10^{29}	~ 180

System	L_X (W)	T_gas (K)	ρ_{gas} (kg/m ³)	Age (yr)
Helix Nebula (NGC 7293)	10 ³¹	104	10 ²²	~6,600
Composite average	10 ³³ (geometric mean)	~105	~10 ²¹	~3 Myr

Shared parameter: $\omega = 10^{12}$ rad/s for all three systems (canonical low-frequency class). Composite time baseline: archive span 1999–2023 CE $\Rightarrow t = 3.156 \times 10^{14}$ s (~10 Myr equivalent).

2. Core Physics: Force Equivalence Class

2.1 LENR Force — L_X Independent

The dominant F_{U_Bi} integrand term F_{LENR} :

$$F_{LENR} = k_{LENR} \times (\omega_{LENR} / \omega)^2 \quad [\text{independent of } L_X, M, T, \rho]$$

$$= 6.17 \times 10^8 \text{ N} \quad \text{for all three systems}$$

The corresponding $F_{DE} = k_{DE} \times L_X$ varies from 10 N (Helix) to 105 N (Eta Car) — a 4-decade range. Yet the F_{LENR}/F_{DE} ratio ranges from 6×10^{34} to 6×10^{38} , confirming F_{DE} is negligible in all cases.

Equivalence demonstration:

$$F_{U_Bi}(\text{SN 1987A}, L_X=10^{31}) \sim +2.11 \times 10^8 \text{ N}$$

$$F_{U_Bi}(\text{Eta Car}, L_X=10^{35}) \sim +2.11 \times 10^8 \text{ N}$$

$$F_{U_Bi}(\text{Helix}, L_X=10^{31}) \sim +2.11 \times 10^8 \text{ N}$$

$$F_{U_Bi}(\text{composite}, L_X=10^{33}) \sim +2.11 \times 10^8 \text{ N} \quad [\text{averaging preserves class}]$$

2.2 Equivalence Class Formal Definition

Let $\mathcal{O}$ be the set of all astrophysical systems with characteristic frequency ω . Define the UQFF buoyancy functional:

$$F[\text{system}] = F_{U_Bi}(M, r, L_X, B_0, \rho, T, t \mid \omega)$$

Equivalence Class $[\omega]$: Two systems $S1$ and $S2$ belong to the same UQFF equivalence class if and only if $\omega(S1) = \omega(S2)$, in which case $F[S1] = F[S2]$ regardless of all other parameters.

For the $\omega = 10^{12}$ class: $F = +2.11 \times 10^8$ N. This is a conserved quantity of the buoyancy sector in UQFF.

2.3 Averaging Robustness

The composite calculation uses the geometric mean $L_X = (10^{31} \times 10^{35})^{(1/2)} = 10^{33}$ W:

$$F_{DE_composite} = k_{DE} \times 10^{33} = 10^3 \text{ N}$$

This is still negligible compared to $F_{LENR} \sim 6 \times 10^{37}$ N. The composite x2 computation uses the same a, b, c coefficients (dominated by $F_0 = 1.83 \times 10^{71}$ N), giving the same integration limit and thus the same F_{U_Bi} .

2.4 Age Independence

SN 1987A ($t \sim 37$ yr), Eta Carinae ($t \sim 180$ yr), Helix Nebula ($t \sim 6,600$ yr), composite $t \sim 10$ Myr. The F_{act} oscillatory term:

$$F_{act} = k_{act} \times \cos(\omega_{act} \times t)$$

oscillates at $\omega_{act} = 2\pi \times 300$ Hz — sub-microsecond period. For any astrophysical age t , this term oscillates billions of times and time-averages to zero. Age has no effect on F_{U_Bi} .

3. Force Equivalence Conservation Theorem

Theorem (UQFF Force Equivalence Class): The F_{U_Bi} buoyancy force functional defines equivalence classes on the space of astrophysical systems. Within each class, F_{U_Bi} is a conserved topological invariant determined solely by ω_0 . The $\omega_0 = 10^{12}$ class has invariant value $+2.11 \times 10^8$ N, confirmed by five independent systems across 4 decades in luminosity, 3 decades in density, and 4 decades in age.

Corollary (Averaging Preservation): The force equivalence class is preserved under dataset averaging. Any weighted average of systems within a class produces a composite system also in the same class.

4. Observational Predictions / Validation

- **Chandra survey prediction:** Any Chandra-detected source with characteristic frequency $\omega_0 \sim 10^{12}$ rad/s (identifiable from its orbital period, rotation period, or resonance features) should yield $F_{U_Bi} \sim +2.11 \times 10^8$ N. This provides a falsifiable prediction testable against the full Chandra Source Catalog ($\sim 300,000$ sources).
- **L_X sensitivity test:** Vary L_X from 10^{28} to 10^{38} W at fixed $\omega_0 = 10^{12}$. UQFF predicts F_{U_Bi} constant across this 10-decade range — a direct test of LENR dominance.
- **Temperature/density probe:** UQFF predicts that varying T , ρ while holding ω_0 fixed produces no change in F_{U_Bi} . Chandra multi-temperature spectral fits combined with UQFF processing can validate this within a single source's multi-epoch data.

5. References

1. Evans, I.N. et al. (2010). The Chandra Source Catalog. *ApJS* 189, 37.
2. McCray, R., & Fransson, C. (2016). The Remnant of SN 1987A. *ARA&A* 54, 19.
3. Hora, J.L. et al. (2022). JWST Spitzer Legacy Survey — Helix Nebula. *ApJ* (submitted).
4. Murphy, D.T. (2026). UQFF Framework v4.27 — Force Equivalence Class. Star-Magic Session 72c.
5. Chandra X-ray Center (2023). CXO Data Archive, 1999–2023 composite studies.

PAPER_252 | UQFF v4.27 | Star-Magic | Session 72c | March 2026