

PAPER_255: PSR J0030+0451 Isolated Neutron Star — Density Regime Positive Buoyancy and F_neutron Dominance

Author: Daniel T. Murphy (daniel.murphy00@gmail.com) **Framework:** UQFF v4.27 — Star-Magic Physics **Source:** CondensedPhysics3.py — PSRJ0030NeutronStarFUBiCalculator (Session 72d, ALMA Cycle 12) **Date:** March 2026 **Series:** Phase 2 Session 72d — §3.x ALMA Cycle 12 Neutron Star UQFF Integrals

$$F_U(r, t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{bi}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, \quad [SSq] = 0.57$$

$$M_J^{\text{UQFF}} = M_J^{\text{Jeans}} \cdot \left(1 - [SSq] \cdot \frac{B^2}{8\pi\rho c_s^2} \right), \quad [SSq] = 0.57$$

Abstract

PSR J0030+0451 is an isolated millisecond pulsar at $\sim 1,100$ light-years, with a mass of approximately $1.4 M_{\text{sun}}$ confined to a radius of ~ 10 km ($r = 104$ m) — the compact geometry of a neutron star. This system is the **first isolated pulsar class** in the CP3 calculator, and it introduces a new UQFF regime defined by the neutron-star-density cross-section parameter $s_n \sim 10^3?$, representing the degenerate nuclear density of neutron star matter.

In the ISM systems of PAPER_250-254, $s_n \sim 10^4$ yields $F_{\text{neutron}} = k_{\text{neutron}} \times s_n = 106$ N. For PSR J0030 at $s_n = 10^3?$, $F_{\text{neutron}} = 10^{10} \times 10^3? = \mathbf{104? N}$ — a difference of 53 orders of magnitude. This neutron force is the dominant UQFF term by far.

The key **uniquely rare discovery** of this paper is that despite this 53-order amplification of F_{neutron} , and despite the compact scale ($r = 104$ m vs $r = 6.17 \times 10^{16}$ m for the SNRs), PSR J0030 is a **positive buoyancy** system: $F_{U_{\text{Bi}}} \sim +2.53 \times 10^{208}$ N. The compact-scale geometry at $\phi_0 = 10^{?12}$ preserves the positive sign. The equivalence class extends across 14 orders of magnitude in radius and 53 orders in s_n .

1. System Parameters

Parameter	Symbol	Value	Units	Source
Distance	d	$\sim 1,100$	ly	Chandra/NICER 2019
Mass	M	$1.4 M_{\text{sun}} = 2.786 \times 10^{30}$	kg	Neutron star canonical

Parameter	Symbol	Value	Units	Source
Neutron star radius	r	104	m	~10 km
X-ray luminosity	L_X	10 ³¹	W	NICER 2019
Surface B field	B0	108	T	Millisecond pulsar typical
System frequency	?0	10 ^{?12}	rad/s	Same as SNR class
Neutron cross-section	s_n	10^{3?}	—	NS density (vs ISM 10²⁴)

2. Core Physics: Neutron-Star Density Regime

2.1 F_neutron — The New Dominant Term

$$F_{\text{neutron}} = k_{\text{neutron}} \times s_n = 10^{1^0} \times 10^{3?} = 104? \text{ N}$$

For comparison:

$$F_{\text{LENR}} (?0=10^{?12}) \sim 6.17 \times 10^{3?} \text{ N}$$

$$F_{\text{neutron}} / F_{\text{LENR}} = 104? / 6.17 \times 10^{3?} \sim 1.6 \times 10?$$

F_neutron exceeds F_LEN R by 9 orders for the neutron star regime. The force hierarchy shifts from LENR-dominant (ISM and SNR systems) to **neutron-dominant** (compact objects):

Force hierarchy at ?0=10^{?12}, s_n=10^{3?}:

$$F_{\text{neutron}} \sim 104? \text{ N} \quad [\text{dominant} - 9 \text{ orders above } F_{\text{LENR}}]$$

$$F_{\text{LENR}} \sim 6 \times 10^{3?} \text{ N} \quad [\text{second}]$$

$$F_{\text{Newt}} \sim GM/r^2 \cdot |x_2| \quad [\text{negligible}]$$

$$F_{\text{res}} \ll F_{\text{LENR}} \quad [\text{DPM invisible} - \text{same conclusion as PAPER}_{251}]$$

2.2 Compact Geometry and P Positive Buoyancy Preservation

Despite the 9-order dominance of F_neutron over F_LEN R, the sign of F_U_Bi remains positive. This is because the compact geometry (r = 104 m) affects the term_gravity = GM/r² and the integration limit x2 in a way that preserves the positive root:

$$\begin{aligned} \text{term_gravity} &= G \cdot M / r^2 = 6.674 \times 10^{-11} \times 2.786 \times 10^{30} / (104)^2 \\ &\sim 1.86 \times 10^6 \text{ m/s}^2 \quad [\text{huge surface gravity}] \end{aligned}$$

The quadratic discriminant $b^2 - 4ac$ with $a = 1.86 \times 10^6$, $b = 4.72 \times 10^{?3}$, $c \sim -1.83 \times 10^{71}$ gives a positive x_2 root (same sign as ISM systems), because the vacuum energy $F_0 = 1.83 \times 10^{71} \text{ N}$ overwhelms the sign-determining coefficient c regardless of the surface gravity scale.

Key result: $x_2 > 0$? $F_{\text{U_Bi_i}} = \text{integrand} \times |x_2| > 0$? **positive buoyancy at $F_{\text{U_Bi}} \sim +2.53 \times 10^{2^8} \text{ N}$.**

2.3 53-Order s_n Range: Equivalence Class Breadth

The s_n parameter spans:

s_n (ISM/SNR systems): $\sim 10^{?4}$? $F_{\text{neutron}} = 10^6$ N [PAPER_250–254]
 s_n (PSR J0030): $\sim 10^{3?}$? $F_{\text{neutron}} = 10^{4?}$ N [this paper]

53 orders of magnitude in s_n , yet both classes show **positive buoyancy at $\Omega = 10^{?12}$** . This confirms that s_n (like B_0 in PAPER_251) does not breach the equivalence class — the Ω parameter remains the exclusive class determinant.

2.4 DPM Resonance at $B_0 = 108$ T

$$\begin{aligned} \text{DPM}_{\text{resonance}} (\text{PSR J0030}) &= 2 \cdot \mu_B \cdot B_0 / (\hbar \cdot \Omega) \\ &= 2 \times 9.274\text{e-}24 \times 108 / (1.0546\text{e-}34 \times 10^{?12}) \\ &\sim 1.76 \times 10^{31} \end{aligned}$$

This is an astronomically large DPM resonance, yet it is still invisible relative to F_{neutron} : $F_{\text{res}}/F_{\text{neutron}} \ll 1$. The DPM invisibility theorem (PAPER_251) extends to the neutron-star-density regime.

3. Extended Force Equivalence Class Theorem

Theorem (UQFF NS-Density Class Extension): The positive buoyancy equivalence class at $\Omega = 10^{?12}$ rad/s includes compact objects with neutron-star densities ($s_n \sim 10^{3?}$) in addition to diffuse ISM systems ($s_n \sim 10^{?4}$). $F_{\text{U_Bi}} \sim +2 \times 10^{2^8}$ N regardless of s_n spanning 53 orders, confirming that s_n is not a class-determinant parameter. The vacuum energy anchor $F_0 = 1.83 \times 10^{71}$ N ensures $x_2 > 0$ for all physically observable values of s_n .

4. ALMA Cycle 12 Observational Context

PSR J0030+0451 is an ALMA Cycle 12 proposal target. Observable UQFF signatures include:

- **Isotopic anomaly:** LENR neutron-capture at $F_{\text{neutron}} = 10^{4?}$ N (53 orders above ISM) predicts elevated $^2\text{H}/^1\text{H} > 10^{?5}$ and $^{13}\text{C}/^{12}\text{C} > 0.01$ in the pulsar wind nebula — detectable with ALMA Band 6 at 230 GHz.
- **EHT polarimetry:** The extreme $\text{DPM}_{\text{resonance}} \sim 1.76 \times 10^{31}$ at $B_0 = 108$ T predicts distinctive helical B-field structure in the pulsar wind, detectable with EHT 20 μas resolution at 230 GHz.
- **NICER hotspot:** PSR J0030+0451 hotspot morphology constrains the NS mass-radius relation; UQFF predicts $F_{\text{U_Bi}}$ positive — consistent with a gravitationally stable bound NS (no anomalous mass loss or unbinding).

5. References

1. Riley, T.E. et al. (2019). A NICER View of PSR J0030+0451: Millisecond Pulsar Parameter Estimation. *ApJ Lett.* 887, L21.
2. Özel, F., & Freire, P. (2016). Masses, Radii, and the Equation of State of Neutron Stars. *ARA&A* 54, 401.
3. Kozima, H. (1998). *The Science of the Cold Fusion Phenomenon*. Elsevier.
4. ALMA Proposal Cycle 12. PSR J0030+0451 — UQFF isotopic anomaly detection (Murphy, D.T. 2026).
5. Murphy, D.T. (2026). UQFF Framework v4.27 — NS Density Regime Discovery. Star-Magic Session 72d.

PAPER_255 | UQFF v4.27 | Star-Magic | Session 72d | March 2026