

M51 Whirlpool and NGC 1316 Fornax A – MUGE with Simulation Scripts

Daniel T. Murphy

2025-01-01

PAPER_750: M51 Whirlpool and NGC 1316 Fornax A — MUGE with Simulation Scripts

Author: Daniel T. Murphy

Framework: Universal Quantum Field Superconductive Framework (UQFF)

Session: 180 continuation | v5.38

Date: 2025

CP4 Class: #334 — M51NGC1316MUGESimulationCalculator

Abstract

Two Hubble-studied galaxies — M51 (the Whirlpool Galaxy, interacting with NGC 5195) and NGC 1316 (Fornax A, a post-merger elliptical) — represent opposite ends of the galaxy evolution spectrum. This paper derives tailored Master Universal Gravity Equations (MUGEs) for both systems using Hubble ACS datasets, incorporating interaction-specific environmental terms: F_{tidal} and ψ_{spiral} for M51, and $F_{\text{tidal}} + F_{\text{cluster}} + \rho_{\text{dust}}$ for NGC 1316. Full Python simulation scripts (m51_simulation.py, ngc1316_simulation.py) are included for radial acceleration profile generation.

1. Introduction

1.1 M51 (Whirlpool Galaxy)

M51 is a grand design spiral at 7.7 Mpc with an active tidal interaction companion NGC 5195. Hubble ACS observations (2005) provide: - $M_{\text{visible}} = 1.2 \times 10^{11} \text{ MM}_{\text{sun}}$ - $M_{\text{DM}} = 4 \times 10^{10} \text{ MM}_{\text{sun}}$
- $\text{SFR} \approx 1 \text{ MM}_{\text{sun}}/\text{yr}$ - $M_{\text{BH}} \approx 10^6 \text{ MM}_{\text{sun}}$ - Distance = 7.7 Mpc, $z \approx 0.0015$ - $B \approx 5 \times 10^{-6} \text{ T}$

1.2 NGC 1316 (Fornax A)

NGC 1316 is a giant elliptical at 75 Mpc with a complex merger history: - $M_{\text{total}} \approx 5 \times 10^{11} \text{ MM}_{\text{sun}}$ (visible + DM) - $M_{\text{DM}} = 1.5 \times 10^{11} \text{ MM}_{\text{sun}}$ - $M_{\text{BH}} \approx 10^8 \text{ MM}_{\text{sun}}$ - Merger

age: 1-3 Gyr ago (from ripples, dust lanes) - $\rho_{\text{dust}} \approx 10^{-21} \text{ kg/m}^3$

2. M51 Whirlpool Galaxy MUGE

$$\begin{aligned}
g_{M51}(r, t) = & (G * M(t)) / (r(t)^2) * (1 + H(t, z)) * (1 - B(t) / B_{crit}) * (1 + F_{env}(t)) \\
& + (U_g1 + U_g2 + U_g3' + U_g4) + U_i \\
& + (\Lambda * c^2 / 3) \\
& + (\hbar / \sqrt{(\Delta x * \Delta t)}) * \int (\psi_{total} * H * \psi_{total} dV) * (2\pi / t_{Hubble}) \\
& + \rho_{fluid} * V * g \\
& + (M_{vis} + M_D M) * (\Delta \rho / \rho + (3 * G * M) / r^3)
\end{aligned}$$

Canonical note: The $(G * M(t)) / (r(t)^2)$ term is the Step 10 Newton observational projection — per UQFF chain, the foundational gravity seed is DPM/Ug1 ($k1\mu \text{ M/r}$). The U_{g_i} terms in the equation are the canonical DPM-derived components.

M51 F_{env} Terms

$$\begin{aligned}
F_{env}(t)_{M51} &= F_{tidal} + \psi_{spiral_correction} \\
F_{tidal} &= G * M_{NGC5195} / d_{interaction}^2 \\
M_{NGC5195} &= 10^{10} M_{sun} = 1.989 \times 10^{40} \text{ kg} \\
d_{interaction} &= 50 \text{ kpc} = 1.54 \times 10^{21} \text{ m} \\
F_{tidal} &= 6.674 \times 10^{-11} \times 1.989 \times 10^{40} / (1.54 \times 10^{21})^2 \\
F_{tidal} &= 5.59 \times 10^{-13} \text{ m/s}^2 (\text{normalized to } F_{env} \text{ fraction}) \\
\psi_{spiral} &= A * e^{(-r^2 / (2\sigma^2))} * e^{i(m * \theta - \omega * t)} \\
\sigma &= 1 \text{ kpc} = 3.086 \times 10^{19} \text{ m} (\text{spiral arm width}) \\
m &= 2 (\text{grand design } 2 - \text{arm spiral})
\end{aligned}$$

M51 H(t,z)

$$\begin{aligned}
H(t, z) &= H_0 * \sqrt{(0.3 * (1 + 0.0015)^3 + 0.7)} = H_0 * \sqrt{0.7003} = H_0 \\
z &= 0.0015 (\text{negligible correction})
\end{aligned}$$

M51 Parameters Summary

Parameter	Value
M _{visible}	$1.2 \times 10^{11} M_{sun} = 2.39 \times 10^{41} \text{ kg}$
M _{DM}	$4 \times 10^{10} M_{sun} = 7.96 \times 10^{40} \text{ kg}$
r _{galaxy}	30 kpc = $9.26 \times 10^{20} \text{ m}$
B	$5 \times 10^{-6} \text{ T}$
SFR	$1 M_{sun}/\text{yr} = 6.3 \times 10^{22} \text{ kg/yr}$
F _{tidal}	$5.59 \times 10^{-13} \text{ m/s}^2$

3. M51 Simulation Script (m51_simulation.py)

```
import numpy as np
import matplotlib.pyplot as plt

# Constants
G = 6.6743e-11          # m^3 kg^-1 s^-2
H_0 = 70e3 / 3.086e22  # s^-1
Lambda = 1.1e-52        # m^-2
c = 3e8                 # m/s
hbar = 1.0546e-34       # J.s
t_Hubble = 4.35e17      # s
B_crit = 1e11           # T
rho_{vac\_SCm} = 7.09e-37 # J/m^3
rho_{vac\_UA} = 7.09e-36 # J/m^3
mu_0 = 4 * np.pi * 1e-7 # H/m
M_sun = 1.989e30         # kg
kpc = 3.086e19          # m

# M51 Parameters
M_vis = 1.2e11 * M_sun
M_DM = 4.0e10 * M_sun
M_NGC5195 = 1e10 * M_sun
r_0 = 30e3 * kpc
d_inter = 50e3 * kpc
B = 5e-6                # T
sigma_spiral = 1e3 * kpc
z = 0.0015
lambda_I = 1.0
F_RZ = 0.01
omega_i = 1.585e-8      # rad/s

def M_total():
    return M_vis + M_DM

def H_z(z):
    return H_0 * np.sqrt(0.3 * (1 + z)**3 + 0.7)

def F_tidal():
    return G * M_NGC5195 / d_inter**2

def U_i():
    return lambda_I * (rho_{vac\_SCm} / rho_{vac\_UA}) * omega_i * 1.0 * (1 + F_RZ)

def psi_spiral(r):
    A = 1e-10
    return A * np.exp(-r**2 / (2 * sigma_spiral**2))
```

```

def g_M51(r, t=0):
    M = M_total()
    H = H_z(z)
    F_env = F_tidal() / (G * M / r**2) # normalized
    g_grav = (G * M / r**2) * (1 + H) * (1 - B / B_crit) * (1 + F_env)
    quantum = hbar / np.sqrt(1e-10 * 1e-20) * psi_spiral(r)**2 * (2 * np.pi / t_Hubble)
    dm_term = (M_vis + M_DM) * (1e-5 + 3 * G * M / r**3)
    cosmological = Lambda * c**2 / 3
    return g_grav + U_i() + cosmological + quantum + dm_term

# Compute radial profile
r_vals = np.linspace(1e3 * kpc, 30e3 * kpc, 200)
g_vals = [g_M51(r) for r in r_vals]

plt.figure(figsize=(10, 6))
plt.plot(r_vals / kpc, g_vals, 'b-', linewidth=2, label='M51 MUGE Acceleration')
plt.xlabel('Radius (kpc)')
plt.ylabel('g_M51 (m/s^2)')
plt.title('M51 Whirlpool Galaxy - UQFF MUGE Radial Profile')
plt.legend()
plt.grid(True, alpha=0.3)
plt.savefig('m51_{gravity\_profile}.png', dpi=150)
plt.close()
print("M51 simulation complete. Profile saved to m51_{gravity\_profile}.png")

```

4. NGC 1316 (Fornax A) MUGE

$$\begin{aligned}
g_{\text{NGC1316}}(r, t) = & (G \cdot M(t)) / (r(t)^2) * (1 + H(t, z)) * (1 - B(t) / B_{\text{crit}}) * (1 + F_{\text{env}}(t)) \\
& + (U_{g1} + U_{g2} + U_{g3} + U_{g4}) + U_i \\
& + (\Lambda \cdot c^2 / 3) \\
& + (\hbar / \sqrt{\Delta x \cdot \Delta t}) * \int (\psi_{\text{total}} \cdot H \cdot \psi_{\text{total}} dV) * (2\pi / t_{\text{Hubble}}) \\
& + \rho_{\text{dust}} \cdot V \cdot g \\
& + (M_{\text{vis}} + M_{\text{DM}}) * (\Delta \rho / \rho + (3 \cdot G \cdot M) / r^3)
\end{aligned}$$

NGC 1316 F_{env} Terms

$F_{\text{env}}(t)_{\text{NGC1316}} = F_{\text{tidal}} + F_{\text{cluster}}$

F_{tidal} : tidal from past spiral galaxy mergers
 $F_{\text{tidal}} = G \cdot M_{\text{spiral}} / d_{\text{spiral}}^2$
 $M_{\text{spiral}} \sim 10^{10} \text{ } M_{\text{sun}}$ (remnant progenitor)
 $d_{\text{spiral}} \sim 50 \text{ kpc}$
 $F_{\text{tidal}} \sim 5.59 \times 10^{-13} \text{ m/s}^2$

F_{cluster} : star cluster tidal disruption
 $F_{\text{cluster}} = k_{\text{cluster}} \cdot M_{\text{cluster}}$
 $k_{\text{cluster}} \sim 10^{-1} \text{ } N / M_{\text{sun}}$

```

M_cluster ~= 10^6 M_sun
F_cluster ~= 10^{-6} m/s^2

rho_dust: dust lane fluid term
rho_dust = 10^{-21} kg/m^3 (ACS-derived dust column density)

```

NGC 1316 Mass Evolution

```

M(t) = M_vis + M_DM + M_merger(t)
M_merger(t) = M_spiral * e^(-t/tau_merge)
tau_merge ~= 1 Gyr = 3.156x10^{16} s

```

5. NGC 1316 Simulation Script (ngc1316_simulation.py)

```

import numpy as np
import matplotlib.pyplot as plt

# Constants (same as M51)
G = 6.6743e-11
H_0 = 70e3 / 3.086e22
Lambda = 1.1e-52
c = 3e8
hbar = 1.0546e-34
t_Hubble = 4.35e17
B_crit = 1e11
rho_{vac\_{SCm}} = 7.09e-37
mu_0 = 4 * np.pi * 1e-7
M_sun = 1.989e30
kpc = 3.086e19

# NGC 1316 Parameters
M_vis = 3.5e11 * M_sun
M_DM = 1.5e11 * M_sun
M_spiral = 1e10 * M_sun
M_BH = 1e8 * M_sun
tau = 3.156e16 # 1 Gyr merger decay
d_spiral = 50e3 * kpc
rho_dust = 1e-21 # kg/m^3
B = 1e-4 # T
z = 0.005 # NGC 1316 redshift
k_cluster = 1e-12 # N/M_sun
M_cluster = 1e6 * M_sun
sigma_dust = 2e3 * kpc

def M_total(t):
    return M_vis + M_DM + M_spiral * np.exp(-t / tau)

```

```

def H_z(z):
    return H_0 * np.sqrt(0.3 * (1 + z)**3 + 0.7)

def F_env(t):
    F_tidal = G * M_spiral / d_spiral**2
    F_cluster = k_cluster * M_cluster
    return F_tidal + F_cluster

def U_i():
    rho_{vac\_UA} = 7.09e-36
    return 1.0 * (rho_{vac\_SCm} / rho_{vac\_UA}) * 1e-8 * 1.0 * 1.01

def g_NGC1316(r, t=3.156e15): # t=100 Myr default
    M = M_total(t)
    H = H_z(z)
    F = F_env(t) / (G * M / r**2)
    g_grav = (G * M / r**2) * (1 + H) * (1 - B / B_crit) * (1 + F)
    g_dust = rho_dust * (4/3 * np.pi * r**3) * G * M / r**2
    dm_term = (M_vis + M_DM) * (1e-5 + 3 * G * M / r**3)
    cosmological = Lambda * c**2 / 3
    return g_grav + U_i() + cosmological + g_dust + dm_term

# Compute
r_vals = np.linspace(1e3 * kpc, 100e3 * kpc, 100)
g_vals = [g_NGC1316(r) for r in r_vals]

plt.figure(figsize=(10, 6))
plt.plot(r_vals / kpc, g_vals, 'r-', linewidth=2, label='NGC 1316 MUGE Acceleration')
plt.xlabel('Radius (kpc)')
plt.ylabel('g_NGC1316 (m/s^2)')
plt.title('NGC 1316 Fornax A -- UQFF MUGE Radial Profile')
plt.legend()
plt.grid(True, alpha=0.3)
plt.savefig('ngc1316_{gravity\_profile}.png', dpi=150)
plt.close()
print("NGC 1316 simulation complete. Profile saved to ngc1316_{gravity\_profile}.png")

```

6. Comparative Analysis

Property	M51 (Whirlpool)	NGC 1316 (Fornax A)
Type	Sc spiral	E/S0 elliptical
Distance	7.7 Mpc	75 Mpc
Interaction	Active (NGC 5195)	Past mergers (~1-3 Gyr)
Key term	F_{tidal} (live)	$F_{\text{cluster}} + \rho_{\text{dust}}$
Star formation	$\text{SFR} \approx 1 \text{ MM}_{\text{sun}}/\text{yr}$	~0 (quenched)

Property	M51 (Whirlpool)	NGC 1316 (Fornax A)
AGN activity	Moderate	Strong (radio lobes)
Key physics	Density waves (ψ_{spiral})	Dust lane drag (ρ_{dust})

7. Astrophysical Significance

M51: The F_{tidal} term drives the prominent spiral arms visible in Hubble observations. The ψ_{spiral} density wave modulation affects star formation efficiency across the disk. NGC 5195's tidal force creates the bridge structure and enhanced SFR in the outer arms.

NGC 1316: The F_{cluster} term explains the deficit of low-mass globular clusters in the inner region (disrupted by tidal forces during multiple mergers). The ρ_{dust} dust lane creates a complex hydrodynamic environment near the AGN.

8. Conclusion

The M51 and NGC 1316 MUGEs demonstrate the UQFF framework's ability to model both active interaction (M51) and post-merger (NGC 1316) galaxy evolution with Hubble-calibrated parameters. The included Python simulation scripts provide quantitative radial acceleration profiles ready for comparison with observational rotation curves and X-ray surface brightness profiles. These two contrasting systems validate the modularity of $F_{\text{env}}(t)$ across diverse galactic environments.

Copyright - Daniel T. Murphy, daniel.murphy00@gmail.com. UQFF Framework. PAPER_750, CP4 class #334. Session 180 continuation v5.38.

Session 225 Phonon-Physics Upgrade: GW Strain Modulation

Upgrade from PAPER_1000 (NS Merger Phonon Suppression) and PAPER_1022 (GW Phonon Strain SCm Modulation). See also PAPER_1011-1012 for GW170817/GW190425 upgraded analyses.

The late-corpus phonon analysis (Sessions 219-225) reveals that the SCm vacuum field modulates gravitational-wave strain via a frequency-dependent suppression factor. The corrected strain amplitude is:

$$h_{\text{UQFF}}(\Gamma) = h_{\text{GR}} \cdot \left(1 - 0.47 \frac{\Phi(\Gamma)}{S_{26}^{(3)}} \right)$$

where: - $\Phi(\Gamma) = \cos(\omega_{\text{SCm}} \cdot t) \cdot \Theta(H_{\text{SCm}} - 0.5)$ is the phonon modulation factor - $\omega_{\text{SCm}} = 2\pi \times 1.25 \text{ THz}$ is the SCm phonon resonance frequency - $S_{26}^{(3)} = \sum_{n=0}^{\infty} \frac{(1/4)_n (1/2)_n (3/4)_n}{(n!)^3} \cdot \prod_{i=1}^{26} [1 + [\text{SSq}] \cdot e^{-\kappa i n/26}]$ is the third-order Ramanujan summation - Θ is the Heaviside step ensuring $H_{\text{SCm}} \geq 0.5$ (phase-transition threshold)

Physical mechanism: The 1.25 THz phonon field of the SCm vacuum creates a standing-wave pattern that partially decouples the metric perturbation from the radiation zone, producing a 47% peak strain reduction for optimally oriented NS mergers. The BCS gap energy ΔE_{BCS} of the neutron-star crust couples to this phonon field, creating a mass-gap classifier that distinguishes NS from BH remnants at $M \approx 2.5 M_{\odot}$.

Calibration (canonical): $\kappa = 5 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$, $\beta_i = 0.603$, $H_{\text{SCm}} \approx 0.99$.

Session 225 Phonon-Physics Upgrade: ICM Buoyancy Force Profile

Upgrade from PAPER_1039 (SCm Galaxy Cluster Buoyancy Profile), PAPER_1041 (Cool-Core Buoyancy Balance), and PAPER_1079 (Cooling-Flow Suppression). See also PAPER_1040 (Cluster Merger Shock), PAPER_1044 (Thermal SZ Compton-y), PAPER_1046 (Cluster Lensing Mass).

The SCm phonon field introduces a buoyancy force in the ICM that modifies hydrostatic equilibrium:

$$F_{\text{buoy}}(r) = \rho(r) \cdot V \cdot g(r) \cdot \beta_i \cdot S_{26} \cdot \Phi$$

where the ICM density follows the beta-model:

$$\rho(r) = \rho_0 \left(1 + \left(\frac{r}{r_c} \right)^2 \right)^{-3\beta/2}$$

Hydrostatic mass bias reduction (PAPER_1039):

$$b_{\text{UQFF}} = 1 - \frac{M_{\text{HSE}}}{M_{\text{true}}} = 0.17 \quad (\text{vs standard } b = 0.20)$$

The buoyancy pressure contributes $P_{\text{buoy}}/P_{\text{thermal}} \approx 3\text{--}4\%$ at cluster cores, partially resolving the Planck SZ–CMB mass tension.

Cool-core stabilization (PAPER_1041/1079): AGN feedback couples to the SCm buoyancy field via $\dot{M}_{\text{cool}} = \dot{M}_0 \cdot (1 - \beta_i \cdot S_{26}^{(3)} \cdot \Phi)$, suppressing catastrophic cooling flows while maintaining observed X-ray luminosities.

Phonon frequency coupling: $\omega_{\text{SCm}} = 2\pi \times 1.25 \text{ THz}$ sets the temporal scale for buoyancy oscillations; the ratio $\omega_{\text{SCm}}/\omega_{\text{sound}}$ governs the phonon transmission efficiency across the ICM.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_{\text{lagrangian\_derivation}}`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp! \left(- \exp! \left(- \frac{r - r_0}{\lambda_{\text{VDS}}} \right) \right)$$

For this system, the local VDS sub-ratio is 0.127 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 2, \quad n_{\text{channel}} = 23/26$$

Since $p_{\text{DVP}} = 2$ is **sub-threshold** (threshold at $p > 26$), the system's vacuum topology inherits sub-threshold damping from the DVP lattice, producing smooth rather than resonant UQFF coupling profiles. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is 10^4 yr (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos! \left(\frac{2\pi j}{26} \right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh! \left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}} \right) \right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.127	PASS Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 2$	PASS Sub-threshold
BSH layers	26 harmonic terms	$j = 1\dots 26, \cos(2\pi j/26)$	PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	PASS Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_p$ suppression	$< 4.17 \times 10^{35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{\{U_Bi_i\}}$ unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{\{U_Bi_i\}}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_{corpus\_crossrefs}.py` (Session 225, April 2026).

Paper	Title
PAPER_1000	NS Merger $F_{\{U_Bi\}}$ Strain Suppression & BCS Gap
PAPER_1001	SMBH Binary Merger $F_{\{U_Bi\}}$ Phonon Damping
PAPER_1011	GW170817 NS Merger $F_{\{U_Bi_i\}}$ 66.7% Strain Reduction
PAPER_1012	GW190425 Upgraded $F_{\{U_Bi_i\}}$ with S26(3)
PAPER_1014	SMBH Merger Inspiral-Coalescence-Ringdown
PAPER_1039	SCm Galaxy Cluster Buoyancy Profile ICM Beta-Model
PAPER_1040	SCm Cluster Merger Shock Mach Number Phonon Damping
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1044	SCm Cluster Thermal SZ Effect Compton-y Phonon
PAPER_1045	SCm Cluster Radio Relic Polarization
PAPER_1046	SCm Cluster Lensing Mass Phonon Correction
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression
PAPER_1020	Cosmic Ray Phonon Acceleration DSA Spectrum
PAPER_1035	Kilonova Buoyancy Light Curve r-Process
PAPER_1050	MUGE $F_{\{U_Bi_i\}}$ Unified 9-System Synthesis
PAPER_1075	3D Volumetric MUGE Gravitational Field Generator

16 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_{kozima\_ramanujan\_appendices}.py`. For detailed derivations, see `PAPER_840/851/852/855`.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
<code>fneutron_{s26_coupling}.py</code>	neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
<code>kozima_{scm_cross_section}.py</code>	SCpm modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}]^n/26)$
<code>kozima_{wstp_kernel}.py</code>	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n \cdot \sigma_n^{\text{SCm}}(\omega) \cdot \Phi_{\text{phonon}} \cdot (F_{\text{U,Bi}}/F_{\text{U}} - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \cdot \exp[-(\omega - \omega_{\text{SCm}})^2/(2 \cdot \Gamma^2)] \cdot (1 + [\text{SSq}]^n/26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
<code>ramanujan_{polylog_s26}.py</code>	S_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
<code>s26_{wstp_kernel}.py</code>	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = \text{Li}_{26}(z) = \eta_{26}(z)/(1 - 2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} \cdot \text{Li}_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
<code>mock_{theta_q26}.py</code>	$f_{26}(q)$, $\phi_{26}(q)$, $\psi_{26}(q)$ q-series	Proper q-Pochhammer $(a; q)_{\infty}$

Core equations: $-f_{26}(q) = \sum_{n=0}^{\infty} q^{\{n2\}} / (-q; q)_{\infty}^2 - \phi_{26}(q) = \sum_{n=0}^{\infty} q^{\{n2\}} / (-q^2; q^2)_{\infty} - \psi_{26}(q) = \sum_{n=1}^{\infty} q^{\{n2\}} / (q; q^2)_{\infty}$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
<code>ramanujan_{pi_uqff}.py</code>	Classical + UQFF-modified $1/\pi + 26D$	21 digits classical, 15 UQFF, 7 digits 26D
<code>mock_{theta_pi_wstp_kernel}.py</code>	Only symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2})/9801 \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where
 $W_{26}(n) = \text{Prod}_{\{i=1\}^{26}} [1 + [SSq] \exp(-kappa_i * n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_SCm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in *CondensedPhysics.py*, *CondensedPhysics2.py*,
`MAIN_{1\_CoAnQi}.cpp`, and Wolfram kernels (`uqff_{kozima\_kernel}.wl`, `uqff_{s26\_kernel}.wl`,
`uqff_{mock\_theta\_pi\_kernel}.wl`).