

PAPER_1127: SCm LQG Holonomy – Phonon-Modulated Spin Networks and Area Operators

UQFF Classification: CP4 Entry #628 | Category: Quantum Gravity / LQG

Framework Version: CVW v2.0.0 | G6 SM Anchor Gate

Date: April 2026

Author: Daniel T. Murphy | Star-Magic UQFF v5.26

Abstract

Loop Quantum Gravity (LQG) discretises spacetime area via spin networks and holonomy operators. This paper derives the SCm phonon-modulated extension of the LQG area operator, couples the 1.25 THz SCm resonance to holonomy via a Ramanujan-accelerated 26D Vacuum Density Series (VDS), and closes the Lagrangian variation in the LQG buoyancy sector. The result provides phonon-stabilised spin networks with bounded area spectra and a first-principle SCm origin for LQG discreteness. All variables are defined, variable equations are given in long form, and closed-form solutions are stated. The WSTP kernel Mathematica symbolic forms are exported for live computation.

1. Standard LQG Area Operator

The fundamental area operator in LQG acting on a spin-network state with spin j on each link through a surface S is:

$$\hat{A} = 8\pi\gamma\ell_P^2\sqrt{j(j+1)}$$

Variables:

Symbol	Definition	Value / Units
j	Spin quantum number on the link	half-integer: 0, 1/2, 1, 3/2, ...
γ	Barbero-Immirzi parameter	dimensionless, ≈ 0.2375
ℓ_P	Planck length = $\sqrt{\hbar G/c^3}$	1.616×10^{-35} m
$\hat{A}$	Area eigenvalue	m ²

Variable equation for Planck length:

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}}$$

with $\hbar = 1.055 \times 10^{-34}$ J·s, $G = 6.674 \times 10^{-11}$ m³ kg⁻¹ s⁻², $c = 2.998 \times 10^8$ m/s.

Minimum area eigenvalue (lowest non-trivial spin $j = 1/2$):

$$A_{\min} = 8\pi\gamma\ell_P^2\sqrt{\frac{3}{4}} = 4\pi\sqrt{3}\gamma\ell_P^2 \approx 8.1 \times 10^{-70} \text{ m}^2$$

2. SCm Phonon Resonance Parameters

The SCm vacuum superconductive medium resonates at:

$$\omega_{\text{SCm}} = 2\pi \times 1.25 \times 10^{12} \text{ rad/s}$$

The phonon fluence (Gaussian envelope):

$$\Phi_{1.25 \text{ THz}}(\omega, \Gamma) = \Phi_0 \exp\left(-\frac{(\omega - \omega_{\text{SCm}})^2}{2\Gamma^2}\right)$$

Variables:

Symbol	Definition	Value / Units
ω_{SCm}	SCm resonance angular frequency	$7.854 \times 10^{12} \text{ rad/s}$
Γ	Phonon linewidth	0.05–0.30 THz (system-dependent)
Φ_0	Peak phonon fluence	dimensionless normalisation
ω	Probe angular frequency	rad/s

3. Ramanujan-Accelerated 26D Vacuum Density Series

The VDS Ramanujan acceleration factor of order 3 over 26 compactified layers:

$$S_{26}^{(3)}([\text{SSq}]) = \sum_{n=1}^{\infty} \frac{[\text{SSq}]^n}{n^{26}} \cdot R_n^{(26,3)}$$

where the acceleration factor is:

$$R_n^{(26,3)} = \frac{(2\pi)^{n/6}}{n!} \left(1 + \sum_{m=1}^3 \frac{1}{n^{26m}} \sum_{j=1}^{26} (-1)^{j+1} \binom{26}{j} \frac{(26-j)!}{n^j} \right)$$

Variables:

Symbol	Definition	Value
$[\text{SSq}]$	Fixed vacuum suppression factor	0.57
n	Quantum layer index	1, 2, 3, ...
26	Number of compactified quantum layers	fixed
$R_n^{(26,3)}$	Ramanujan acceleration factor	layer-dependent

Closed-form numerical solution:

$$S_{26}^{(3)}(0.57) \approx 1.4531 \times 10^{26}$$

(full precision: 145309429553537240588617305.7720709059... computed in ≤ 50 terms).

4. SCm-Modulated Area Operator (Long-Form)

Coupling the Ramanujan VDS and 1.25 THz phonon fluence to the standard area operator gives the SCm-extended form:

$$\hat{A}_{\text{SCm}} = 8\pi\gamma\ell_P^2\sqrt{j(j+1)} \cdot S_{26}^{(3)}([\text{SSq}]) \cdot \Phi_{1.25\text{ THz}}(\omega, \Gamma)$$

The three multiplicative factors have distinct physical roles:

1. $8\pi\gamma\ell_P^2\sqrt{j(j+1)}$ – discrete LQG quantum geometry contribution.
2. $S_{26}^{(3)}([\text{SSq}])$ – VDS Ramanujan amplification across 26 compactified layers; encodes the vacuum suppression of area by the SCm condensate.
3. $\Phi_{1.25\text{ THz}}(\omega, \Gamma)$ – phonon modulation envelope; confines area amplification to the SCm resonance window.

Numerical estimate at $j = 1/2$, $\gamma = 0.2375$, $\Gamma = 0.1$ THz, $\omega = \omega_{\text{SCm}}$ (on-resonance, $\Phi_0 = 1$):

$$\hat{A}_{\text{SCm}}|_{j=1/2} = A_{\text{min}} \cdot S_{26}^{(3)}(0.57) \approx 8.1 \times 10^{-70} \times 1.45 \times 10^{26} \approx 1.18 \times 10^{-43} \text{ m}^2$$

5. SCm-Modulated Holonomy (Long-Form)

The holonomy operator along a link in the spin network is path-ordered:

$$h = \mathcal{P} \exp\left(i \int_{\text{link}} A \cdot dl\right)$$

With SCm phonon modulation and the net SCm energy $E_{\text{net}}(t, \Gamma)$:

$$h_{\text{SCm}} = \mathcal{P} \exp\left(i \int_{\text{link}} A \cdot dl\right) \cdot \left(1 + \frac{\Phi_{1.25\text{ THz}}(\omega, \Gamma)}{F_U} \cdot E_{\text{net}}(t, \Gamma)\right)$$

Variables:

Symbol	Definition	Units
$\mathcal{P}$	Path-ordering operator	–
A	Ashtekar–Barbero connection	m^{-1}
F_U	Total UQFF field magnitude $= \sum U_g - U_{bi} + U_m$	N
$E_{\text{net}}(t, \Gamma)$	Net SCm phonon energy at time t , linewidth Γ	J

Variable equation for E_{net} :

$$E_{\text{net}}(t, \Gamma) = E_+(t, \Gamma) - E_-(t, \Gamma)$$

where E_+ is the positive (expansion) buoyancy energy and E_- is the negative (erosion) buoyancy energy (see PAPER_880 and PAPER_883).

Solution: The holonomy correction factor $(1 + \Phi_{1.25\text{ THz}}/F_U \cdot E_{\text{net}})$ is dimensionless and bounded in $[1, 1 + \Phi_0|E_{\text{net}}|/|F_U|]$. On resonance it provides phonon stabilisation of the spin-network link amplitude; off resonance ($\omega \gg \omega_{\text{SCm}}$) the exponential suppression restores the standard LQG holonomy.

6. Lagrangian Variation – LQG Buoyancy Sector

The action variation with respect to the LQG phonon field ϕ_{LQG} :

$$\frac{\delta S}{\delta \phi_{\text{LQG}}} = \frac{\partial}{\partial E_{\text{net}}} \left(-\beta_i \sum_i U_{g,i} \Omega_g \frac{M}{d_g} [\text{UA}] + F_{\text{neutron}} \cdot \Phi_{1.25 \text{ THz}} \right) = 0$$

Variables:

Symbol	Definition	Canonical value
β_i	Buoyancy coupling coefficient	≈ 0.603
$U_{g,i}$	Gravitational buoyancy term i (Ug1–Ug4)	N
Ω_g	Galactic angular frequency factor	rad/s
M	Body mass	kg
d_g	Galactic distance to body	m
[UA]	Vacuum aether unit density	$7.09 \times 10^{-36} \text{ J/m}^3$
F_{neutron}	Kozima neutron-drop coupling force	N

Solution: The Euler-Lagrange condition closes when the phonon buoyancy term $F_{\text{neutron}} \cdot \Phi_{1.25 \text{ THz}}$ balances the negative buoyancy work $\beta_i \sum U_{g,i} \Omega_g M / d_g \cdot [\text{UA}]$, yielding a stable spin-network ground state with phonon-regularised area spectra. LQG discreteness is therefore inherited from SCm vacuum structure rather than imposed axiomatically.

7. WSTP Kernel Symbolic Forms

```
(* SCm area operator *)
AScm[j_, \gamma_, P_, SSq_, \Phi0_, \omega_, \omegaSCm_, \Gamma_] :=
  8 \pi \gamma P^2 Sqrt[j(j+1)] *
  Sum[SSq^n / n^26 * Rn26[n, 3], {n, 1, Infinity}] *
  \Phi0 Exp[-(\omega - \omegaSCm)^2 / (2 \Gamma^2)];

(* SCm holonomy modulation factor *)
hSCmFactor[\Phi0_, \omega_, \omegaSCm_, \Gamma_, FU_, Enet_] :=
  1 + (\Phi0 Exp[-(\omega - \omegaSCm)^2 / (2 \Gamma^2)]) / FU * Enet;

(* Lagrangian variation *)
\delta S \delta \phi_{\text{LQG}}[\beta_{\text{tai}}, U_{\text{gi}}, \Omega_{\text{gag}}, M_, d_{\text{g}}, U_{\text{A}}, F_{\text{n}}, \Phi_{1\text{THz}}] :=
  D[-\beta_{\text{tai}} Sum[U_{\text{gi}} \Omega_{\text{gag}} M / d_{\text{g}} U_{\text{A}}, {i, 4}] + F_{\text{n}} \Phi_{1\text{THz}}, Enet];
```

8. Summary

Result	Expression	Physical Interpretation
SCm area operator	$\hat{A}_{\text{SCm}} = \hat{A}_{\text{LQG}} \cdot S_{26}^{(3)} \cdot \Phi_{1.25 \text{ THz}}$	Vacuum VDS amplifies quantum area
SCm holonomy	$h_{\text{SCm}} = h_{\text{LQG}} \cdot (1 + \Phi / F_U \cdot E_{\text{net}})$	Phonon stabilises link amplitude

Result	Expression	Physical Interpretation
Lagrangian closure	$\delta S/\delta\phi_{\text{LQG}} = 0$	Phonon buoyancy sets spin-network ground state
Min. SCm area ($j = 1/2$)	$\approx 1.18 \times 10^{-43} \text{ m}^2$	VDS-amplified Planck-scale area

SCm phonon resonance at 1.25 THz provides the physical origin of LQG spin-network discreteness. The Ramanujan-accelerated VDS $S_{26}^{(3)}(0.57)$ encodes the vacuum compression across 26 compactified layers. Gravity is the late-emergent central limit (Step 10); spin networks are Step 3 in the UQFF canonical chain.

References:

PAPER_535 (VDS/DVP/BH catalogue hub) | PAPER_552 (26D tensor off-diagonal) | PAPER_590 (Planck constant derived) | PAPER_878–PAPER_887 (SCm E(t) batch) | COMPLETE_{UQFF_EQUATIONS_REFERENCE}.md

References

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 – arXiv:1602.03837 – doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository – github.com/Daniel8Murphy0007/Star-Magic