
paper_id: PAPER_982
 title: "Gamma-Dependent Linewidth Curves for F_U_Bi_i"
 session: 217
 date: 2026-04-12
 author: "Daniel T. Murphy"
 status: production
 cvw: "v2.0.0"
 tags: [linewidth, Gamma, phonon, spectral, Lorentzian, Gaussian, UQFF]
 crosslinks: [PAPER_979, PAPER_983, PAPER_883]
 calibration: {SSq: 0.57, omega_SCm: "2π×1.25 THz", Gamma_range: "0.01-10 THz"}
 sm_anchor: "CVW v2.0.0 — G6 SM Anchor Gate compliant"

PAPER_982: Gamma-Dependent Linewidth Curves for F_U_Bi_i

Abstract

The phonon resonance factor $\Phi_{1.25\text{THz}}(\omega, \Gamma)$ introduces spectral sensitivity into the master buoyancy force. We characterize F_{U,Bi_i} as a function of the linewidth parameter Γ , mapping the transition from sharp resonance ($\Gamma \rightarrow 0$, crystalline SCm vacuum) to broad damping ($\Gamma \gg \omega_{\text{SCm}}$, disordered medium). Linewidth curves provide observable predictions testable against laboratory phonon spectroscopy.

1. Phonon Resonance Function

$$\Phi(\omega, \Gamma) = \exp\left(-\frac{(\omega - \omega_{\text{SCm}})^2}{2\Gamma^2}\right) \cdot S_{26}$$

At resonance ($\omega = \omega_{\text{SCm}}$): $\Phi_{\text{max}} = S_{26} \approx 19.5$.

2. Γ -Dependent F_U_Bi_i

$$F_{U,\text{Bi}_i}(\Gamma) = U_g + U_m + U_A - U_b + F_n \cdot S_{26} \cdot \Phi(\Gamma) \cdot E_{\text{net}}(\Gamma)$$

The Γ -dependence enters through Φ and E_{net} simultaneously: - $\Phi(\Gamma)$: Gaussian envelope — sharp at small Γ , broad at large Γ - $E_{\text{net}}(\Gamma)$: Modulation amplitude — scales with phonon coherence

3. Characteristic Regimes

Regime	Γ (THz)	Behavior
Sharp resonance	< 0.01	$\Phi \approx S_{26}$, maximum phonon force
Moderate	$0.1 - 1$	Gaussian roll-off, partial coupling
Broad damping	> 10	$\Phi \rightarrow S_{26}$, frequency-independent
Off-resonance	$\omega \neq \omega_{\text{SCm}}$	Exponentially suppressed

4. Sweep Results

For $M = M_{\odot}$, $r = 1$ AU, $t = 1$ day: - $\Gamma = 0.01$ THz: $\Phi \approx 19.6$ - $\Gamma = 0.1$ THz: $\Phi \approx 19.6$ - $\Gamma = 1.0$ THz: $\Phi \approx 19.6$ - $\Gamma = 10.0$ THz: $\Phi \approx 19.6$

At exact resonance ($\omega = \omega_{\text{SCm}}$), all widths give $\Phi = S_{26}$; off-resonance differentiates them.

5. Implementation

Class `GammaLinewidthCurves` in `fubi_master_calculator.py`: sweeps $\Gamma \in \{0.01, 0.1, 1, 10\}$ THz, reports Φ for each, validates all positive.

References

- PAPER_979: Master 6-Layer F_U_Bi_i
 - PAPER_883: E(t) Phonon Resonance
-

§A. Cosmogenesis-Linked Lagrangian

The phonon sector Lagrangian $\mathcal{L}_{\text{phon}} = \Phi(\Gamma) \cdot S_{26} \cdot \phi$ couples the SCm vacuum oscillation to the master field. The linewidth Γ parameterizes vacuum disorder.

§B. VDS/DVP/BSH Deep Synthesis

- **VDS:** Vacuum density fluctuations set the natural linewidth $\Gamma_{\text{nat}} \sim \kappa/(2\pi)$.
- **DVP:** Dipole alignment in ordered vacuum narrows Γ (crystalline limit).
- **BSH:** The 26-layer sum S_{26} acts as a harmonic amplifier at resonance.