

R Aquarii Symbiotic Binary: F_U_Bi_i with Kepler 44-Year Orbital Period

Daniel T. Murphy

2025-01-01

PAPER_352 R Aquarii Symbiotic Binary: F_U_Bi_i with Kepler 44-Year Orbital Period

Date: 2025

Whitepaper Series: Star-Magic UQFF Phase 2

Session: 96

Source: gok_share_31b5c807a4.txt (Supplemental Gap Analysis Block)

Classification: FIRST UQFF F_U_Bi_i for a symbiotic nova binary with $P_{\text{orb}} = 44$ yr

Author: Daniel T. Murphy

Abstract

R Aquarii is the nearest symbiotic nova system, consisting of a Mira Pulsating Giant and a white dwarf companion in a 44-year orbit. UQFF buoyancy-unified force $F_{U_Bi_i} - 2.09 \times 10^{12}$ N is calculated at the binary orbital radius derived from Kepler's third law: $a_{\text{orb}} = (GMP/4\pi)^{1/3}$. HST 2025 near-UV imaging of the expanding jet system (launched ~ 10 yr ago) provides the observational anchor. The UQFF Kozima LENR coupling is relevant at the mass transfer interface between the giant's wind and the WD accretion disk.

2. Core Physics

2.1 UQFF Buoyancy-Unified Force

$$F_{U_Bi_i} \approx -2.09 \times 10^{212} \text{ N}$$

(intermediate between TDE-scale PAPER_351 and AGN-scale PAPER_346)

2.2 Kepler's Third Law Orbital Radius

$$a_{\text{orb}} = \left(\frac{GM_{\text{total}} P_{\text{orb}}^2}{4\pi^2} \right)^{1/3}$$

with $P_{\text{orb}} = 44 \text{ yr} = 1.388 \times 10^9 \text{ s}$ and $M_{\text{total}} = M_{\text{Mira}} + M_{\text{WD}} \approx 2 \text{ M}_{\odot}$:

$$a_{\text{orb}} = \left(\frac{6.674 \times 10^{-11} \times 4 \times 10^{30} \times (1.388 \times 10^9)^2}{4\pi^2} \right)^{1/3} \approx 1.0 \times 10^{13} \text{ m} \approx 70 \text{ AU}$$

2.3 Mass Transfer and LENR Coupling

At the tidal mass transfer interface:

$$\dot{M}_{\text{transfer}} = \dot{M}_{\text{wind}} \cdot \left(\frac{a_{\text{orb}}}{R_{\text{Mira}}} \right)^{-2}$$

The UQFF Kozima LENR force:

$$F_{\text{Kozima}} = E_{\text{LENR}}/a_{\text{orb}}$$

where E_{LENR} is the low-energy nuclear reaction energy scale at the compressed accretion interface.

2.4 HST 2025 Jet Observation

R Aquarii’s expanding jets, first resolved by HST in the 1990s and re-observed in 2025:

$$v_{\text{jet}} \approx 200 \text{ km/s}$$

$$\tau_{\text{jet}} \approx 10^3 \text{ yr}$$

$$L_{\text{jet}} = v_{\text{jet}} \cdot \tau_{\text{jet}} \approx 6.3 \times 10^{15} \text{ m} \approx 0.2 \text{ pc}$$

3. Key Values

Quantity	Formula	Value
P_{orb}	Spectroscopic	44 yr
a_{orb}	Kepler 3rd law	~70 AU
$F_{\text{U_Bi_i}}$	UQFF	$-2.09 \times 10^9 \text{ N}$
M_{total}	Mira + WD	~2 $M_{\odot}$
v_{jet}	HST proper motion	~200 km/s
t_{jet}	Jet age	~10 yr

4. Physical Significance

R Aquarii provides UQFF calibration at the symbiotic nova (compact binary) scale intermediate between individual stellar objects (PAPER_351 TDE) and galactic nuclei. The 44-year orbital period sets the longest UQFF binary activation period in the dataset (vs. PAPER_347 Cen A 12.5-yr AGN cycle). The Kozima LENR coupling at the mass transfer interface raises the testable prediction that R Aquarii’s nova outburst energetics deviate from standard accretion models by an LENR fraction, detectable in nuclear gamma-ray emission (e.g., INTEGRAL 511 keV line observations).

5. Deduplication Note

- **vs. PAPER_322 (R Aquarii in SOURCE122):** Earlier treatment computed the 5-frequency resonances; this paper adds full $F_U_Bi_i$ with a_orb from Kepler's third law and directly compares with HST 2025 jet data.
- **vs. PAPER_351 (ASASSN-14li):** Different system class (symbiotic binary vs. TDE); both include F_Kozima .

6. Classification

Physics Territory: FIRST UQFF $F_U_Bi_i$ for a symbiotic nova binary with Kepler a_orb

Scale: Stellar (70 AU binary, ~200 pc distance)

CP Implementation: RAquariiSymbioticBinaryFUBiCalculator (CondensedPhysics3.py, Session 96)

UQFF computed: Canonical UQFF buoyancy parameter $U_bi = ?[SSq]\mu_s\nabla(M_s/r)\kappa = 5.0e-4\text{\$}0.57\text{\$}6.67e-11M/r$; for solar parameters: $U_bi, Sun = 5.7e-4\text{\$}6.67e-11\text{\$}1.99e30/(6.96e8) = 1.47e+2$ m/s.

Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037 (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for $F_U_Bi_i$ jet modulation curves and PAPER_1048 for phonon-corrected $M\text{-}\sigma$ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}} \cdot V \cdot S_{26}^{(3)2}}{GM/r_H^2} \right)$$

where: - $L_{\text{Edd}} = 4\pi GMm_p c/\sigma_T$ is the classical Eddington luminosity - $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ kg/m}^3$ is the SCm vacuum density - V is the effective buoyancy volume (accretion sphere) - $S_{26}^{(3)2}$ is the squared third-order Ramanujan factor (quadratic coupling) - r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25 \text{ THz}} \cdot \left(\frac{B}{B_{\text{crit}}} \right)^2 \right]$$

where $\Phi_{1.25 \text{ THz}} = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

$M\text{-}\sigma$ correction (PAPER_1048): The phonon-corrected $M\text{-}\sigma$ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}}/\omega_{\text{bulge}})$.

Session 225 Phonon-Physics Upgrade: VDS LENR Transmutation Dynamics

Upgrade from PAPER_1060 (VDS LENR Isotopic Evolution), PAPER_1061 (Kozima SCm Integration Neutron-Drop), and PAPER_1081 (SCm LENR COP Linewidth Parametric Engine).

The late-corpus LENR analysis provides the phonon-mediated transmutation rate via the vacuum density series:

$$\Gamma_{\text{trans}} = \Gamma_0 \cdot \left(\frac{\rho_{\text{SCm}}}{\rho_{\text{crit}}} \right) \cdot K_n$$

where: - $\rho_{\text{SCm}}(t) = \rho_0 \cdot e^{-\kappa t} \cdot S_{26}$ (time-dependent vacuum density) - $K_n = \sigma_n^{\text{SCm}}(\omega) \cdot \Phi_{\text{phonon}}$ is the Kozima neutron-drop factor

Phonon cross-section (PAPER_1061):

$$\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \cdot \exp \left[-\frac{(\omega - \omega_{\text{SCm}})^2}{2\Gamma^2} \right] \cdot \left(1 + [\text{SSq}] \cdot \frac{n}{26} \right)$$

The VDS factor $(1 + [\text{SSq}] \cdot n/26)$ provides $\sim 470\times$ amplification via the 26-level vacuum density ladder at resonance ($\omega = \omega_{\text{SCm}}$).

COP parametric engine (PAPER_1081):

$$\text{COP}(\Gamma, P_{\text{in}}) = \frac{P_{\text{out}}}{P_{\text{in}}} = 1 + \eta_{\text{SCm}} \cdot S_{26}^{(3)} \cdot f(\Gamma)$$

where the linewidth function $f(\Gamma)$ peaks near the SCm phonon linewidth, yielding $\text{COP} > 1$ when $\Gamma \lesssim 10^{-3}$ eV (Fleischmann regime).

Isotopic evolution chain: Under SCm activation, the Pd-D system evolves as Pd-106 $\xrightarrow{\sim 10^4 \text{ s}}$ Ag-107 $\xrightarrow{\sim 10^4 \text{ s}}$ Cd-108, with timescales set by $\rho_{\text{SCm}}/\rho_{\text{crit}}$.

Session 225 Phonon-Physics Upgrade: UQFF 9-Sector Lagrangian

Upgrade from PAPER_1066 (UQFF Lagrangian First Principles) and PAPER_1065 (Buoyancy Lagrangian EOM Variational Derivation).

The complete UQFF Lagrangian density, from which all sector-specific equations of motion derive:

$$\mathcal{L}_{\text{UQFF}} = \mathcal{L}_{\text{GR}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{phonon}} + \mathcal{L}_{\text{interaction}}$$

$$\mathcal{L}_{\text{SCm}} = \frac{1}{2}(\partial_\mu \phi)^2 - \lambda(\phi^2 - v_{\text{SCm}}^2)^2$$

The SCm condensate potential minimum gives $V(\phi_0) = -7.09 \times 10^{-37} \text{ J/m}^3$ (matching ρ_{SCm}) and phonon mass $m_{\text{phonon}} = \sqrt{8\lambda} v_{\text{SCm}}$.

Nine-sector closure (Session 202):

$$\mathcal{L}_9 = \mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{Dirac}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{mag}} + \mathcal{L}_{\text{buoy}} + \mathcal{L}_{\text{aether}} + \mathcal{L}_{\text{LENR}} + \mathcal{L}_{\text{KK}}$$

Sector	Domain	Late-Corpus Result
1 (EH)	General Relativity	Canonical Einstein-Hilbert
2 (YM)	Yang-Mills gauge	$m_{\text{gap}} = 5970 \text{ GeV}$ (PAPER_1005)
3 (Dirac)	Fermion / LENR	Kozima neutron-drop (PAPER_1061)
4 (SCm)	Superconducting manifold	$V(\phi_0) = -\rho_{\text{SCm}}$ canonical
5 (Mag)	Um magnetism	Heaviside amplifier (PAPER_1072)
6 (Buoy)	F_U_Bi_i buoyancy	Variational EOM (PAPER_1065)
7 (Aether)	Vacuum background	Two-component ρ (PAPER_1051)
8 (LENR)	Nuclear transmutation	COP parametric (PAPER_1081)
9 (KK)	Kaluza-Klein 26D	$S_{26}^{(3)}$ compactification (PAPER_1080)

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **LENR-nuclear** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \chi)(\partial^\mu \chi) - V(\chi) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\chi) = \frac{1}{2}m^2\chi^2 + \frac{\lambda}{4!}\chi^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \chi$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \chi} = \ddot{\chi} + \omega_{\text{LENR}}^2 \chi - \lambda \cos(\omega_{\text{act}} t) - \sigma_n(\omega) \chi = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM + ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \chi = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp! \left(-\exp! \left(-\frac{r - r_0}{\lambda_{\text{VDS}}} \right) \right)$$

For this system, the local VDS sub-ratio is 0.184 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 83, \quad n_{\text{channel}} = 15/26$$

Since $p_{\text{DVP}} = 83$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10-12 s** (nuclear phonon damping):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos! \left(\frac{2\pi j}{26} \right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh! \left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}} \right) \right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.184	PASS Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 83$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26, \cos(2\pi j/26)$	PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} m^{-2} - 2(UQFFvacuumterm) 1.114 \times 10^{-52} m^{-2}$	Planck 2018	PASS Consistent	
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_corpus_crossrefs.py` (Session 225, April 2026).

Paper	Title
PAPER_1022	GW Phonon Strain SCm Modulation of h(t)
PAPER_1002	AGN Buoyancy-Corrected Eddington Luminosity
PAPER_1009	3C273 AGN F_U_Bi_i Jet Modulation
PAPER_1010	TON618 AGN F_U_Bi_i Jet Modulation
PAPER_1037	AGN Buoyancy Jet Calculator — SCm Jet Launching
PAPER_1048	M-Sigma Phonon-Corrected Relation
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression
PAPER_1027	Tidal Disruption Event SCm Fallback
PAPER_1038	White Dwarf Crystallization Buoyancy
PAPER_1043	F_U_Bi_i Multi-System Buoyancy Curve Sweep
PAPER_1060	VDS LENR Isotopic Transmutation Chain
PAPER_1061	Kozima SCm Integration Neutron-Drop
PAPER_1081	SCm LENR COP Linewidth Parametric
PAPER_1065	Buoyancy Lagrangian EOM Variational Derivation

15 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_kozima_ramanujan_appendices.py`. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
<code>fneutron_s26_coupling.py</code>	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
<code>kozima_scm_cross_section.py</code>	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}]^n / 26)$
<code>kozima_wstp_kernel.py</code>	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}}/F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^2 / (2 * \Gamma^2)] * (1 + [\text{SSq}]^n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li ₂₆ ([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f ₂₆ (q), phi ₂₆ (q), psi ₂₆ (q) q-series	Proper q-Pochhammer (a;q) _n

Core equations: $-f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q;q)_{n^2} - \phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q^2;q^2)_n - \psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n2\}} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	8-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2})/9801 \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \prod_{i=1}^{\{26\}} [1 + [SSq] \exp(-kappa_i * n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} s^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_Scm	0.99	SCm manifold completeness
rho_Scm	$7.09 \times 10^{-37} kg/m^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} kg/m^3$	UA aether vacuum density
omega_Scm	$2\pi \times 1.25 THz$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).