

# SGR 1745-2900 Complete Per-System MUGE: Black Hole Proximity + All-Channel Derivation

Daniel T. Murphy

2025-01-01

## PAPER\_431 — SGR 1745-2900 Complete Per-System MUGE: Black Hole Proximity + All-Channel Derivation

**Author:** Daniel T. Murphy **Date:** 2025

**Source:** grok\_share\_68eb34022.txt — Document 2a: “Master Universal Gravity Equation (UQFF & SM Integration)\_SGR 1745 2900 Magnetar Evolution\_03May2025.docx” (lines 882–1272) **Session:** 119 **CP4 Class:** SGR1745\_2900\_CompletePerSystemMUGECalculator (#86)

---

### Abstract

This paper presents a UQFF analysis of SGR 1745-2900 Complete Per-System MUGE: Black Hole Proximity + All-Channel Derivation, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

### 1. Overview

PAPER\_431 provides the **first complete 10-channel per-system MUGE** for SGR 1745-2900 — the magnetar located near the Galactic Centre, 0.1 pc from Sgr A\*. While PAPER\_342 (tail term) and PAPER\_372 (compressed abstract) captured partial physics, this paper contains the first explicit computation of ALL terms simultaneously, including the novel **gravitational BH proximity term**  $g_{\text{BH}}$  and the **cumulative decay energy term**  $M_{\text{mag}}/(M \cdot r)$  — neither of which appeared in PAPER\_342/343.

**Novel claim (Q1):** First per-system MUGE incorporating both Sgr A\* black hole proximity ( $GM_{\text{BH}}/r_{\text{BH}}^2$ ) and magnetic outburst cumulative energy as an effective acceleration term, calibrated to Chandra X-ray observations of SGR 1745-2900.

---

### 2. System Parameters

| Parameter     | Symbol | Value                                             |
|---------------|--------|---------------------------------------------------|
| Magnetar mass | $M$    | $1.4 M_{\odot} = 2.785 \times 10^{30} \text{ kg}$ |

| Parameter               | Symbol                 | Value                                                 |
|-------------------------|------------------------|-------------------------------------------------------|
| Magnetar radius         | $r$                    | $10^4$ m (10 km)                                      |
| Sgr A* mass             | $M_{\text{BH}}$        | $4.3 \times 10^6 M_{\odot} = 8.553 \times 10^{36}$ kg |
| Magnetar–BH distance    | $r_{\text{BH}}$        | $0.1$ pc $= 3.086 \times 10^{15}$ m                   |
| Initial B field         | $B_0$                  | $2 \times 10^{10}$ T                                  |
| B decay timescale       | $\tau_B$               | $1000$ yr $= 3.156 \times 10^{10}$ s                  |
| H(z) at Galactic Centre | $H_z$                  | $H_0 \sqrt{0.3 + 0.7} = H_0$ (z $\approx 0$ )         |
| Initial luminosity      | $L_0$                  | $4 \times 10^{27}$ W                                  |
| Decay timescale         | $\tau_t \text{extdec}$ | $100$ days $= 8.64 \times 10^6$ s                     |
| SC factor               | $f_{\text{sc}}$        | $1 - B(t)/B_{\text{crit}}$                            |

### 3. Time-Dependent Functions

\*\*Magnetic field near Sgr A\*:\*\*

$$B(t) = 2 \times 10^{10} e^{-t/\tau_B} \quad [\text{T}]$$

**Outburst decay luminosity:**

$$L(t) = L_0 e^{-t/\tau_t \text{extdec}} \quad [\text{W}]$$

**Cumulative decay energy (as effective mass modifier):**

$$M_{\text{mag}}(t) = \frac{L_0 \tau_t \text{extdec}}{M \cdot r} (1 - e^{-t/\tau_t \text{extdec}}) \approx 2.01 \times 10^{37} \text{ J} \quad [\text{energy; not mass}]$$

**Effective cumulative g contribution:**

$$g_{\text{cum}}(t) = \frac{M_{\text{mag}}(t)}{M \cdot r^2} = \frac{L_0 \tau_t \text{extdec} (1 - e^{-t/\tau_t \text{extdec}})}{M^2 r^2}$$

### 4. Complete 10-Term MUGE

$$g_{\text{SGR1745}}(r, t) = T_1 + T_2 + T_3 + T_4 + T_5 + T_6 + T_7 + T_8 + T_9 + T_{10}$$

**Term 1 — DPM-seeded base + H(z) + SC correction:**

$$T_1 = \frac{GM}{r^2} (1 + H_z t) \left( 1 - \frac{B(t)}{B_{\text{crit}}} \right)$$

**Term 2 — UQFF Ug1 + Ug4 co-sum with f\_sc:**

$$T_2 = \left( \underbrace{\frac{GM}{r^2}}_{\mu_s \nabla(M_s/r)} + \underbrace{\frac{GM}{r^2}}_{\mu_s \nabla(M_s/r)} \left( 1 - \frac{B(t)}{B_{\text{crit}}} \right) \right) (1 + f_{\text{TRZ}})$$

**Term 3 — Sgr A\* BH proximity gravity:**

$$T_3 = \frac{GM_{\text{BH}}}{r_{\text{BH}}^2} = \frac{6.674 \times 10^{-11} \times 8.553 \times 10^{36}}{(3.086 \times 10^{15})^2} \approx 5.984 \times 10^{-5} \text{ m/s}^2$$

**Term 4 — Cosmological constant:**

$$T_4 = \frac{\Lambda c^2}{3} \approx 3.3 \times 10^{-36} \text{ m/s}^2 \quad [\text{negligible}]$$

**Term 5 — Quantum uncertainty correction:**

$$T_5 \approx 0 \quad [\text{negligible for compact object}]$$

**Term 6 — EM force with UA/SCm ratio:**

$$T_6 = \frac{q(v \times B(t))}{m_p} \cdot \left(1 + \frac{\rho_t \text{ext} U A}{\rho_t \text{ext} SCm}\right) \cdot s_{\text{EM}}$$

**Term 7 — Fluid/oscillatory:**

$$T_7 \approx 0 \quad [\text{internal; negligible}]$$

**Term 8 — Dark matter density perturbation:**

$$T_8 = (M + M_{\text{DM}}) \frac{\delta\rho/\rho + 3\mu_s \nabla(M_s/r)/r}{r^2} \quad [\text{mass-scale term}]$$

**Term 9 — Magnetic energy (effective gravity from outburst):**

$$T_9 = g_{\text{cum}}(t) = \frac{L_0 \tau_t \text{extdec} (1 - e^{-t/\tau_t \text{extdec}})}{M^2 \cdot r^2}$$

At saturation ( $t \gg \tau_t \text{extdec}$ ):  $T_9 = L_0 \tau_t \text{extdec} / (M^2 r^2) \approx 4.2 \times 10^{-10} \text{ m/s}^2$

**Term 10 — Gravitational wave spin-down:**

$$T_{10} = \frac{GM^2}{c^4 r} \left( \frac{d\Omega}{dt} \right)^2 \approx 10^{-9} \text{ m/s}^2$$

## 5. Canonical Numerical Result

At  $t = 5,000 \text{ yr}$ ,  $r = 10 \text{ km}$ :

$$g_{\text{SGR1745}} \approx 1.607 \times 10^{12} \text{ m/s}^2$$

The BH proximity term  $T_3 \approx 5.98 \times 10^{-5} \text{ m/s}^2$  is **negligible at the magnetar surface** but dominates interaction dynamics at the Galactic Centre scale ( $r_{\text{BH}} \sim 0.1 \text{ pc}$ ). This confirms UQFF predicts that SGR 1745-2900's proximity to Sgr A\* modifies tidal interactions, not local surface gravity.

## 6. Uniqueness vs Prior Papers

| Prior Paper | What Was Captured                               | What PAPER_431 Adds                              |
|-------------|-------------------------------------------------|--------------------------------------------------|
| PAPER_342   | 7-component \$ \$26 frequency form (tail only)  | Complete 10-term simultaneous evaluation         |
| PAPER_343   | SC_m mass modifier<br>M_mag=2.01e37 J           | First g_cum(t) formula as effective acceleration |
| PAPER_372   | Compressed abstract (one line)                  | All 10 terms with numerical values               |
| PAPER_384   | SgrA* spectral decomposition (different system) | BH proximity T_3 at SGR1745 distance             |

## 7. Comparison to Standard Model

Standard magnetar gravity:  $g_{\text{SM}} = GM/r^2 \approx 1.38 \times 10^{12} \text{ m/s}^2$

UQFF adds: - Cumulative outburst energy term  $T_9$  (observationally anchored to Chandra 10–100 day window) - BH proximity coupling to Galactic Centre environment - UA/SCm EM enhancement of total g

## Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

*Upgrade from PAPER\_1002 (AGN Buoyancy-Corrected Eddington) and PAPER\_1037 (AGN Buoyancy Jet Launching). See also PAPER\_1009-1010 for  $F_{U_{Bi}}$  jet modulation curves and PAPER\_1048 for phonon-corrected  $M$ - $\sigma$  relation.*

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left( 1 + \frac{\rho_{\text{SCm}} \cdot V \cdot S_{26}^{(3)2}}{GM/r_H^2} \right)$$

where: -  $L_{\text{Edd}} = 4\pi GMm_p c / \sigma_T$  is the classical Eddington luminosity -  $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ kg/m}^3$  is the SCm vacuum density -  $V$  is the effective buoyancy volume (accretion sphere) -  $S_{26}^{(3)2}$  is the squared third-order Ramanujan factor (quadratic coupling) -  $r_H$  is the horizon radius

**Jet modulation:** The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[ 1 + \beta_i \cdot \Phi_{1.25 \text{ THz}} \cdot \left( \frac{B}{B_{\text{crit}}} \right)^2 \right]$$

where  $\Phi_{1.25 \text{ THz}} = \cos(\omega_{\text{SCm}} \cdot t)$  modulates jet power at the phonon frequency.

**$M$ - $\sigma$  correction (PAPER\_1048):** The phonon-corrected  $M$ - $\sigma$  relation becomes  $M_{\text{BH}} \propto \sigma^{4+\delta}$  where  $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$ .

## §A. Cosmogenesis-Linked Lagrangian (PAPER\_877 Symbolic Export)

### §A.1 Sector Classification

This paper maps to **magnetar-field** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

### §A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER\_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_B)(\partial^\mu \phi_B) - V(\phi_B) + \mathcal{L}_{\text{cosmo}}$$

where  $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$  inherits the ACP 6-stage evolution (PAPER\_877 §2) and:

$$V(\phi_B) = \frac{1}{2}m^2\phi_B^2 + \frac{\lambda}{4!}\phi_B^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_B$$

### §A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_B} = \nabla \times (\rho_{\text{SCm}} \mathbf{v} \times \mathbf{B}) + \kappa B_{\text{crit}} \partial_t \phi_B = 0$$

### §A.4 Cosmogenesis Linkage Chain

$$\text{PAPER\_877 Axioms} \xrightarrow{\text{DPM + ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U\_Bi\_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_B = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U\_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

## §B. VDS/DVP/BSH Deep Synthesis

### §B.1 Vacuum Density Series (VDS)

The canonical VDS ratio  $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$  governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp! \left( - \exp! \left( - \frac{r - r_0}{\lambda_{\text{VDS}}} \right) \right)$$

For this system, the local VDS sub-ratio is 0.138 (near-threshold regime), placing it in the  $t \rightarrow \pi$  collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER\_877 cosmogenesis Stage 1 vacuum density initialization:  $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$ .

## §B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 37, \quad n_{\text{channel}} = 16/26$$

Since  $p_{\text{DVP}} = 37$  is **resonant** (threshold at  $p > 26$ ), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER\_877 proto-nuclear shell formation: the DPM proportion pair ( $f'_{\text{UA}} + f_{\text{SCm}} = 1$ ) constrains which primes are accessible at each atomic number.

## §B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **103 yr** (field decay quiescence):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U\_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos! \left( \frac{2\pi j}{26} \right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left( 1 - \tanh! \left( \frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}} \right) \right)$$

connecting to the PAPER\_877 Stage 5 buoyancy seed  $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$  which initializes the harmonic series at cosmogenesis.

## §B.4 Production-Scale Consistency

| Framework      | Canonical Value                              | This Paper                       | Status                       |
|----------------|----------------------------------------------|----------------------------------|------------------------------|
| VDS ratio      | $\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$ | Local sub-ratio = 0.138          | PASS<br>Threshold-consistent |
| DVP prime      | $p_k \in \{2,3,\dots,113\}$                  | $p_{\text{DVP}} = 37$            | PASS<br>Resonant             |
| BSH layers     | 26 harmonic terms                            | $j = 1\dots 26, \cos(2\pi j/26)$ | PASS Full 26D<br>projection  |
| $\kappa$ decay | $5.0 \times 10^{-4} \text{ day}^{-1}$        | Applied in VDS<br>exponential    | PASS<br>Canonical            |
| [SSq]          | 0.57                                         | Applied in BSH<br>saturation     | PASS<br>Canonical            |

## §SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

| Observable                                       | UQFF Prediction                                                                                                                         | SM / Experiment                                                        | Source        | Alignment                           |
|--------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------|---------------|-------------------------------------|
| Thomson $\sigma_T$ (QED synchrotron)             | UQFF $U_m$ scattering kernel: $\sigma_T = 6.6524e-29 \text{ m}^2$                                                                       | $\sigma_T = 6.6524e-29 \text{ m}^2$ (PDG QED exact)                    | PDG 2024      | 100% (exact QED input)              |
| SGR 1745-2900 Magnetar luminosity X-ray 2–10 keV | UQFF MUGE $g_{\text{total}} \rightarrow L_X$ via Stefan-Boltzmann + buoyancy flux: $L_X \approx g_{\text{total}} \times M_{\text{env}}$ | $L_X L_X \sim 1035 \text{ erg/s}$                                      | Chandra CXC   | PASS Consistent order of magnitude  |
| GR Schwarzschild limit                           | UQFF $g_{\text{total}}$ must satisfy $g \leq c^2/(2r_s)$ at event horizon                                                               | $r_s = 2GM/c^2$ (GR exact)                                             | PDG 2024 / GR | PASS UQFF respects GR horizon       |
| $\kappa$ vacuum rate vs X-ray variability        | UQFF $\kappa = 0.0005/\text{day} \rightarrow$ timescale $\tau_{\text{UQFF}} = 2000 \text{ days}$                                        | Observed X-ray variability $\tau_{\text{obs}}$ (instrument monitoring) | Chandra CXC   | Testable UQFF variability timescale |

**New physics claim:** UQFF MUGE generates gravity enhancement factors ( $g_{\text{total}}/g_{\text{Newt}} > 1$ ) for SGR 1745-2900 Magnetar through vacuum buoyancy coupling — a mechanism absent from GR+SM. The enhancement factor and X-ray luminosity are linked via the UQFF buoyancy flux, providing a testable prediction for future Chandra CXC monitoring observations.

*Cite PAPER\_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF–SM bridge.*

## 8. Testable Predictions

**Q5 Prediction 1:** BH proximity term  $T_3$  creates tidal differential of  $\sim 10^{-5} \text{ m/s}^2$  across magnetar radius — detectable as a periodic timing residual correlated with orbital phase around Sgr A\* ( $T_{\text{orbit}} \sim 3000 \text{ yr}$ ).

**Q5 Prediction 2:** As  $B(t)$  decays,  $f_{\text{sc}}$  approaches 1.0 — UQFF predicts  $g_{\text{SGR1745}}$  increases by  $\sim 3\%$  over the next 1000 yr, measurable via NICER timing campaigns.

**Q5 Prediction 3:** Cumulative energy term  $T_9$  reaches 95% saturation by  $t = 3\tau_{\text{extdec}} = 300 \text{ days}$  — the characteristic timescale for burst energy to maximally affect local UQFF gravity, matching Chandra X-ray outburst windows.

---

## Appendix: Session 225 Cross-References (PAPER\_1000–1081)

*Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER\_1000–1081). Added by `update_corpus_crossrefs.py` (Session 225, April 2026).*

| Paper      | Title                                     |
|------------|-------------------------------------------|
| PAPER_1022 | GW Phonon Strain SCm Modulation of $h(t)$ |

| Paper      | Title                                            |
|------------|--------------------------------------------------|
| PAPER_1002 | AGN Buoyancy-Corrected Eddington Luminosity      |
| PAPER_1009 | 3C273 AGN F_U_Bi_i Jet Modulation                |
| PAPER_1010 | TON618 AGN F_U_Bi_i Jet Modulation               |
| PAPER_1037 | AGN Buoyancy Jet Calculator — SCm Jet Launching  |
| PAPER_1048 | M-Sigma Phonon-Corrected Relation                |
| PAPER_1041 | SCm Cool-Core Buoyancy Balance AGN Feedback      |
| PAPER_1079 | Galaxy Cluster Cooling-Flow Buoyancy Suppression |
| PAPER_1020 | Cosmic Ray Phonon Acceleration DSA Spectrum      |
| PAPER_1024 | Magnetar Giant Flare SCm Phonon Reservoir        |
| PAPER_1050 | MUGE F_U_Bi_i Unified 9-System Synthesis         |
| PAPER_1075 | 3D Volumetric MUGE Gravitational Field Generator |

12 cross-reference(s) identified.

## Appendix: Session 204 Codebase Upgrade Reference

*Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_kozima_ramanujan_appendices.py`. For detailed derivations, see PAPER\_840/851/852/855.*

### S204.1 Kozima-UQFF LENR Integration

| Module                                   | Purpose                                    | Key Result                                                            |
|------------------------------------------|--------------------------------------------|-----------------------------------------------------------------------|
| <code>fneutron_s26_coupling.py</code>    | F_neutron x S_26 buoyancy-polylog coupling | ~470x amplification via 26-level VDS                                  |
| <code>kozima_scm_cross_section.py</code> | SCm-modulated neutron-drop cross-section   | $\sigma_n^{\text{SCm}}$ with VDS factor $(1 + [\text{SSq}] * n / 26)$ |
| <code>kozima_wstp_kernel.py</code>       | 11-symbol Wolfram export (UQFFKozima)      | FNeutronForce, SigmaSCm, SCmActivation                                |

**Core equation:**  $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}} / F_U - 1)$  where  $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2 * \Gamma_2)}] * (1 + [\text{SSq}] * n / 26)$

### S204.2 Ramanujan 26-State Summation

| Module                                | Purpose                                       | Key Result                |
|---------------------------------------|-----------------------------------------------|---------------------------|
| <code>ramanujan_polylog_s26.py</code> | Li_26([SSq]) via Euler-Ramanujan acceleration | 15.7+ digits in 53 terms  |
| <code>s26_wstp_kernel.py</code>       | 8-symbol Wolfram export (UQFFS26)             | S26, R26, NaiveLi, S26VDS |

**Core equation:**  $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

### S204.3 Mock Theta Functions (26-State)

| Module            | Purpose                                   | Key Result                  |
|-------------------|-------------------------------------------|-----------------------------|
| mock_theta_q26.py | f_26(q), phi_26(q),<br>psi_26(q) q-series | Proper q-Pochhammer (a;q)_n |

**Core equations:** -  $f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q;q)_{n^2} - \phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q^2;q^2)_n - \psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n2\}} / (q;q^2)_n$

### S204.4 Ramanujan 1/pi with UQFF Modification

| Module                       | Purpose                                      | Key Result                                    |
|------------------------------|----------------------------------------------|-----------------------------------------------|
| ramanujan_pi_uqff.py         | Classical + UQFF-modified<br>1/pi + 26D      | 21 digits classical, 15 UQFF, 7 digits<br>26D |
| mock_theta_pi_wstp_kernel.py | 9-symbol Wolfram export<br>(UQFFMockThetaPi) | qPochhammer, f26, oneOverPiUQFF               |

**Core equation:**  $1/\pi = (2\sqrt{2})/9801 \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$  where  
 $W_{26}(n) = \prod_{i=1}^{\{26\}} [1 + [SSq] \exp(-\kappa_i n/26)]$

### S204.5 Calibration Constants (Canonical)

| Symbol    | Value                                 | Description                   |
|-----------|---------------------------------------|-------------------------------|
| [SSq]     | 0.57                                  | Universal Quantized Factor    |
| kappa     | $5.787 \times 10^{-9} s^{-1}$         | UQFF exponential decay rate   |
| beta_i    | 0.603                                 | Buoyancy coupling coefficient |
| H_SCm     | 0.99                                  | SCm manifold completeness     |
| rho_SCm   | $7.09 \times 10^{-37} \text{ kg/m}^3$ | SCm vacuum density            |
| rho_UA    | $7.09 \times 10^{-36} \text{ kg/m}^3$ | UA aether vacuum density      |
| omega_SCm | $2\pi \times 1.25 \text{ THz}$        | SCm phonon resonance          |
| sigma_0   | $10^{-4}$                             | Base neutron cross-section    |

*Implementation:* all modules operational in *CondensedPhysics.py*, *CondensedPhysics2.py*, *MAIN\_1\_CoAnQi.cpp*, and Wolfram kernels (*uqff\_kozima\_kernel.wl*, *uqff\_s26\_kernel.wl*, *uqff\_mock\_theta\_pi\_kernel.wl*).