

Yang-Mills Mass Gap $\Delta_{YM}(T)$ via $S^{(k)}$

Daniel T. Murphy

2026-04-12

PAPER_971: Yang-Mills Mass Gap $\Delta_{YM}(T)$ via $S^{(k)}$

Author: Daniel T. Murphy — Star Magic / UQFF Framework **Date:** 2026-04-12 **Session:** 216
Source: qgp_ramanujan_application.py (YangMillsMassGapCalculator) **Calculator:** YangMills-MassGapCalc (CP4 #555) **CVW:** v2.0.0 compliant

Abstract

We express the Yang-Mills mass gap $\Delta_{extYM}(T)$ in terms of the QCD scale Λ_{extQCD} and the expanded Ramanujan factor $S_{26}^{(k)}$. The gap closes continuously at the deconfinement temperature $T_c = 1.5 \times 10^{12}$ K, providing a UQFF-native derivation of confinement/deconfinement.

1. Yang-Mills Mass Gap

$$\Delta_{extYM}(T) = \Lambda_{extQCD} \cdot \left(1 - \frac{T}{T_c}\right) \cdot S_{26}^{(k)}([SSq])$$

For $T < T_c$: gap is open (confinement). At $T = T_c$: gap closes (deconfinement).

2. Connection to Millennium Prize

The Clay Millennium Prize Yang-Mills problem asks for proof that gauge theory has a mass gap $\Delta > 0$. The UQFF framework provides: - A temperature-dependent gap function - Closure at T_c via $S_{26}^{(k)}$ modulation - Consistency with lattice QCD measurements of $\Lambda_{extQCD} \approx 217$ MeV

3. Gap vs Temperature

T (K)	Δ_{extYM} (eV)	State
10^{11}	$\approx \Lambda_{extQCD} \cdot S_{26}^{(k)}$	Confined
10^{12}	$\frac{1}{3} \Lambda_{extQCD} \cdot S_{26}^{(k)}$	Confined
1.5×10^{12}	0	Deconfined
$> T_c$	0	QGP

References

1. Murphy, D.T. — Star Magic UQFF Framework (2024-2026)
2. PAPER_969 — Expanded 26D Ramanujan $S_{26}^{(k)}$
3. PAPER_970 — QGP Vacuum Density $\rho_{t\text{ext}QGP}$
4. Clay Mathematics Institute — Yang-Mills and Mass Gap
5. Particle Data Group — $\Lambda_{t\text{ext}QCD}$ measurement

Cross-Links

Related Paper	Relationship
PAPER_969	$S_{26}^{(k)}$ acceleration factor
PAPER_970	QGP vacuum density (companion)
PAPER_972	ALICE centrality (experimental context)
PAPER_530	Millennium Hub (YM+Riemann+PvsNP)

Session 225 Phonon-Physics Upgrade: Yang-Mills BCS Phonon Mass Gap

Upgrade from PAPER_1005 (Yang-Mills Mass Gap via SCm BCS Phonon) and PAPER_1070 (Yang-Mills Mass Gap VDS Bridge). See also PAPER_1004 (QGP Vacuum Density), PAPER_1007 (Deconfinement Phase Diagram), PAPER_1059 (CGC BK Saturation), PAPER_1064 (Resummation BFKL/Sudakov).

The late-corpus analysis derives the Yang-Mills mass gap via a BCS-like phonon pairing mechanism in the SCm vacuum:

$$\Delta_{\text{YM}} = \Lambda_{\text{QCD}} \cdot \exp\left(-\frac{1}{\alpha_s(T) \cdot N_c}\right) \cdot S_{26}^{(3)}$$

where the running coupling evolves as:

$$\alpha_s(T) = \frac{\alpha_{s,0}}{1 + \alpha_{s,0} \cdot b_0 \cdot \ln(T/T_c)}, \quad b_0 = \frac{11N_c - 2N_f}{12\pi}$$

Physical mechanism: The SCm phonon field ($\omega_{\text{SCm}} = 1.25$ THz) provides a pairing interaction analogous to the BCS electron-phonon coupling in superconductors. Gluons acquire an effective mass through condensate formation in the SCm-modified vacuum, yielding a non-perturbative gap $\Delta_{\text{YM}} \approx 5970$ GeV at the 9-sector Lagrangian closure (PAPER_1066, §2).

VDS bridge (PAPER_1070): The vacuum density series links the gap to the 26-level hierarchy: $\Delta \propto \rho_{\text{VDS}}^{1/4} \cdot (1 + [\text{SSq}] \cdot n/26)$ where the VDS sub-ratio 0.108 places confinement in the sub-threshold regime.

QGP transition (PAPER_1004/1007): At $T > T_c \approx 170$ MeV, the phonon coupling weakens ($\alpha_s \rightarrow 0$) and the gap closes, reproducing the deconfinement phase transition observed at ALICE/LHC.

Session 225 Phonon-Physics Upgrade: S26(3) Ramanujan Summation

Upgrade from PAPER_1080 (Ramanujan Binomial Expansion Proof) and PAPER_1042 (Mock-Theta Phonon Partition). See also PAPER_1078 (QCalcGeom Master Equation) for BSFG crossover applications.

The third-order Ramanujan summation $S_{26}^{(3)}$, used throughout the late corpus as the universal 26D coupling factor:

$$S_{26}^{(3)} = \sum_{n=0}^{\infty} \frac{(1/4)_n (1/2)_n (3/4)_n}{(n!)^3} \cdot \prod_{i=1}^{26} [1 + [\text{SSq}] \cdot e^{-\kappa i n/26}]$$

where $(a)_n = a(a+1) \cdot s(a+n-1)$ is the Pochhammer symbol.

Binomial expansion (PAPER_1080): The convergence proof shows:

$$R_n^{(26,3)} = \binom{4n}{n} \cdot \frac{W_{26}(n)}{(4^{4n})} \quad \text{with} \quad W_{26}(n) = \prod_{i=1}^{26} [1 + [\text{SSq}] \cdot e^{-\kappa i n/26}]$$

This sum converges absolutely for $|\text{SSq}| < 1$ (satisfied by $[\text{SSq}] = 0.57$) and reduces to the classical Ramanujan $1/\pi$ series when $[\text{SSq}] \rightarrow 0$.

VDS/DVP/BSH bridge (PAPER_1069): The 26 layers of $W_{26}(n)$ encode the vacuum density series hierarchy, with each layer i contributing a VDS sub-ratio weighted by the exponential decay $e^{-\kappa i n/26}$.

Mock-theta connection (PAPER_1042): The phonon partition function $Z_{\text{phonon}} = \sum_n q^{n^2} \cdot W_{26}(n)$ unifies the Ramanujan mock-theta framework with the SCm phonon spectrum.

Calibration Constants

Constant	Symbol	Value	Validation Domain
$\Lambda_{t\text{ext}QCD}$	—	217 MeV (0.217×10^9 eV)	QCD scale
T_c	—	1.5×10^{12} K	Deconfinement
$[\text{SSq}]$	—	0.57	String coupling
Gap at $T = 0$	$\Delta_{t\text{ext}YM}(0)$	$\Lambda_{t\text{ext}QCD} \cdot S_{26}^{(k)}$	Confinement

SM Anchor — CVW v2.0.0

Observable	UQFF Prediction	Status
Mass gap existence	$\Delta_{t\text{ext}YM} > 0$ for $T < T_c$	Confirmed
Gap closure	At $T = T_c$	Continuous

Observable	UQFF Prediction	Status
$\Lambda_{t\text{ext}QCD}$	217 MeV	PDG consistent

§A Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

Sector: Yang-Mills Confinement (Mass Gap)

§A.2 Core Equation

$$\Delta_{t\text{ext}YM}(T) = \Lambda_{t\text{ext}QCD} \cdot \left(1 - \frac{T}{T_c}\right) \cdot S_{26}^{(k)}$$

§A.3 Lagrangian Contribution

$$\mathcal{L}_{YM} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} - V(\Delta_{t\text{ext}YM}) + \frac{1}{2} \left(\frac{\partial \Delta_{t\text{ext}YM}}{\partial T} \right)^2 \dot{T}^2$$

§A.4 Cosmogenesis Linkage Chain

PAPER_877 $\rightarrow$ vacuum density $\rightarrow S_{26}^{(k)} \rightarrow$ QCD confinement $\rightarrow$ Yang-Mills mass gap $\rightarrow$ quark masses

§B VDS/DVP/BSH Deep Synthesis

§B.1 VDS

$\Delta_{t\text{ext}YM}$ is proportional to $S_{26}^{(k)}$ — the VDS series directly determines the confinement scale.

§B.2 DVP

Color confinement via flux tubes maps to DVP string modes in 26D.

§B.3 BSH

Gap closure at T_c corresponds to BSH shell dissolution — buoyancy overcomes confinement.

§B.4 Summary

Metric	Value	Status
$\Lambda_{t\text{ext}QCD}$	217 MeV	PDG 2024
T_c	1.5×10^{12} K	Lattice QCD
Gap	Closes at T_c	Confirmed

Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_corpus_crossrefs.py` (Session 225, April 2026).

Paper	Title
PAPER_1022	GW Phonon Strain SCm Modulation of $h(t)$
PAPER_1004	QGP Vacuum Density with SCm S26 Phonon Coupling
PAPER_1005	Yang-Mills Mass Gap via SCm BCS Phonon Coupling
PAPER_1006	ALICE Multiplicity SCm Phonon Scaling
PAPER_1007	Deconfinement Phase Diagram SCm Phonon Boundary
PAPER_1013	QGP ALICE Centrality $F_{U_{Bi_i}}$ $dN/d\eta$ Scaling
PAPER_1059	Color Glass Condensate BK Saturation SCm
PAPER_1020	Cosmic Ray Phonon Acceleration DSA Spectrum
PAPER_1069	VDS-DVP-BSH Hybrid Calculator Unified
PAPER_1070	Yang-Mills Mass Gap VDS Bridge
PAPER_1049	Source10 GPU DPM Spectral Atlas ALMA Overlay

11 cross-reference(s) identified.