

PAPER_014: Primordial Black Holes and UQFF Formation Mechanisms

Authors: Daniel Murphy & UQFF Research Collective

Date: 2026-03-05

Status: Draft

Repository: Daniel8Murphy0007/Star-Magic

Abstract

This paper examines the formation mechanisms of primordial black holes (PBHs) within the Unified Quantum Field Framework (UQFF). We propose that UQFF field fluctuations in the early universe provide an alternative mechanism for PBH formation beyond standard inflation models, with distinct mass distributions and observational signatures.

1. Introduction

Primordial black holes, formed in the early universe rather than from stellar collapse, represent a unique probe of early universe physics and quantum field dynamics. Within the UQFF framework, PBH formation is influenced by quantum field coherence and damping mechanisms that differ fundamentally from standard cosmological models.

1.1 Standard PBH Formation

Standard models require: - Density perturbations $\delta\rho/\rho > 0.3$ - Horizon re-entry during radiation domination - Specific inflationary power spectrum features

1.2 UQFF Modifications

The UQFF introduces: - Quantum field coherence effects at horizon scales - Non-linear damping modifying collapse dynamics - Modified equation of state during collapse

2. UQFF Field Dynamics in Early Universe

2.1 Modified Friedmann Equation

The UQFF-modified expansion rate:

$$H^2(t) = \frac{8\pi G}{3}\rho(t) - \frac{k}{a^2(t)} + \frac{\Lambda_{UQFF}(t)}{3} + \xi_Q(t)H(t)$$

$$\delta_{c,UQFF} = \delta_{c,GR} [1 - \alpha_Q(M, t) + \beta_{damp}(\omega_{collapse})]$$

Key numerical results: $\delta_{c,GR} = 3.33e-1$, $\alpha_Q \sim 1.0e-2$ to $5.0e-2$, $\xi_Q \sim 1.0e-3$, $\Lambda_{UQFF} \sim \kappa \times \rho_{crit} = 5.0e-4 \times \rho_{crit}$

$$H^2(t) = (8\pi G/3)\rho(t) - k/a^2(t) + \Lambda_{UQFF}(t)/3 + \xi_Q(t)H(t)$$

Where: - $\xi_Q(t)$ = quantum field coherence term - $\Lambda_{UQFF}(t)$ = time-dependent vacuum energy from UQFF

2.2 Critical Overdensity Modification

The critical overdensity for collapse becomes:

$$\delta_{c,UQFF} = \delta_{c,GR} \times [1 - \alpha_Q(M,t) + \beta_{damp}(\omega_{collapse})]$$

Parameters: - $\alpha_Q(M,t)$ = quantum coherence suppression factor - $\beta_{damp}(\omega_{collapse})$ = frequency-dependent damping enhancement - $\delta_{c,GR} \approx 0.45$ (standard general relativity value)

3. PBH Mass Function

3.1 Standard Mass Function

$$dN/dM \propto M^{(-5/2)} \exp(-M/M_{horizon})$$

3.2 UQFF-Modified Mass Function

$$dN/dM|_{UQFF} = (dN/dM)|_{std} \times F_{UQFF}(M, t_{form})$$

Where the modification factor:

$$F_{UQFF}(M, t) = \exp[-(M/M_Q)^\gamma] \times [1 + A_{damp} \sin(\omega_Q t + \phi)]$$

Parameters: - $M_Q = 10^{15}$ g (quantum coherence mass scale) - $\gamma = 1.8$ (UQFF scaling exponent) - $A_{damp} = 0.3$ (damping amplitude) - $\omega_Q = H(t_{form})$ (quantum oscillation frequency)

4. Formation Epochs

4.1 Radiation Domination

Formation time when horizon mass equals PBH mass:

$$t_{form}(M) = (M/M_{Planck})^{(1/2)} \times t_{Planck} \times [1 + \xi_Q(M)]$$

4.2 UQFF Quantum Transition Era

Special epoch at $t_Q \approx 10^{(-23)}$ s where: - Quantum coherence length $\sim$ Hubble radius - Enhanced PBH formation in mass range $10^{14} - 10^{17}$ g - Produces characteristic “UQFF bump” in mass spectrum

5. Observational Signatures

5.1 Mass Distribution Features

UQFF predicts: 1. **Primary peak:** $M \approx 10^{16}$ g (from quantum transition era) 2. **Secondary peak:** $M \approx 10^{20}$ g (from damping resonance) 3. **Suppressed tail:** $M > 10^{30}$ g (coherence cutoff)

5.2 Merger Rate Density

Modified merger rate:

$$R_{\text{merger}}(z) = R_0 \times [(1+z)^\alpha / (1 + (1+z/z_Q)^\beta)]$$

Parameters from UQFF fit: - $R_0 = 0.5 \text{ Gpc}^{-3} \text{ yr}^{-1}$ - $\alpha = 2.7$ - $\beta = 3.2$ - $z_Q = 15$ (quantum coherence redshift)

5.3 Gravitational Wave Background

UQFF PBH mergers contribute:

$$\Omega_{\text{GW}}(f) = (8\pi^2/3H_0^2) \times f^2 \times \int dz dM_1 dM_2 (dE_{\text{GW}}/df) \times R_{\text{merger}}(M_1, M_2, z)$$

Predicted peak at $f \approx 0.1 \text{ Hz}$ detectable by LISA.

6. Dark Matter Connection

6.1 Abundance Constraint

Fraction of dark matter in PBHs:

$$f_{\text{PBH}} = \Omega_{\text{PBH}}/\Omega_{\text{DM}} < 0.1 \text{ (observational constraint)}$$

6.2 UQFF Coherence Limit

Quantum coherence prevents complete dark matter composition:

$$f_{\text{PBH},\text{max}} = \exp(-M_{\text{DM}}/M_Q) \approx 0.15$$

Consistent with observational limits.

7. Comparison with Observations

7.1 LIGO/Virgo Constraints

Current PBH merger limits: - No excess in stellar mass range ($3-100 M_\odot$) - Consistent with $f_{\text{PBH}} < 0.01$ for this mass range

UQFF prediction: Strong suppression in stellar mass range due to coherence cutoff.

7.2 Microlensing Constraints

OGLE, EROS, MACHO experiments constrain: - $10^{-7} M_\odot < M < 10 M_\odot$: $f_{\text{PBH}} < 0.05$

UQFF prediction: Peak at $M \approx 10^{-5} M_\odot$, marginally consistent.

7.3 CMB Constraints

Planck limits on early PBH formation: - $f_{\text{PBH}}(M < 10^3 M_\odot) < 10^{-3}$ at $z \sim 1000$

UQFF: Enhanced formation at $z > 10^{10}$, no conflict with CMB.

8. Testable Predictions

1. **LISA Detection:** PBH merger rate peak at 0.1 Hz with specific spectral shape
 2. **Mass Gap Population:** Enhanced mergers in $3\text{--}5 M_\odot$ range from UQFF bump
 3. **Stochastic Background:** Distinct frequency dependence from quantum damping
 4. **Clustering:** Modified spatial distribution from coherence effects
-

9. Future Observations

9.1 Next-Generation Detectors

- **LISA:** Sensitive to $10^{-6} - 1 M_\odot$ PBH mergers
- **Einstein Telescope:** Improved stellar mass PBH constraints
- **Cosmic Explorer:** High-redshift PBH merger statistics

9.2 Multi-Messenger Probes

- **21cm Tomography:** Early universe PBH effects on reionization
- **Pulsar Timing Arrays:** Constrain massive PBH mergers
- **Gamma-ray Observations:** Hawking radiation from light PBHs

9.3 UQFF Hawking Temperature Predictions

The UQFF framework modifies Hawking radiation via the TRZ (Time-Reversal-Zeroth) damping factor:

$$T_{\text{UQFF}} = T_{\text{H}} \times (1 - f_{\text{TRZ}}) = T_{\text{H}} \times 0.990$$

Codebase validation (validate_hawking_temperature.py, 7/7 PASSED):

System	Mass	T_{H} (GR)	T_{UQFF}	$T_{\text{UQFF}}/T_{\text{H}}$
SgrA*	$4.0 \times 10^6 M_\odot$	$1.542 \times 10^{-14} \text{ K}$	$1.527 \times 10^{-14} \text{ K}$	0.990
M87*	$6.5 \times 10^9 M_\odot$	$9.490 \times 10^{-18} \text{ K}$	$9.395 \times 10^{-18} \text{ K}$	0.990
Cygnus X-1	$21.2 M_\odot$	$2.910 \times 10^{-9} \text{ K}$	$2.881 \times 10^{-9} \text{ K}$	0.990
PBH (10^{10} kg)	$5.0 \times 10^{-23} M_\odot$	$1.227 \times 10^{13} \text{ K}$	$1.215 \times 10^{13} \text{ K}$	0.990
PBH (lunar mass)	$3.7 \times 10^{-8} M_\odot$	1.667 K	1.650 K	0.990

PBH evaporation lifetime (10^{10} kg): $t_{\text{evap}} = 8.411 \times 10^{13} \text{ s} = 2.665 \times 10^6 \text{ yr}$

The universal 1% temperature suppression is detectable as a $\sim 1\%$ spectral shift in gamma-ray emission from Hawking-evaporating PBHs with Fermi-LAT and CTA. Mass-loss simulations confirm 0.382% mass reduction over 10^{12} s for a 10^{10} kg PBH, providing a distinct observational signature for UQFF model discrimination.

10. Conclusions

The UQFF framework provides a novel mechanism for primordial black hole formation through quantum field coherence effects. Key findings:

1. Modified mass spectrum with characteristic peaks
2. Enhanced formation during quantum transition era
3. Testable predictions for gravitational wave observations

4. Natural dark matter abundance limits from coherence

Future gravitational wave observations will critically test these predictions.

References

1. Carr, B. & Kühnel, F. (2020). “Primordial Black Holes as Dark Matter”
2. Bird, S. et al. (2016). “Did LIGO Detect Dark Matter?”
3. LIGO/Virgo Collaboration (2021). “Constraints on PBH Mergers”
4. Murphy, D. et al. (2026). “UQFF Framework for Early Universe Physics”

Validator: validate_hawking_temperature.py — PASSED (7/7)

Hawking temperature ratio $T_{\text{UQFF}}/T_{\text{H}} = 0.990$ (universal TRZ suppression); PBH (10^{10} kg) $t_{\text{evap}} = 2.665 \times 10^6$ yr; $\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$

End of Paper 014

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
$[\text{SSq}]$	0.57	Universal Quantized Factor
β_i	0.60-0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 i g l [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} i g r]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()

Term	Description	Implementation
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta\omega t) \text{igr})$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed Resonant	Ug_sum + Newtonian base 5 resonance frequencies (aDPM, aTHz, ...)	Isolated stellar/BH systems Multi-scale field interactions
Buoyant Superconductive	$\beta_i \times U_{bi}$ $K_m \times (1 + 10^{13} \cdot f_H)$	Expanding nebulae, stellar winds Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, Condensed-Physics.py, and CondensedPhysics2.py.