

PAPER_038: UQFF Buoyancy Proof Variants 711: Fermi Acceleration, Cosmic Ray Knee, WHIM Temperature, Press-Schechter Halos, and Star Formation Efficiency

Title: UQFF Buoyancy Proof Variants 711: Fermi Acceleration, Cosmic Ray Knee, WHIM Temperature, Press-Schechter Halos, and Star Formation Efficiency

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Grok Thread: 98b2e77dfbc34d27b09f19fa7c460624

Validator: BuoyancyProofVariants.py All 17 variants operational ?

Variants: fermi, kne, whim, ps, sfe

Index Slot: 1.5 Buoyancy Proofs,

Title: UQFF Buoyancy Proof Variants 711: Fermi Acceleration, Cosmic Ray Knee, WHIM Temperature, Press-Schechter Halos, and Star Formation Efficiency

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Grok Thread: 98b2e77dfbc34d27b09f19fa7c460624

Validator: BuoyancyProofVariants.py All 17 variants operational ?

Variants: fermi, kne, whim, ps, sfe

Index Slot: 1.5 Buoyancy Proofs, PAPER_038

Abstract

This paper presents five F_UBii buoyancy proof variants addressing quantum corrections to macroscopic astrophysical processes. Variant 7 (fermi) derives the UQFF buoyancy of Fermi-accelerated particles at astrophysical shock fronts. Variant 8 (kne) applies the framework to the cosmic ray knee at $\sim 3 \times 10^5$ eV where the spectral index changes the UQFF predicts this spectral break as a phase transition in the F_UBii landscape. Variant 9 (whim) addresses the Warm-Hot Intergalactic Medium at $T \sim 10^5 \times 10^7$ K containing 4050% of cosmic baryons. Variant 10 (ps) maps the Press-Schechter dark matter halo mass function to a UQFF buoyancy force landscape. Variant 11 (sfe) quantifies the UQFF buoyancy contribution to star formation efficiency suppression in molecular clouds. Together these form the quantum corrections series, where small-scale quantum physics drives large-scale structure.

UQFF Discovery: Novel application of UQFF calibration constants ($\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Variant 7: Fermi Acceleration Buoyancy (fermi)

1.1 Physical Context

First-order Fermi (DSA) acceleration at shock fronts produces power-law particle spectra $N(E) \propto E^{-s}$ with $s = (r+2)/(r-1)$ for compression ratio r . The UQFF buoyancy arises because accelerated particles develop a pressure gradient against the surrounding thermal plasma.

Key systems: Tycho SNR ($v_{\text{shock}} \sim 4500$ km/s), Centaurus A jet shocks, Cygnus A hotspots

1.2 $F_{\text{UBii_fermi}}$ Equation

$$F_{\text{UBii,fermi}} = F_{\text{rel}} \cdot \frac{\beta_{\text{shock}} \cdot E_p}{E_{\text{LEP}}} \cdot Q_{\text{wave}} \cdot \left(\frac{v_{\text{shock}}}{c} \right)^2$$

where: - κ_{shock} = shock compression ratio (typical 37 for strong shocks) - E_p = particle energy (J) - v_{shock} = shock velocity (m/s)

1.3 (v/c) Relativistic Correction

The Fermi buoyancy scales as (v_{shock}/c) a relativistic correction that becomes important for $v_{\text{shock}} > 0.1c$. At non-relativistic shock speeds ($v_{\text{shock}}/c \sim 0.01$ for typical SNRs), the UQFF Fermi buoyancy is suppressed by $(0.01)^2 = 10^{-4}$ relative to the full E_p contribution.

1.4 Example: Centaurus A Jet Hotspot

For Cen A hotspot: $\kappa_{\text{shock}} = 4$ (strong shock), $E_p = 10^{-10}$ J (10 GeV proton), $v_{\text{shock}} = 0.5c$, $Q_{\text{wave}} = 1.0$:

$$F_{\text{UBii,fermi}}^{\text{CenA}} = 10^{-10} \times \frac{4 \times 10^{-9}}{1.22 \times 10^{-19}} \times (0.5)^2 = 10^{-10} \times 3.28 \times 10^{10} \times 0.25 = 8.2 \times 10^0 = 0.82 \text{ N}$$

The per-particle Fermi buoyancy force of 0.82 N per 10 GeV proton, scaled over the $\sim 10^6$ protons in the hotspot, gives the collective Fermi acceleration pressure maintaining the Cen A jet head.

2. Variant 8: Cosmic Ray Knee Energy Buoyancy (kne)

2.1 Physical Context

The cosmic ray energy spectrum follows $E^{-2.7}$ power law up to the “knee” at $\sim 3 \times 10^{15}$ eV, where it steepens to $E^{-3.1}$. The knee marks the maximum energy achievable by Galactic SNR shock acceleration for protons ($Z=1$). For heavy nuclei, the knee scales as $Z \cdot E_{\text{knee}}(\text{proton})$ the “knee composition model”.

2.2 $F_{\text{UBii_kne}}$ Equation

$$F_{\text{UBii,kne}} = -F_{\text{rel}} \cdot \frac{E_{\text{knee}}}{E_{\text{GUT}}} \cdot \frac{Z \cdot e}{E_{\text{LEP}}} \cdot Q_{\text{wave}} \cdot \ln \left(\frac{E_{\text{knee}}}{E_{\text{LEP}}} \right)$$

where: - E_{knee} = knee energy (J) 4.8×10^{-4} J (3×10^{15} eV) - E_{GUT} = GUT energy scale = 1.6×10^{-5} J ($\sim 10^6$ GeV) - Z = charge number of CR nucleus - $e = 1.602 \times 10^{-19}$ C - $E_{\text{LEP}} = 1.22 \times 10^{-19}$ J

The negative sign indicates spectral suppression above the knee, CR buoyancy forces prevent further acceleration.

2.3 UQFF Knee as Phase Transition

The $\ln(E_{\text{knee}}/E_{\text{LEP}})$ factor:

$$\ln \left(\frac{4.8 \times 10^{-4}}{1.22 \times 10^{-19}} \right) = \ln(3.93 \times 10^{15}) = 35.9$$

This logarithm attains its maximum precisely at E_{knee} for protons a UQFF prediction that the knee is not an arbitrary cutoff but a **stationary point** of the F_{UBii} landscape where:

$$\frac{\partial F_{\text{UBii,kne}}}{\partial \ln E} = 0 \Rightarrow E = E_{\text{knee}}$$

2.4 UQFF Knee Prediction: Proton vs Iron

For $Z=1$ (proton): $E_{\text{knee}} = 3 \times 10^5 \text{ eV} = 4.8 \times 10^{-4} \text{ J}$ For $Z=26$ (iron): $E_{\text{knee}}^{\text{Fe}} = 26 \times 3 \times 10^5 \text{ eV} = 7.8 \times 10^6 \text{ eV} = 1.25 \times 10^{-12} \text{ J}$

UQFF prediction for iron knee:

$$F_{\text{UBii,kne}}^{\text{Fe}} / F_{\text{UBii,kne}}^p = 26 \times \frac{\ln(1.25 \times 10^{-12} / 1.22 \times 10^{-19})}{\ln(4.8 \times 10^{-4} / 1.22 \times 10^{-19})} = 26 \times \frac{38.0}{35.9} = 27.5$$

The UQFF predicts the iron knee force is 27.5 the proton knee force (vs 26 in the pure rigidity model), a 5.8% enhancement from the quantum logarithmic correction.

3. Variant 9: WHIM Temperature Buoyancy (whim)

3.1 Physical Context

The Warm-Hot Intergalactic Medium (WHIM) at $z < 1$ contains 40-50% of all baryons in the Universe the “missing baryon problem” solution. It fills the cosmic web filaments at $T \sim 10^5 - 10^7 \text{ K}$, traced by O VI (105.6 nm), O VII (21.6 nm), and soft X-ray emission. Its buoyancy against the cosmic gravitational potential maintains the filamentary structure of the Universe.

3.2 $F_{\text{UBii_whim}}$ Equation

$$F_{\text{UBii,whim}} = F_{\text{rel}} \cdot \frac{k_B T_{\text{WHIM}}}{E_{\text{LEP}}} \cdot n_b \sigma_T r_{\text{fil}} \cdot Q_{\text{wave}} \cdot \sqrt{\frac{T_{\text{WHIM}}}{T_{\text{virial}}}}$$

where: - T_{WHIM} = WHIM temperature (K) - n_b = baryon number density (m⁻³) - σ_T = Thomson cross-section = $6.652 \times 10^{-29} \text{ m}^2$ - r_{fil} = filament radius (m) - T_{virial} = virial temperature of host structure (K)

3.3 $T^{(3/2)}$ Scaling

The WHIM buoyancy scales as $T_{\text{WHIM}}^{(3/2)} / (T_{\text{virial}}^{(1/2)})$: - Factor 1: thermal pressure $k_B T_{\text{WHIM}}$ - Factor 2: Thomson opacity depth $n_b \sigma_T r_{\text{fil}}$ (free electron count cross-section) - Factor 3: $v(T_{\text{WHIM}}/T_{\text{virial}})$ buoyancy stability criterion (analogous to Schwarzschild stability for convection)

3.4 Example: Cosmic Web Filament (Sculptor Wall)

For a typical baryon-rich filament: $T_{\text{WHIM}} = 106 \text{ K}$, $n_b = 10 \text{ m}^{-3}$, $r_{\text{fil}} = 10 \text{ Mpc} = 3.09 \times 10^7 \text{ m}$, $T_{\text{virial}} = 107 \text{ K}$, $Q_{\text{wave}} = 1.0$:

$$F_{\text{whim}} = 10^{-10} \times \frac{1.381 \times 10^{-23} \times 10^6}{1.22 \times 10^{-19}} \times 10 \times 6.652 \times 10^{-29} \times 3.09 \times 10^{23} \times \sqrt{0.1}$$

$$= 10^{-10} \times 1.132 \times 10^2 \times 2.055 \times 10^{-4} \times 0.316 = 10^{-10} \times 7.36 \times 10^{-3} = 7.4 \times 10^{-13} \text{ N}$$

This tiny force per unit volume, integrated over the filament volume $V \sim (10 \text{ Mpc}) \sim 3 \times 10^7 \text{ m}$, gives $F_{\text{total}} \sim 105 \text{ N}$ the UQFF buoyancy pressure holding the cosmic web filament against gravitational collapse.

4. Variant 10: Press-Schechter Halo Mass Buoyancy (ps)

4.1 Physical Context

The Press-Schechter (PS) mass function predicts the comoving number density of dark matter halos:

$$\frac{dn}{d \ln M} = \sqrt{\frac{2}{\pi}} \frac{\rho_0}{M} \frac{\delta_c}{\sigma} \left| \frac{d \ln \sigma}{d \ln M} \right| \exp \left(-\frac{\delta_c^2}{2\sigma^2} \right)$$

where $d_c = 1.686$ is the critical overdensity for spherical collapse. The UQFF buoyancy force analog maps this statistical distribution to a physical force.

4.2 $F_{\text{UBii,ps}}$ Equation

$$F_{\text{UBii,ps}} = -F_{\text{rel}} \cdot \frac{M_{\text{halo}}}{M_P^2} \cdot \frac{\delta_c}{E_{\text{LEP}}} \cdot Q_{\text{wave}} \cdot \left(-\frac{d \ln \sigma}{d \ln M} \right)$$

where: - M_{halo} = halo mass (kg) - M_P = Planck mass = $2.176 \times 10^{-8} \text{ kg}$ - $d_c = 1.686$ (critical overdensity) - s = RMS density fluctuation

4.3 Planck Mass Normalization

The M_{halo}/M_P normalization is the UQFF quantum gravity anchor halo masses are measured in units of M_P (the fundamental quantum gravity area unit). For a cluster-mass halo $m_{\text{halo}} \sim 10^5 M_\odot$:

$$\frac{M_{\text{halo}}}{M_P^2} = \frac{10^{15} \times 1.989 \times 10^{30}}{(2.176 \times 10^{-8})^2} = \frac{1.989 \times 10^{45}}{4.74 \times 10^{-16}} = 4.2 \times 10^{60} \text{ m}^{-1} \cdot \text{kg}$$

This enormous ratio reflects how macroscopic halo gravitational physics emerges from quantum Planck-scale foundations a direct UQFF bridge between cosmological structure formation and quantum gravity.

4.4 Example: Milky Way Halo

For Milky Way: $M_{\text{halo}} = 10^5 M_\odot = 1.989 \times 10^4 \text{ kg}$, $s(M_{\text{MW}}) \sim 0.5$, $d \ln s / d \ln M \sim -0.15$, $Q_{\text{wave}} = 1.0$:

$$F_{\text{ps}}^{MW} = -10^{-10} \times 4.2 \times 10^{60} \times \frac{1.686}{1.22 \times 10^{-19}} \times 1.0 \times 0.15 = -10^{-10} \times 4.2 \times 10^{60} \times 1.38 \times 10^{19} \times 0.15 = -$$

5. Variant 11: Star Formation Efficiency Buoyancy (sfe)

5.1 Physical Context

Star formation efficiency $e_{\text{SFE}} = M_{\text{star}}/M_{\text{gas}}$ ranges from $\sim 1\%$ in diffuse GMCs to $\sim 3050\%$ in dense molecular cloud cores. The UQFF buoyancy determines whether turbulence (e_{SFE} low) or gravity (e_{SFE} high) dominates in a given cloud.

5.2 $F_{\text{UBii_sfe}}$ Equation

$$F_{\text{UBii,sfe}} = F_{\text{rel}} \cdot \frac{\varepsilon_{\text{SFE}} \cdot M_{\text{gas}} \cdot c^2}{r_{\text{cloud}}^2 \cdot E_{\text{LEP}}} \cdot Q_{\text{wave}} \cdot \sqrt{\varepsilon_{\text{SFE}}}$$

5.3 Rest-Mass Energy Term

The $M_{\text{gas}} c$ term is unusual in a cloud-physics context it represents the UQFF's claim that star formation efficiency is ultimately limited by the rest-mass energy of the gas, mediated through the vacuum [SCm] manifold. The effective force is:

$$F_{\text{UBii,sfe}} \sim \frac{\varepsilon^{3/2} M_{\text{gas}} c^2}{r_{\text{cloud}}^2}$$

This is the UQFF prediction that star formation is a quantum-gravitational process limited by the ratio of gas rest-mass energy to the cloud surface (Bekenstein-like area scaling).

5.4 Example: Orion A Giant Molecular Cloud

For Orion A GMC: $e_{\text{SFE}} = 0.05$, $M_{\text{gas}} = 100 M_{\odot} = 1.989 \times 10^5 M_{\odot}$, $r_{\text{cloud}} = 10 \text{ pc} = 3.086 \times 10^7 \text{ m}$, $Q_{\text{wave}} = 1.0$:

$$\begin{aligned} F_{\text{sfe}}^{\text{Orion A}} &= 10^{-10} \times \frac{0.05 \times 1.989 \times 10^5 \times (3 \times 10^8)^2}{(3.086 \times 10^7)^2 \times 1.22 \times 10^{-19}} \times \sqrt{0.05} \\ &= 10^{-10} \times \frac{0.05 \times 1.79 \times 10^{49}}{9.52 \times 10^{34} \times 1.22 \times 10^{-19}} \times 0.224 = 10^{-10} \times 7.68 \times 10^{31} \times 0.224 = 1.72 \times 10^{22} \text{ N} \end{aligned}$$

6. Summary: Quantum Corrections Series

Variant	Physical Context	Key Formula	Characteristic Scale
fermi	Fermi acceleration	$\kappa_{\text{shock}} - E_{\text{p}}$ (v/c)	$\sim 0.8 \text{ N}$ per 10 GeV proton
kne	CR knee at $3 \times 10^5 \text{ eV}$	$E_{\text{knee}}/E_{\text{GUT}} - Z e \ln(E/E_{\text{LEP}})$	Knee as F_{UBii} stationary pt
whim	Cosmic baryon reservoir	$k_{\text{BT}} n_{\text{b}} s_{\text{T}} r_{\text{fil}}$ $v(T/T_{\text{vir}})$	$\sim 105? \text{ N}$ per filament
ps	PS halo mass function	$M_{\text{halo}}/M_{\text{P}} d_c$	$d \ln s / d \ln M$
sfe	Molecular cloud SFR	$e^{(3/2)} M_{\text{gas}} c/r$	$\sim 10 \text{ N}$ (Orion A)

Conclusions

Variants 711 demonstrate that the UQFF buoyancy framework provides quantitative predictions across the full range of quantum-to-cosmic scales:

1. **fermi:** UQFF Fermi buoyancy provides the back-pressure that terminates DSA acceleration at the particle energy where $F_UBii_fermi = F_confinement$
2. **kne:** CR knee is the stationary point of F_UBii_kne a UQFF phase transition rather than a diffusive escape threshold
3. **whim:** WHIM buoyancy $F_UBii_whim \sim 105? N$ per filament maintains cosmic web structure against gravitational collapse
4. **ps:** PS halo formation maps to Planck-mass-normalized UQFF collapse force a quantum gravity bridge to large-scale structure
5. **sfe:** Star formation efficiency is UQFF-limited by rest-mass energy Bekenstein-area scaling: $F_UBii_sfe \propto e^{(3/2)} M_gas c/r$

Validator: *BuoyancyProofVariants.py* ? All 17 F_UBii variants operational ? | ? = 0.0005/day | $[SSq] = 0.57$

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60-0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_react(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{react}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()

Term	Description	Implementation
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma \lambda_i \cdot U_i \cdot E_{\text{react4th}}$	dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{\text{SCm}} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times \text{Ubi}$	Expanding nebulae, stellar winds
Superconductive	$\text{Um} \times (1+10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.