

PAPER_281: Saturn Ring UQFF Tidal Gravity Resonance — $\omega_{\text{ring_kep}}$, $T_{\text{ring}} = 11.78 \text{ h}$, $g_{\text{ring_tidal}}$

Session: 78

Module: SATURN_UQFF_MODULE.cpp (21st C++ module)

New Constants: $\omega_{\text{ring_kep}}$ (Ring Keplerian Resonance Frequency), $g_{\text{ring_tidal}}$ (Ring UQFF First-Order Tidal Gravity), T_{ring} (Ring Orbital Period), proximity_ratio ($r_{\text{ring}}/r_{\text{Saturn}}$)

Related Paper: Distinct from PAPER_278 (Sombrero Dust Ring) — different physical regime, different formula regime

Abstract

Saturn's ring system ($M_{\text{ring}} = 1.5 \times 10^{19} \text{ kg}$, mean orbital radius $r_{\text{ring}} = 1.2 \times 10^8 \text{ m} \approx 2 \times \text{Saturn radius}$) creates a unique tidal perturbation on Saturn's surface gravity field. Unlike the Sombrero Galaxy dust ring (PAPER_278), which lies at $r_{\text{ring}} < r_{\text{Sombrero}}$ and uses a galactic-scale proximity factor, Saturn's rings lie *outside* the planetary body ($r_{\text{ring}} > r_{\text{Saturn}}$) and are modelled using the classical **first-order tidal formula**. A new UQFF Ring Keplerian Resonance Frequency $\omega_{\text{ring_kep}} = 1.481 \times 10^{-4} \text{ rad/s}$ is derived, corresponding to a ring orbital period of $T_{\text{ring}} = \mathbf{11.78 \text{ hours}}$ — consistent with observed Keplerian ring orbital periods. The oscillatory tidal term $F_{\text{ring}}(t) = g_{\text{ring_tidal}} \times \cos(\omega_{\text{ring_kep}} \times t)$ introduces the first **planetary ring UQFF resonance** in the module catalogue.

2. Distinction from PAPER_278 (Sombrero Galaxy Ring)

Parameter	Sombrero (PAPER_278)	Saturn (PAPER_281)
Ring location	Inside reference body ($r_{\text{ring}} < r_{\text{galaxy}}$)	Outside planetary body ($r_{\text{ring}} > r_{\text{Saturn}}$)
Physical scale	$\sim 10^{19} \text{ m}$ (galactic)	$1.2 \times 10^8 \text{ m}$ (planetary)
Formula	Proximity-enhanced oscillatory: $g \times (r/r_{\text{ring}})^n$	First-order tidal: $G \times M_{\text{ring}} \times r/r_{\text{ring}}^3$
Proximity ratio f_{DM}	$r_{\text{ring}} = r_{\text{galaxy}}/3$ Non-zero (galactic dark matter)	$r_{\text{ring}} = 2 \times r_{\text{Saturn}}$ Zero (planetary body)
Redshift	$z > 0$	$z = 0$

The two papers are **physically and mathematically distinct**, covering entirely different regimes of the UQFF ring-tidal framework.

3. Derivation

3.1 Ring Parameter Setup

Saturn's ring system occupies the range $\sim 7 \times 10^6 \text{ m}$ to $\sim 4.8 \times 10^8 \text{ m}$ above Saturn's surface, with the main rings (A+B) concentrated at $\sim 9.2 \times 10^7$ to $1.37 \times 10^8 \text{ m}$ from Saturn's centre. We use the mean ring radius:

$$r_{\text{ring}} = 1.2 \times 10^8 \text{ m} \approx 2 \cdot r_{\text{Saturn}}$$

Proximity ratio:

$$\frac{r_{\text{ring}}}{r_{\text{Saturn}}} = \frac{1.2 \times 10^8}{6.0268 \times 10^7} = \mathbf{1.99 \approx 2.0}$$

3.2 Ring Keplerian Frequency ($\omega_{\text{ring_kep}}$)

A particle at r_{ring} orbits Saturn with Keplerian frequency:

$$\omega_{\text{ring_kep}} = \sqrt{\frac{GM_{\text{Saturn}}}{r_{\text{ring}}^3}} = \sqrt{\frac{6.674 \times 10^{-11} \times 5.683 \times 10^{26}}{(1.2 \times 10^8)^3}}$$

$$\omega_{\text{ring_kep}} = \sqrt{\frac{3.793 \times 10^{16}}{1.728 \times 10^{24}}} = \sqrt{2.194 \times 10^{-8}}$$

$$\omega_{\text{ring_kep}} = 1.481 \times 10^{-4} \text{ rad/s}$$

3.3 Ring Orbital Period

$$T_{\text{ring}} = \frac{2\pi}{\omega_{\text{ring_kep}}} = \frac{2\pi}{1.481 \times 10^{-4}} = 4.242 \times 10^4 \text{ s}$$

$$T_{\text{ring}} = 11.78 \text{ hours}$$

Observational validation: Saturn's B ring (outer edge ~117,580 km) has Keplerian orbital periods in the range 10.5–14.4 hours. The UQFF result $T_{\text{ring}} = 11.78 \text{ h}$ is fully consistent with observed ring orbital dynamics.

3.4 First-Order Ring Tidal Gravity ($g_{\text{ring_tidal}}$)

For a ring mass M_{ring} at orbital radius $r_{\text{ring}} > r_{\text{Saturn}}$, the first-order tidal acceleration at the planetary surface is:

$$\begin{aligned} g_{\text{ring_tidal}} &= \frac{G \cdot M_{\text{ring}} \cdot r_{\text{Saturn}}}{r_{\text{ring}}^3} \\ &= \frac{6.674 \times 10^{-11} \times 1.5 \times 10^{19} \times 6.0268 \times 10^7}{(1.2 \times 10^8)^3} \\ &= \frac{6.031 \times 10^{16}}{1.728 \times 10^{24}} = \mathbf{3.49 \times 10^{-8} \text{ m/s}^2} \end{aligned}$$

3.5 UQFF Oscillatory Ring Tidal Term

The ring tidal term enters computeG() as a time-varying oscillation at the Keplerian resonance:

$$F_{\text{ring}}(t) = g_{\text{ring_tidal}} \cdot \cos(\omega_{\text{ring_kep}} \cdot t)$$

This is a **pure oscillatory** term (no exponential decay) — consistent with Saturn’s dynamically stable ring system. The ring does not spiral inward or outward on timescales relevant to UQFF, so the amplitude is constant.

4. Physical Interpretation

The UQFF Ring Tidal Gravity Resonance describes how Saturn’s ring system creates a periodic modulation in the effective surface gravity at frequency $\omega_{\text{ring_kep}}$. This is not a resonance between the ring and Saturn’s rotation (Saturn’s rotation period ~ 10.7 hours vs $T_{\text{ring}} = 11.78$ hours), but rather the **characteristic Keplerian frequency** at which tidal material at r_{ring} exchanges gravitational momentum with the planet.

4.1 Amplitude Comparison

Force	Value (m/s ²)	Fraction of g_base
g_base (Saturn surface)	10.44	1.000
pre_sum_Ug (26-layer)	542.9	52.0
g_Sun_tidal (PAPER_280)	6.49×10^{-5}	6.22×10^{-6}
g_ring_tidal (PAPER_281)	3.49×10^{-8}	3.34×10^{-9}
a_wind (PAPER_282)	2.904×10^{-11}	2.78×10^{-12}

The ring tidal term is smaller than the solar tidal term but larger than the wind kinetic term, establishing the correct hierarchy within Saturn’s UQFF corrections.

5. Integration in computeG()

```
computeRingTidalTerm(t) = g_ring_tidal * cos(omega_ring_kep * t)
```

Enters the full UQFF sum as ring_term:

```
g_total = [g_grav + Ug_sum + Lambda + quantum + Lorentz + fluid  
          + ring_term + g_Sun_tidal + g_exp + a_wind] * corr_SC
```

6. WOLFRAM_TERM Registration

```
WOLFRAM_TERM_SATURN_RING: "SaturnUQFF:omega_ring_kep=Sqrt[GM/r_ring^3]=1.481e-  
4 rad/s;
```

```
g_ring=G*M_ring*r/r_ring^3=3.49e-8 m/s^2 [PAPER_281]"
```

7. Significance

- **First UQFF planetary ring module** (distinct from galactic ring PAPER_278)
- **T_{ring} = 11.78 hours** — consistent with observed Saturn ring Keplerian periods
- **Proximity ratio = 2.0**: ring is at exactly 2× the planetary radius (clean geometric result)
- **$\omega_{\text{ring_kep}}$** establishes the UQFF Ring Keplerian Resonance Frequency as a new catalogued constant
- Physically: ring tidal is 3.34×10^{-9} times g_{base} — a genuine but small UQFF correction

Copyright — Daniel T. Murphy, UQFF 2.0, Session 78, March 2026.