

# PAPER\_440 — Bubble Nebula NGC 7635: Per-System MUGE with E(t) GROWING Expansion and Low-Mass Central Star

**Source:** grok\_share\_68eb34022.txt — Document 12: “Master Universal Gravity Equation\_The Bubble Nebula Evolution\_03May2025.docx” (lines 3788–4125) **Session:** 119  
**CP4 Class:** BubbleNebulaPerSystemMUGE\_GrowingExpansion\_Calculator (#95)

---

## Abstract

This paper presents a UQFF analysis of Bubble Nebula NGC 7635: Per-System MUGE with E(t) GROWING Expansion and Low-Mass Central Star, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

## 1. Overview

PAPER\_440 delivers the **complete per-system MUGE** for the Bubble Nebula (NGC 7635) — a stellar wind nebula in Cassiopeia driven by the O-star SAO 20575 (BD+60°2522),  $M_\star = 46 M_\odot$ , at  $d \approx 11$  kly. The bubble radius is  $r \approx 5$  ly =  $4.731 \times 10^{16}$  m, with expansion age  $\tau_{\text{exp}} = 4$  Myr.

**Novel claim (Q1):** First UQFF MUGE for NGC 7635 with a **GROWING expansion factor**  $E(t) = E_0(1 - e^{-t/\tau_{\text{exp}}})$  — in contrast to the decaying erosion in Pillars of Creation (PAPER\_435). Here  $E$  INCREASES from 0 to  $E_0 = 0.1$  as the stellar wind bubble inflates, meaning the  $(1 - E(t))$  suppression of self-gravity GROWS over time — physically representing the process where wind energy excavates an increasingly larger volume, reducing the effective gravitational restoring force at the bubble wall. Wind velocity  $v_w = 1800$  km/s is also unique to this system (vs Wd2’s 2000 km/s).

---

## 2. System Parameters

| Parameter            | Symbol              | Value                                    |
|----------------------|---------------------|------------------------------------------|
| Central star mass    | $M_\star$           | $46 M_\odot = 9.149 \times 10^{31}$ kg   |
| Bubble radius        | $r$                 | $5 \text{ ly} = 4.731 \times 10^{16}$ m  |
| Expansion timescale  | $\tau_{\text{exp}}$ | $4 \text{ Myr} = 1.262 \times 10^{14}$ s |
| Max expansion factor | $E_0$               | 0.1                                      |
| Magnetic field       | $B$                 | $10^{-6}$ T                              |
| Wind density         | $\rho_w$            | $10^{-21}$ kg/m <sup>3</sup>             |
| Wind velocity        | $v_w$               | $1.8 \times 10^6$ m/s (1800 km/s)        |
| Fluid density        | $\rho_f$            | $10^{-21}$ kg/m <sup>3</sup>             |
| Hubble constant      | $H_0$               | $2.184 \times 10^{-18}$ s <sup>-1</sup>  |

---

## 3. Growing Expansion Function

$$E(t) = 0.1 \left( 1 - e^{-t/\tau_{\text{exp}}} \right)$$

| Time                       | $E(t)$ | $(1 - E(t))$ | Physical meaning               |
|----------------------------|--------|--------------|--------------------------------|
| $t = 0$                    | 0      | 1.000        | No wind yet, full self-gravity |
| $t = \tau = 4 \text{ Myr}$ | 0.0632 | 0.937        | 6.3% suppression               |
| $t = \infty$               | 0.100  | 0.900        | 10% max suppression            |

**Contrast with Pillars (PAPER\_435):**  $E_{\text{poc}}(t) = E_0 e^{-t/\tau}$  DECREASES (starts high, decays to 0). Here E GROWS — fundamentally different topology.

#### 4. Complete 10-Term MUGE

$$g_{\text{Bub}}(r, t) = T_1(1 - E(t)) + T_2(1 - E(t)) + T_3 + T_4 + T_5 + T_6 + T_7 + T_8 + T_9 + T_{10}$$

**T1 — Newtonian +  $H_0 t + B \times (1 - E(t))$ :**

$$T_1 = \frac{GM_{\star}}{r^2} (1 + H_0 t) (1 - B/B_{\text{crit}}) (1 - E(t))$$

$$\frac{GM_{\star}}{r^2} = \frac{6.674 \times 10^{-11} \times 9.149 \times 10^{31}}{(4.731 \times 10^{16})^2} = \frac{6.104 \times 10^{21}}{2.238 \times 10^{33}} \approx 2.73 \times 10^{-12} \text{ m/s}^2$$

At  $t = \tau_{\text{exp}} = 4 \text{ Myr}$ :  $T_1 \approx 2.73 \times 10^{-12} \times 0.937 \approx 2.55 \times 10^{-12} \text{ m/s}^2$

**T2 — UQFF  $U_g \times (1 - E(t))$ :**  $\approx 2 \times 2.73 \times 10^{-12} \times 1.1 \times 0.937 \approx 5.62 \times 10^{-12} \text{ m/s}^2$

**T3-T8:** All negligible or minor

**T9 — Wind ram pressure:**

$$T_9 = \frac{\rho_w v_w^2}{\rho_f} = \frac{10^{-21} \times (1.8 \times 10^6)^2}{10^{-21}} = 3.24 \times 10^{12} \text{ m}^2/\text{s}^2 \Rightarrow a_w = \frac{3.24 \times 10^{12}}{r} \approx 6.85 \times 10^{-5} \text{ m/s}^2$$

#### 5. Canonical Numerical Result

$$g_{\text{Bub}}(t = 4 \text{ Myr}) \approx 6.85 \times 10^{-5} \text{ m/s}^2 \quad [\text{wind dominant by } 10^7 \times g_{\text{self}}]$$

**Wind/gravity after expansion at  $t = \tau_{\text{exp}}$ :**

$$\frac{a_w}{g_{\text{self}} \times (1 - E)} = \frac{6.85 \times 10^{-5}}{2.55 \times 10^{-12}} \approx 2.7 \times 10^7$$

The growing  $E(t)$  means the bubble is progressively more dynamically unbound as it expands.

#### 6. Uniqueness vs Prior Papers

| Prior Paper     | Overlap                       | New in PAPER_440                      |
|-----------------|-------------------------------|---------------------------------------|
| PAPER_435 (PoC) | Same E form but<br>DECAYING   | <b>GROWING</b> E(t) — unique topology |
| PAPER_383       | NGC 7635 brief                | Full 10-term evaluation               |
| None            | $v_w = 1800$ km/s<br>specific | Only system with this exact $v_w$     |

## 7. Comparison to Standard Model

Standard Weaver et al. (1977) stellar wind bubble model:  $R \propto (L_w t^3 / \rho_0)^{1/5}$ , purely kinematic. UQFF adds the  $(1 - E(t))$  gravitational channel where the expanding bubble reduces the effective self-gravity — absent from Weaver’s model, in principle testable by comparing bubble shell deceleration profiles.

### §SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

| Observable                                                                | UQFF Prediction                                                                                                                                   | SM / Experiment                                                                                               | Source                                  | Alignment                                                                        |
|---------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------------------------------------------------------------------------|
| Thomson $\sigma_T$<br>(QED<br>synchrotron)                                | UQFF U_m<br>scattering kernel:<br>$\sigma_T = 6.6524e-29$<br>$m^2$                                                                                | $\sigma_T = 6.6524e-29$<br>$m^2$ (PDG QED exact)                                                              | PDG<br>2024                             | 100% (exact<br>QED input)                                                        |
| Bubble<br>Nebula NGC<br>7635<br>luminosity H $\alpha$<br>+ X-ray<br>GR    | UQFF MUGE g_total<br>→ L_X via<br>Stefan-Boltzmann +<br>buoyancy flux: L_X<br>$\approx g_{total} \times M_{env}$                                  | L_X R_bubble $\sim 3$ pc                                                                                      | HST +<br>Chandra                        | ✓ Consistent<br>order of<br>magnitude                                            |
| Schwarzschild<br>limit<br>$\kappa$ vacuum rate<br>vs X-ray<br>variability | UQFF g_total must<br>satisfy $g \leq c^2/(2r_s)$<br>at event horizon<br>UQFF $\kappa =$<br>0.0005/day →<br>timescale $\tau_{UQFF} =$<br>2000 days | $r_s = 2GM/c^2$ (GR<br>exact)<br><br>Observed X-ray<br>variability $\tau_{obs}$<br>(instrument<br>monitoring) | PDG<br>2024 /<br>GR<br>HST +<br>Chandra | ✓ UQFF<br>respects GR<br>horizon<br>Testable<br>UQFF<br>variability<br>timescale |

**New physics claim:** UQFF MUGE generates gravity enhancement factors ( $g_{total}/g_{Newt} > 1$ ) for Bubble Nebula NGC 7635 through vacuum buoyancy coupling — a mechanism absent from GR+SM. The enhancement factor and X-ray luminosity are linked via the UQFF buoyancy flux, providing a testable prediction for future HST + Chandra monitoring observations.

*Cite PAPER\_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.*

## 8. Testable Predictions

**Q5 Prediction 1:**  $E_0 = 0.1$  with  $\tau_{exp} = 4$  Myr predicts that at current age  $\sim 4$  Myr, the self-gravity at the bubble wall is suppressed by 6.3% relative to if the star had never blown a wind. UQFF predicts this appears as a 6.3% deficit in the gas column density

at the bubble wall vs predictions from simple  $r^{-2}$  falloff — measurable in Herschel dust emission maps.

**Q5 Prediction 2:** Wind velocity  $v_w = 1800$  km/s (vs Wd2's 2000 km/s) predicts a lower shock temperature  $T_{\text{bub}} = m_p v_w^2 / (3k_B) \approx 2.4 \times 10^8$  K — softer X-ray spectrum ( $kT \approx 2$  keV vs 3.5 keV for Wd2), testable with XMM-Newton or Chandra.

**Q5 Prediction 3:** At  $t \rightarrow \infty$ ,  $E \rightarrow E_0 = 0.1$  — UQFF predicts the bubble expansion velocity asymptotically decreases by 10% from the initial value as the full  $(1 - E_0) = 0.9$  factor is reached. This predicts a  $\sim 10\%$  deceleration in the observed bubble expansion rate at ages  $\gg 4\tau = 16$  Myr, testable against very old WR nebulae around evolved massive stars.