

PAPER_084: Information Paradox Resolution in the 26D UQFF Framework via Holographic Channel Encoding

Session: 0

Title: Information Paradox Resolution in the 26D UQFF Framework via Holographic Channel Encoding

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: MAIN_1_CoAnQi.cpp Batch 21 (Information Paradox, 26D channels), validate_phase3.py

Index Slot: 1.11 Black Hole Physics & Hawking Radiation,

Title: Information Paradox Resolution in the 26D UQFF Framework via Holographic Channel Encoding

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: MAIN_1_CoAnQi.cpp Batch 21 (Information Paradox, 26D channels), validate_phase3.py

Index Slot: 1.11 Black Hole Physics & Hawking Radiation, PAPER_084

Abstract

The black hole information paradox where unitarity demands information escapes evaporation but Hawking radiation appears thermal is resolved in the UQFF through 26-dimensional holographic encoding. Each of the 26 UQFF spatial dimensions carries an independent quantum information channel. During evaporation, infalling information is redistributed across channels 126, with channels 2126 (the ultra-compact layers, beyond the observable 4D membrane) acting as non-local storage. The 4D observer measures a thermal Hawking spectrum, but the full 26D state is pure. Batch 21 (Jan 28, 2026) implemented the InformationParadoxModule with Hawking radiation Page curves and 26D channel encoding.

UQFF Discovery: Novel application of UQFF calibration constants ($\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Standard Paradox

Hawking (1976): if T_H is exactly thermal, S_{BH} decreases from $S=A/4$ to $S=0$? pureness is lost. Unitarity requires $S_{\text{entanglement}}$ (Hawking radiation with BH) to follow the Page curve: first increasing (early time), then decreasing back to 0 (late time).

2. 26D Holographic Channel Resolution

Channel Structure

The UQFF 26-layer compressed gravity framework assigns each dimensional layer a quantum channel capacity:

$$I_k = -\text{Tr}[\rho_k \log \rho_k] \quad k = 1, \dots, 26$$

Total information: $I_{\text{total}} = \sum_{k=1}^{26} I_k = S_{\text{BH,initial}}$

Channel Assignment During Evaporation

Layer Group	Channels	Storage Type	4D Observable
14	UQFF observable	Hawking photons	Thermal spectrum
518	UQFF extended	Sub-Planckian modes	Not observable
1924	UQFF deep structure	Non-local entanglement	Nil
2526	Cosmic Egg layers	Non-local pure state	Information preserved

Channels 2526 host the **complete pre-collapse pure state** throughout evaporation, ensuring global unitarity while channels 14 produce the observed thermal spectrum.

3. UQFF Page Curve

The UQFF Page curve modifies the standard quantum extremal surface (QES) prediction:

$$S_{\text{UQFF}}(t) = \min \left[S_{\text{thermal}}(t), S_{\text{BH}}(t) + \sum_{k=25}^{26} I_k (1 - e^{-\kappa t}) \right]$$

Where: - $S_{\text{thermal}}(t)$ = early-time entropy of Hawking radiation - $S_{\text{BH}}(t)$ = Bekenstein-Hawking entropy (decreasing) - The $e^{-\kappa t}$ term comes directly from the UQFF $\kappa = 0.0005/\text{day}$ decay - Page time t_P occurs when $S_{\text{thermal}} = S_{\text{BH}} + \sum_{k=25}^{26} I_k$

UQFF Page Time

$$t_P^{\text{UQFF}} = t_P^{\text{GR}} \times e^{+\kappa \cdot t_{\text{evap}}} \approx t_P^{\text{GR}} \times (1 + \kappa \cdot t_{\text{evap}})$$

For stellar BH ($t_{\text{evap}} \sim 1074$ s): correction factor is astronomically large physically this means that within any finite observation time, the UQFF Page curve looks **thermal** to a 4D observer, with the information recovery pushed beyond any accessible time. This is consistent with the absence of observational evidence for information recovery in Hawking radiation.

4. Firewall Prevention

The AMPS firewall paradox requires choosing between a smooth horizon (infalling observer) or pure Hawking radiation. The UQFF resolves this through **channels 1924 non-local entanglement**: the infalling observer encounters a smoothly modified vacuum (Superconductive mode, $E_{\text{react}} 10^?$) rather than a firewall, while the external observer's Hawking radiation becomes approximately (not exactly) thermal.

Summary

Aspect	Standard QM	UQFF 26D	Resolution
Information storage	In BH?	Channels 2526	Non-local, always preserved
Page curve	QES prediction	UQFF ?-modified	Extended beyond t_{evap}
Firewall	AMPS paradox	SCm smooth horizon	Channels 1924 entanglement
4D observer	Thermal Hawking	Approximately thermal	Consistent

Source: `MAIN_1_CoAnQi.cpp` Batch 21 (`InformationParadoxModule`) | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
$[SSq]$	0.57	Universal Quantized Factor
β_i	0.60-0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma\lambda_i\cdot U_i\cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{\text{SCm}} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434\cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$U_m \times (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, Condensed-Physics.py, and CondensedPhysics2.py.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **BH-gravity** sector of the 9-sector UQFF Lagrangian (see uqff_lagrangian_derivation.py).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_\mu \phi_{\text{BH}})(\partial^\mu \phi_{\text{BH}}) - V(\phi_{\text{BH}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{BH}}) = \frac{1}{2}m^2\phi_{\text{BH}}^2 + \frac{\lambda}{4!}\phi_{\text{BH}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{BH}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{BH}}} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \rho_{\text{vac},[\text{SCm}]}g_{\mu\nu} + F_{U_Bi_i}/r^2 = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{BH}} =$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.083 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 97, \quad n_{\text{channel}} = 7/26$$

Since $p_{\text{DVP}} = 97$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii

where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is $10^6 \mathbf{M_BH/M_\odot \text{ yr}}$ (quasi-normal mode ringdown):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_\odot}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.083	✓ Threshold-consistent
DVP prime BSH layers	$p_k \in \{2,3,\dots,113\}$ 26 harmonic terms	$p_{\text{DVP}} = 97$ $j = 1\dots 26,$ $\cos(2\pi j/26)$	✓ Resonant ✓ Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	✓ Canonical
[SSq]	0.57	Applied in BSH saturation	✓ Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}] * n / 26)$
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}} / F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2 * \Gamma_2)}] * (1 + [\text{SSq}] * n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z) / (1 - 2^{\{1-26\}}) + 2^{\{1-26\} / (1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: - $f_{26}(q) = \sum_{n=0}^{25} q^{\binom{n}{2}} / (-q;q)_{26}$ - $\phi_{26}(q) = \sum_{n=0}^{25} q^{\binom{n}{2}} / (-q^2;q^2)_{26}$ - $\psi_{26}(q) = \sum_{n=1}^{26} q^{\binom{n}{2}} / (q;q^2)_{26}$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	Symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$
where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq] \exp(-\kappa_{ppai} * n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_SCm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).