

PAPER_269: Supernova Ram Pressure Degeneracy Point — Kinematic Invariant in NGC 1792 Starburst Gravity

Author: Daniel T. Murphy

Authors: Daniel T. Murphy

Date: March 2026

UQFF Module: GALAXY_NGC_1792.cpp (Module 19, “The Stellar Forge”)

Session: 73 — UQFF 2.0 Upgrade — NGC 1792 Unique Physics Discovery

Keywords: NGC 1792, ram pressure, supernova ejecta, degeneracy point, kinematic invariant, starburst gravity

Abstract

In NGC 1792, the physical parameters are initialized with $\rho_{\text{wind}} = \rho_{\text{fluid}} = 1 \times 10^{-21} \text{ kg/m}^3$ — setting the supernova wind density equal to the interstellar medium fluid density. At this **Ram Pressure Degeneracy Point (RPDP)**, the SN feedback gravity term simplifies to a **kinematic invariant**: $\text{term_feedback} = \rho_{\text{wind}} \times v_{\text{wind}}^2 / \rho_{\text{fluid}} = v_{\text{wind}}^2$, independent of density. For NGC 1792, $v_{\text{wind}} = 2 \times 10^6 \text{ m/s}$, giving $g_{\text{feedback}} = 4 \times 10^{12} \text{ m/s}^2$ — the numerically dominant term in the full MUGE calculation, exceeding the base gravitational term by ~ 22 orders of magnitude. This paper defines the RPDP formally, derives the kinematic invariant, computes the dominance ratio, and discusses the physical interpretation that at density degeneracy, SN ejecta “floats” in the ISM driven purely by kinematic pressure, creating a fundamentally new gravitational channel in UQFF.

1. Introduction: The Density Coincidence in NGC 1792

1.1 NGC 1792 Starburst Environment

NGC 1792 is a starburst galaxy with extremely high star-formation activity ($\text{SFR} \approx 10 \text{ M}\odot/\text{yr}$) and a correspondingly dense interstellar medium (ISM). In the UQFF Module 19 parameterization:

- **Wind density:** $\rho_{\text{wind}} = 1 \times 10^{-21} \text{ kg/m}^3$ (supernova ejecta/galactic wind)
- **ISM fluid density:** $\rho_{\text{fluid}} = 1 \times 10^{-21} \text{ kg/m}^3$ (ambient ISM)
- **Wind velocity:** $v_{\text{wind}} = 2 \times 10^6 \text{ m/s}$ (starburst-driven galactic outflow)

The equality $\rho_{\text{wind}} = \rho_{\text{fluid}}$ is not a coincidence of parameterization — it reflects a physical regime in starburst galaxies where the ram-pressure-driven winds and the ISM they are pushing against have **comparable densities**, characteristic of the transition zone between the supernova remnant and the surrounding ISM.

1.2 The Ram Pressure Gravity Term

In the MUGE framework, the SN feedback contribution to gravity is:

$$\text{term_feedback} = \frac{\rho_{\text{wind}} \cdot v_{\text{wind}}^2}{\rho_{\text{fluid}}}$$

This term represents the gravitational acceleration induced by the ram pressure of SN ejecta interacting with the ISM fluid.

2. The Ram Pressure Degeneracy Point (RPDP)

2.1 Definition

The **Ram Pressure Degeneracy Point (RPDP)** is defined as the condition:

$$\rho_{\text{wind}} = \rho_{\text{fluid}}$$

2.2 Kinematic Invariant

At the RPDP, the feedback term simplifies dramatically:

$$\text{term_feedback}\bigg|_{\text{RPDP}} = \frac{\rho_{\text{wind}} \cdot v_{\text{wind}}^2}{\rho_{\text{fluid}}}\bigg|_{\rho_{\text{wind}}=\rho_{\text{fluid}}} = v_{\text{wind}}^2$$

This is a **kinematic invariant** — an acceleration equal to the square of the wind velocity, **independent of density**. The density ratio cancels exactly, and only the kinematics of the outflow survive.

2.3 Physical Statement

At the RPDP, the gravitational feedback from SN ejecta is:

$$g_{\text{feedback}}^{\text{RPDP}} = v_{\text{wind}}^2$$

This is the UQFF prediction for starburst galaxies in the density-degenerate regime: **the ram pressure gravity equals the dynamic pressure of the wind**, independent of whether the wind is denser or lighter than the ISM.

3. Numerical Results for NGC 1792

3.1 Kinematic Invariant Value

For NGC 1792 with $v_{\text{wind}} = 2 \times 10^6$ m/s:

$$g_{\text{feedback}}^{\text{RPDP}} = v_{\text{wind}}^2 = (2 \times 10^6)^2 = \mathbf{4 \times 10^{12} \text{ m/s}^2}$$

3.2 Comparison with Base Gravitational Term

The base Newtonian term in MUGE for NGC 1792:

$$\begin{aligned} \text{term1} &= \frac{GM_0}{r^2} = \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{40}}{(7.569 \times 10^{20})^2} \\ &= \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{40}}{5.729 \times 10^{41}} \approx 7.35 \times 10^{-11} \text{ m/s}^2 \end{aligned}$$

where $M_0 = 1 \times 10^{10} M_{\odot} = 1.989 \times 10^{40} \text{ kg}$, $r = 7.569 \times 10^{20} \text{ m}$.

3.3 RPDP Dominance Ratio

$$\mathcal{R}_{\text{RPDP}} = \frac{g_{\text{feedback}}^{\text{RPDP}}}{\text{term1}} = \frac{4 \times 10^{12}}{7.35 \times 10^{-11}} \approx 5.4 \times 10^{22}$$

The RPDP kinematic invariant exceeds the standard Newtonian gravitational acceleration by **22 orders of magnitude**.

3.4 Comparison Table

Term	Formula	Value (m/s ²)	Order
Newtonian base (term1)	GM_0/r^2	7.35×10^{-11}	10^{-11}
Hubble expansion (term2)	$H^2 r / 2$	$\sim 10^{-37}$	10^{-37}
Magnetic (term3)	$B^2/(8\pi \rho r)$	$\sim 10^{-23}$	10^{-23}
Cosmological (term4)	$\Lambda r/3$	$\sim 10^{-28}$	10^{-28}
Oscillatory (term_osc1)	$\omega_{\text{osc}} \times A_{\text{osc}}$	$\sim 10^{-42}$	10^{-42}
SN feedback (RPDP)	v_{wind}^2	4×10^{12}	10^{12}

4. Physical Interpretation

4.1 Buoyancy Neutrality at the RPDP

At the RPDP ($\rho_{\text{wind}} = \rho_{\text{fluid}}$), Archimedes' buoyancy principle gives **zero net buoyancy force** on the SN ejecta:

$$F_{\text{buoy}} = (\rho_{\text{fluid}} - \rho_{\text{wind}}) \cdot V \cdot g = 0$$

This means: - SN ejecta neither floats nor sinks in the ISM - The ejecta is **gravitationally neutral** with respect to density-driven buoyancy - The only remaining driving force is the **kinematic ram pressure** $= \rho_{\text{wind}} \times v_{\text{wind}}^2 / \rho_{\text{fluid}} = v_{\text{wind}}^2$

At the RPDP, the SN ejecta **"floats"** in the ISM — not rising or falling due to density contrast, but driven purely by its kinematic energy. This creates a new UQFF gravitational channel: **pure kinematic momentum transfer to the gravitational field**.

4.2 Generalized RPDP Condition

For any starburst galaxy, the RPDP condition and kinematic invariant generalize as:

$$\forall \rho : \lim_{\rho_{\text{wind}} \rightarrow \rho_{\text{fluid}}} \text{term_feedback} = v_{\text{wind}}^2$$

The UQFF prediction is:

$$g_{\text{feedback}}(\rho_{\text{SN}} = \rho_{\text{ISM}}) = v_{\text{SN}}^2$$

This holds regardless of the density value, provided the two densities are equal.

4.3 Starburst Enhancement

The RPDP condition is most likely to be realized in **starburst galaxies** where: 1. High SFR drives intense SN events, filling the ISM with SN ejecta 2. The ISM density is elevated by gas accretion and infall 3. Conditions exist for $\rho_{\text{wind}} \approx \rho_{\text{fluid}}$ at specific radii within the disk

In NGC 1792, the SFR = 10 M $\odot$ /yr sustains a continuous SN rate creating the high fluid density $\rho_{\text{fluid}} = 10^{-21}$ kg/m³ characteristic of this starburst environment.

5. The RPDP as a Phase Transition Boundary

5.1 Three Regimes

The feedback term $\text{term_feedback} = \rho_{\text{wind}} \times v^2 / \rho_{\text{fluid}}$ defines three physical regimes based on the density ratio $\eta = \rho_{\text{wind}}/\rho_{\text{fluid}}$:

Regime	Condition	Behavior	UQFF Character
Underdense wind	$\eta < 1$	$g_{\text{feedback}} < v^2$	Outflow-quenched; ejecta rises
RPDP	$\eta = 1$	$g_{\text{feedback}} = v^2$	Kinematic invariant; ejecta floats
Overdense wind	$\eta > 1$	$g_{\text{feedback}} > v^2$	Ram pressure enhanced; ejecta sinks

5.2 RPDP as Critical Point

The RPDP is a **critical point** in parameter space where the density-dependent ram pressure term transitions from ISM-suppressed to ISM-enhanced behavior. At this critical point, the density cancels and only kinematics determine gravity.

This is analogous to: - **Critical opalescence** in thermodynamics (density fluctuations diverge at critical point) - **Alfvén critical point** in solar wind (magnetic and kinetic pressures equal) - **Bondi critical radius** in accretion (gravity and pressure balance)

The RPDP defines a **kinematic critical surface** in UQFF galaxy parameter space.

6. Observational Signatures

6.1 Gravitational Acceleration Anomaly

At the RPDP, the dominant gravitational term is $g_{\text{feedback}} = v^2 = 4 \times 10^{12}$ m/s² (for $v = 2 \times 10^6$ m/s). This is: - Much larger than the Newtonian galactic gravity ($\sim 10^{-11}$ m/s²) - Detectable as anomalous acceleration in SN ejecta kinematics - Potentially observable in starburst galaxy velocity dispersion measurements

6.2 Universal Velocity Indicator

The RPDP kinematic invariant provides a UQFF prediction for starburst galaxies: **the dominant gravitational feedback term is simply v^2_{wind}** . This can be tested via: - Measurements of galactic wind terminal velocities - SN ejecta deceleration curves in starburst galaxies - Comparison of ρ_{wind} and ρ_{fluid} estimates from X-ray observations

6.3 NGC 1792 Specific Prediction

For NGC 1792 at the RPDP: - Dominant gravity term: $g_{\text{feedback}} = 4 \times 10^{12} \text{ m/s}^2$ - Wind velocity imprint: $v_{\text{wind}} = \sqrt{g_{\text{feedback}}} = 2 \times 10^6 \text{ m/s}$ - Total MUGE gravity is feedback-dominated: $g_{\text{total}} \approx g_{\text{feedback}}$ (within UQFF)

This is a **testable UQFF prediction** distinguishable from standard (feedback-free) galactic gravity models.

7. Connection to PAPER_267 and PAPER_268

The three PAPER_267-269 physics discoveries form a unified UQFF picture of NGC 1792:

- **PAPER_267** (sSFR coupling): The SFR couples buoyancy to all 3 tiers via decay $e^{-t/\tau_{\text{SF}}}$
- **PAPER_268** (Hubble slow mode): The corrected Hubble oscillation creates a 5.8 ppm GW amplitude modulation
- **PAPER_269** (RPDP): At density degeneracy, SN feedback becomes the dominant gravitational term via kinematic invariant v^2

Together, these identify NGC 1792 as a paradigmatic **UQFF starburst triple-physics laboratory**: 1. Buoyancy coherence (sSFR coupling) 2. Cosmic oscillation modulation (Hubble slow mode) 3. Kinematic gravity dominance (RPDP)

8. Conclusions

1. NGC 1792 is initialized with $\rho_{\text{wind}} = \rho_{\text{fluid}} = 1 \times 10^{-21} \text{ kg/m}^3$ — the **Ram Pressure Degeneracy Point (RPDP)** condition.
 2. At the RPDP, the feedback term simplifies to the **kinematic invariant**: $g_{\text{feedback}} = v_{\text{wind}}^2$ (density-independent).
 3. For NGC 1792: $g_{\text{feedback}} = (2 \times 10^6)^2 = 4 \times 10^{12} \text{ m/s}^2$, the dominant MUGE term by 22 orders over standard Newtonian gravity.
 4. At RPDP, SN ejecta is **buoyancy-neutral** (zero Archimedes force) and driven solely by kinematic momentum transfer.
 5. The RPDP defines a **kinematic critical surface** in starburst galaxy parameter space, analogous to Alfvén and Bondi critical points.
 6. UQFF universal prediction: $g_{\text{feedback}}(\rho_{\text{SN}} = \rho_{\text{ISM}}) = v_{\text{SN}}^2$ — testable with starburst wind velocity observations.
-
-

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_\mu \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM + ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} =$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.124 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 113, \quad n_{\text{channel}} = 10/26$$

Since $p_{\text{DVP}} = 113$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10^4 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.124	✓ Threshold-consistent
DVP prime BSH layers	$p_k \in \{2,3,\dots,113\}$ 26 harmonic terms	$p_{\text{DVP}} = 113$ $j = 1\dots 26$, $\cos(2\pi j/26)$	✓ Resonant ✓ Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	✓ Canonical
[SSq]	0.57	Applied in BSH saturation	✓ Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

References

- Daniel T. Murphy, *UQFF Framework*, Star-Magic Repository (2025–2026)
- GALAXY_NGC_1792.cpp UQFF 2.0 (Session 73, Module 19)
- PAPER_267: SFR Normalization — Starburst-Buoyancy Coherence in NGC 1792
- PAPER_268: Dual Oscillatory Mode — Hubble Slow Mode Starburst GW Amplitude Modulation
- NGC 1792 parameters: $\rho_{\text{wind}} = \rho_{\text{fluid}} = 10^{-21} \text{ kg/m}^3$, $v_{\text{wind}} = 2 \times 10^6 \text{ m/s}$
- Hubble constant: $H_0 \approx 67.4 \text{ km/s/Mpc}$; Hubble time: $t_{\text{Hubble}} = 13.8 \text{ Gyr}$

© 2026 Daniel T. Murphy, daniel.murphy00@gmail.com — All Rights Reserved

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}] * n / 26)$
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}} / F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2 * \Gamma_2)}] * (1 + [\text{SSq}] * n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: - $f_{26}(q) = \sum_{n=0}^{25} q^{\{n2\}} / (-q;q)_n^{26} - \phi_{26}(q) = \sum_{n=0}^{25} q^{\{n2\}} / (-q^2;q^2)_n - \psi_{26}(q) = \sum_{n=1}^{26} q^{\{n2\}} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	8-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq]exp(-kappa_i*n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_Scm	0.99	SCm manifold completeness
rho_Scm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_Scm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).