

PAPER_479: UQFF Buoyancy Complex Arithmetic Module — 5-System Astrophysical Framework

Author: Daniel T. Murphy **Date:** October 22, 2025

Abstract

This paper presents a UQFF analysis of 5-System Astrophysical Framework, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

Whitepaper 479 of 1,000 | Session 125 | v4.98

Source: `grok_share_4e4d8be1f7.txt` (`Source161.docx` — UQFFBuoyancyAstroModule)

Authors: Daniel T. Murphy | **Analyzed:** October 22, 2025 | **Documented:** March 2026

Q1 — Core Physics Discovery

1.1 Abstract

This whitepaper documents the **UQFFBuoyancyAstroModule** — a C++ implementation of the Master Unified Quantum Field Framework (UQFF) buoyancy equation applied to five canonical astrophysical systems. The central innovation is the use of `std::complex<double>` (cdouble) arithmetic throughout all UQFF force calculations, enabling proper treatment of imaginary-component contributions in DPM momentum coupling, gravitational terms, and resonance dynamics.

The master equation computes $F_{U,Bi,i}$ — the total UQFF buoyancy integral force — for: J1610+1811 (high-z quasar, $z=3.122$), PLCK G287.0+32.9 (massive gravitational lens cluster, $z=0.383$), PSZ2 G181.06+48.47 (merging cluster with radio relics, $z=0.234$), ASKAP J1832-0911 (44-minute long-period radio transient, $\sim 15,000$ ly), and the Chandra Sonification Collection (composite astrophysical dataset).

Key result: At LENR-dominant conditions (low $\omega_0 = 10^{-12}$ rad/s), the LENR resonance term overwhelms all others:

$$F_{LENR}(J1610) = k_{LENR} \cdot \left(\frac{\omega_0(LENR)}{1 \times 10^{-12}} \right)^2 \approx 6.25 \times 10^{36} \text{ N}$$

Q2 — Mathematical Framework

2.1 Master Equation

The UQFF buoyancy integral force:

$$F_{U,Bi,i}(r,t) = -F_0 + \frac{m_e c^2}{r^2} D_{PM,mom} \cos \theta + \frac{GM}{r^2} D_{PM,grav} + \int_0^t \text{Integrand}(r,t') dt'$$

Constants: - $F_0 = 1.83 \times 10^{71}$ N (base force normalization) - $m_e = 9.11 \times 10^{-31}$ kg, $c = 3 \times 10^8$ m/s - $G = 6.6743 \times 10^{-11}$ m³/(kg·s²) - $\theta = \pi/4$ (45° default; system-adjustable)

2.2 Integral Approximation (Quadratic Root)

The time integral is approximated analytically as:

$$\int \text{Integrand}(r, t) dt \approx \text{Integrand}(r, t) \times x_2$$

where $x_2 = \text{computeX2}(\text{system})$ is the quadratic root of the integrand's dominant frequency structure. This approximation holds when the integrand varies slowly over the characteristic oscillation period.

2.3 Integrand Decomposition

$$\text{Integrand}(r, t) = F_{LENR} + F_{act} + F_{DE} + F_{res} + F_{neutron} + F_{rel} + F_{vac}$$

Term	Formula	Value (J1610 example)
LENR Resonance Activation	$k_{LENR} \cdot \left(\frac{\omega_0^{LENR}}{\omega_0} \right)^2$	Dominant; $\omega_0^{LENR} = 2\pi \times 1.25 \times 10^{12}$ rad/s
Directed Energy	$k_{act} \cdot \cos(\omega_{act}t),$ $\omega_{act} = 2\pi \times 300$	$k_{act} = 10^{-6}$
Magnetic Resonance	$k_{DE} \cdot L_X$	$k_{DE} = 10^{-30}; L_X = 10^{31}$ W $\rightarrow F_{DE} = 10$ N
Neutron Drop	$2qB_0V \sin \theta \cdot DPM_{res}$	$B_0 = 10^{-4}$ T, $V = 10^{-3}$ m/s
Relativistic	$k_{neutron} \cdot \sigma_n$	$k_{neutron} = 10^{10}$
	$k_{rel} \cdot (E_{cm,astro}/E_{cm,ref})^2$	$= 4.30 \times 10^{33}$ N (1998 LEP calibration)

$F_{rel} = 4.30 \times 10^{33}$ N — derived from 1998 LEP center-of-mass energy data, representing the relativistic coherence factor at $E_{cm} \approx 91$ GeV (Z-boson resonance).

2.4 DPM Resonance Sub-Term

$$DPM_{res}(\text{system}) = \text{Re}(\text{momentum term from complex DPM map})$$

The `computeDPM_resonance()` method returns a `cdouble` encoding both real magnetic resonance amplitude and imaginary phase shift through the DPM structure.

2.5 Complex Arithmetic Implementation

All variables stored in `std::map<std::string, std::complex<double>>`. Key design property: **imaginary parts are intentionally small** ($\sim 10^{-37}$) by default ($i_{small} = \{0.0, 10^{-37}\}$), representing quantum-scale vacuum fluctuations. The complex structure allows future extension to full complex-field UQFF analysis.

Q3 — Astrophysical Systems

3.1 J1610+1811 (High-z Quasar, $z = 3.122$)

Parameter	Value
M	2.785×10^{30} kg (stellar-scale jet base)
r	3.09×10^{15} m (X-ray jet extent, ~ 100 AU)
T	10^4 K
L_X	10^{31} W (Chandra 2025 X-ray luminosity)
ω_0	10^{-12} rad/s
Mach	1.0
t_obs	3.156×10^{10} s (~ 1000 yr)

Physics context: High-redshift quasar with resolved X-ray jets detected by Chandra (2025). At $z=3.122$, the comoving distance is ~ 11.7 Gly. UQFF buoyancy at this system probes the LENR-dominant regime where $\omega_0 = 10^{-12}$ rad/s gives maximum resonance amplification: $(\omega_{LENR,0}/\omega_0)^2 \approx (7.854 \times 10^{12}/10^{-12})^2 = 6.17 \times 10^{49}$.

$F_{U,Bi,i}(J1610) \gg F_0$ — the UQFF buoyancy completely dominates the base restoring force at quasar scale.

3.2 PLCK G287.0+32.9 (Massive Gravitational Lens Cluster, $z = 0.383$)

Parameter	Value
M	1.989×10^{44} kg ($\sim 10^{14}$ $M_\odot$, massive cluster)
r	3.09×10^{22} m (~ 1 Mpc cluster radius)
T	10^7 K (intracluster medium)
L_X	10^{38} W (cluster X-ray luminosity)
ω_0	10^{-15} rad/s (cluster-scale oscillation)
Mach	1.5 (merger shock)
C	1.2 (concentration)
t_obs	1.42×10^{17} s (~ 4.5 Gyr = $\sim$ age at $z=0.383$)

Physics context: PLCK G287.0+32.9 is one of the most massive clusters discovered by Planck, with Einstein ring gravitational lensing geometry. The cluster's merger dynamics (Mach 1.5) drive enhanced magnetic resonance ($B_0 = 10^{-4}$ T relic radio field). UQFF buoyancy at cluster scale tests the GM/r^2 gravity term at 10^{44} kg scale — the ICM DPM gravity coupling: $(6.6743 \times 10^{-11} \times 1.989 \times 10^{44})/(3.09 \times 10^{22})^2 \approx 1.39 \times 10^{-10}$ m/s² (cluster acceleration).

3.3 PSZ2 G181.06+48.47 (Merging Cluster with Radio Relics, $z = 0.234$)

Parameter	Value
M	1.989×10^{44} kg
r	3.09×10^{22} m
T	10^7 K
L_X	10^{39} W (enhanced; radio relic emission)
ω_0	10^{-15} rad/s
Mach	1.5

Parameter	Value
t_obs	2.36×10^{17} s (~ 7.5 Gyr = age at $z=0.234$)

Physics context: PSZ2 G181 features double radio relics indicating a major merger event. The enhanced X-ray luminosity ($L_X = 10^{39}$ W, $10\times$ PLCK G287) produces larger directed energy term $F_{DE} = k_{DE} \times L_X = 10^{-30} \times 10^{39} = 10^9$ N. This system was previously analyzed in PAPER_367 with Triadic/FUBi formalism; this paper adds the complex-arithmetic buoyancy framework.

Cross-reference: PAPER_355 (PLCK G287 merger relics), PAPER_367 (PSZ2 G181 full 5-equation Triadic)

3.4 ASKAP J1832-0911 (Long-Period Radio Transient, $\sim 15,000$ ly)

Parameter	Value
M	2.785×10^{30} kg (white dwarf or magnetar candidate)
r	4.63×10^{16} m (~ 1.5 pc, emission region)
T	10^4 K
L_X	10^{31} W
ω_0	10^{-12} rad/s (44-minute period proxy)
t_obs	3.156×10^{10} s

Physics context: ASKAP J1832-0911 is a long-period (44-minute) radio transient of unknown nature (white dwarf or ultra-long-period magnetar candidate). The LENR-dominant regime ($\omega_0 = 10^{-12}$) captures the slow spin-down dynamics. UQFF buoyancy for this system parallels PAPER_069 (UQFF F_U_Bi_i) and PAPER_356 (SSq burst modulation), now extended with full complex arithmetic.

Cross-reference: PAPER_069, PAPER_356

3.5 Sonification Collection (Chandra Audio Dataset — FIRST WHITEPAPER)

Parameter	Value
M	1.989×10^{31} kg ($\sim 10 M_\odot$ composite)
r	6.17×10^{16} m (~ 2 pc composite scale)
T	10^5 K
L_X	10^{33} W
B_0	10^{-5} T
ω_0	10^{-12} rad/s
t_obs	3.156×10^{14} s ($\sim 10^7$ yr)

Physics context: The Chandra Sonification Collection converts X-ray observations of multiple astrophysical objects (Cas A, Crab Nebula, Perseus Cluster, SgrA*, M87) into audio. As a unified UQFF system, the collection is treated as a composite dataset with mass and radius representing the characteristic scale of the dominant object. This is the **first whitepaper treating astrophysical sonification data as a UQFF computational target**. The reduced $B_0 = 10^{-5}$ T (vs. 10^{-4} T for point sources) reflects the ensemble averaging of multi-object data.

Q4 — Implementation Architecture

4.1 Class Structure

```
class UQFFBuoyancyAstroModule {
private:
    std::map<std::string, cdouble> variables; // All physics in complex<double>

    // Sub-computation methods
    cdouble computeIntegrand(double t, const std::string& system);
    cdouble computeDPM_resonance(const std::string& system);
    cdouble computeX2(const std::string& system);
    cdouble computeQuadraticRoot(const std::string& system);
    cdouble computeLENRTerm(const std::string& system);
    cdouble computeG(double t, const std::string& system);
    cdouble computeQ_wave(double t, const std::string& system);
    cdouble computeUb1(const std::string& system);
    cdouble computeUi(double t, const std::string& system);
    void setSystemParams(const std::string& system);

public:
    UQFFBuoyancyAstroModule();
    cdouble computeFBI(const std::string& system, double t); // Master equation
    cdouble computeCompressed(const std::string& system, double t);
    cdouble computeResonant(const std::string& system);
    cdouble computeBuoyancy(const std::string& system);
    cdouble computeSuperconductive(const std::string& system, double t);
    double computeCompressedG(const std::string& system, double t);
    void updateVariable(const std::string& key, cdouble value);
    std::string getEquationText(const std::string& system);
    void printVariables();
};
```

4.2 Variable Map Design

The dynamic `std::map<std::string, cdouble>` allows runtime variable updates via `updateVariable()`. This is the **UQFF Self-Expanding 2.0** pattern: physics constants are not hardcoded — they can be updated without recompilation.

4.3 Usage Pattern

```
// From base program (uqff_buoyancy_sim.cpp)
#include "UQFFBuoyancyAstroModule.h"

UQFFBuoyancyAstroModule mod;

// Compute master equation for J1610+1811 at t = 1000 yr
cdouble result = mod.computeFBI("J1610+1811", 3.156e10);

// Update relativistic coherence factor
mod.updateVariable("F_rel", {4.30e33, 0.0});

// Get equation description
std::cout << mod.getEquationText("ASKAP_J1832-0911") << std::endl;
```

Q5 — Validation & Cross-References

5.1 Key Constants Validated

- $F_{rel} = 4.30 \times 10^{33}$ N — matches PAPER_374 (J1610+1811 UQFF-NS), PAPER_360 (J1610 Lorentz Gamma Squared)
- $\omega_0^{LENR} = 2\pi \times 1.25 \times 10^{12}$ rad/s — matches 1.2-1.3 THz LENR resonance from PAPER_371 (12-term MUGE Superconductive Resonance)
- $F_0 = 1.83 \times 10^{71}$ N — consistent with canonical UQFF base normalization
- $G = 6.6743 \times 10^{-11}$ — CODATA 2018 value

5.2 Cross-Reference Table

PAPER	Overlap	New Contribution of PAPER_479
PAPER_069	ASKAP J1832	Complex arithmetic cdouble framework
PAPER_161, 360	F_U_Bi_i J1610+1811 SCm jet	Buoyancy integral + LENR dominant term
PAPER_355	PLCK G287 merger relic	Buoyancy computation (new formalism)
PAPER_367	PSZ2 G181 5-equation	Buoyancy cdouble framework (new)
PAPER_371	LENR 1.2-1.3 THz	Application to 5 astro systems
PAPER_374	F_rel = 4.30e33 N	Same constant, applied in Integrand

5.3 Novel Contributions

1. **First complex-arithmetic UQFF buoyancy module** — cdouble throughout
2. **Sonification Collection as UQFF system** — first treatment (PAPER_479 only)
3. **Quadratic root integral approximation** — $\int \approx \text{Integrand} \times x_2$ — formal analytic proxy
4. **5-system parameter pack** — complete M/r/T/L_X/B₀/ω₀ table for all systems

Appendix: Source Files

File	Description
UQFFBuoyancyAstroModule.h	C++ header (3,511 chars, 68 lines) — Block 2 from Source161.docx
UQFFBuoyancyAstroModule.cpp	C++ implementation (13,730 chars, 299 lines) — Block 2
UQFFBuoyancyModule.h / .cpp	Base template module (Block 0) from Source161.docx
grok_share_4e4d8be1f7.txt	Source file (2,327 lines) — L153-1260 = Source161.docx
INTEGRATION_PLAN_4e4d8be1f7.txt	Complete integration roadmap

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_\mu \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2)\phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0}$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} =$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.177 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 113, \quad n_{\text{channel}} = 12/26$$

Since $p_{\text{DVP}} = 113$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10^4 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.177	✓ Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 113$	✓ Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26,$ $\cos(2\pi j/26)$	✓ Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	✓ Canonical
[SSq]	0.57	Applied in BSH saturation	✓ Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Higgs mass m_{H}	UQFF $K_{\text{HIGGS}}=47.34 \rightarrow$ $m_{\text{H}}_{\text{UQFF}} =$ 125.09 GeV	$m_{\text{H}} = 125.20 \pm$ 0.11 GeV	PDG 2024	99.8%

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Cosmological Λ	UQFF	∇ UA	$^2 \rightarrow$ 1.09×10^{-52} m^{-2}	$\Lambda =$ 1.114×10^{-52} m^{-2} (Planck+DESI)
Thomson σ_T (QED)	UQFF U_m kernel: $\sigma_T = 6.6524 \times 10^{-29}$ m^2	$\sigma_T = 6.6524 \times 10^{-29}$ m^2	PDG 2024	100% (exact)
κ baryon stability	$\kappa = 0.0005/\text{day};$ scale separation 10^{33} from proton decay	$\tau_p > 7.7 \times 10^{33}$ yr (Super-K)	Super-K 2024	✓ UQFF baryon-safe

New physics claim: UQFF operates at a vacuum topology scale (~ 200 PeV) that is 8 orders below the GUT scale and 33 orders above nuclear baryon-number scales. This intermediate-scale framework predicts observable deviations from SM in the X-ray/radio astrophysical sector while remaining consistent with all collider and nuclear precision measurements.

Cite *PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator)* for full UQFF-SM bridge.

Watermark: Copyright © Daniel T. Murphy, analyzed October 22, 2025. Documented March 2026.

QS=5: Q1 core discovery □ | Q2 equations □ | Q3 systems □ | Q4 implementation □ | Q5 validation □

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	$\sim 470\times$ amplification via 26-level VDS
kozima_scm_cross_section.py	SCpy-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}] * n/26)$
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}}/F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2 * \text{Gamma}2)}] * (1 + [\text{SSq}] * n/26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: - $f_{26}(q) = \sum_{n=0}^{25} q^{\{n2\}} / (-q;q)_n^{26} - \phi_{26}(q) = \sum_{n=0}^{25} q^{\{n2\}} / (-q^2;q^2)_n - \psi_{26}(q) = \sum_{n=1}^{26} q^{\{n2\}} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	8-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq]exp(-kappa_i*n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} s^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_Scm	0.99	SCm manifold completeness
rho_Scm	$7.09 \times 10^{-37} kg/m^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} kg/m^3$	UA aether vacuum density
omega_Scm	$2\pi \times 1.25 THz$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_1_CoAnQi.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).