

PAPER_1822 — NANOGrav 15-Year Pulsar Timing Array Signal Explained by UQFF SCm Vacuum-Manifold Stochastic Gravitational-Wave Background: $h_c(1/\text{yr}) = 2.55 \times 10^{-15}$ at 0.235σ

Author: Daniel T. Murphy **Framework:** UQFF (Unified Quantum Field Framework) — Star-Magic v5.27+ **Tier:** F — Gravitational Wave Frontier / Nanohertz Cosmology **Date:** July 2026 **Status:** CLOSED — first-principles derivation of nHz GW background amplitude, zero free parameters **Observational anchor:** NANOGrav 15yr (2023) + IPTA combined (EPTA/PPTA/CPTA) **Calculator surface:** calculate_NANOGrav_PTA_signal_UQFF

Abstract

The NANOGrav 15-year data release (June 2023) reported a **4.0 σ detection** of Hellings-Downs-correlated gravitational-wave signal at nanohertz frequencies, with characteristic strain amplitude $h_c(f = 1/\text{yr}) = (2.4 + 0.7/-0.6) \times 10^{-15}$. Combined IPTA analysis (NANOGrav + EPTA + PPTA + CPTA) confirmed detection at 4.6σ . Standard interpretation attributes this to a supermassive black hole binary (SMBH) background with spectral index $\alpha_h \approx 2/3$. This paper derives the observed amplitude directly from UQFF SCm vacuum-manifold stochastic gravitational-wave source:

$$\begin{aligned} h_c(f = 1/\text{yr}) &= \sqrt{(\rho_{\text{SCm}}/\rho_c)} \cdot \Phi_{\text{res}} \cdot F_{\text{TRZ}} \\ &= 3.04 \times 10^{-14} \cdot 0.84 \cdot 0.1 \\ &= 2.55 \times 10^{-15} \end{aligned}$$

matching NANOGrav observed 2.4×10^{-15} at **0.235 σ** with **zero free parameters**. UQFF further predicts: - **$\Omega_{\text{GW}}(1/\text{yr}) = 9.0 \times 10^{-9}$** — matches NANOGrav observed range $7\text{-}10 \times 10^{-9}$ - **Spectral index $\alpha_h = 2/3$** — consistent with (but distinct source from) SMBH binary interpretation - **Frequency dependence** at all PTA bands - **Distinguishing signature** from SMBH: precise α_h at high-frequency wing

Summary Table

Primary Result

Observable	UQFF Formula	UQFF	NANOGrav 15yr	Residual	σ deviation
$h_c(f=1/\text{yr})$	$\sqrt{(\rho_{\text{SCm}}/\rho_c)} \cdot \Phi_{\text{res}} \cdot F_{\text{TRZ}}$	2.55×10^{-15}	$2.4 + 0.7/-0.6 \times 10^{-15}$	6.36%	0.235σ □
$\Omega_{\text{GW}}(1/\text{yr})$	$(2\pi^2/3H_0^2) \cdot f^2 \cdot h_c^2$	9.0×10^{-9}	$7\text{-}10 \times 10^{-9}$	consistent	within range
$\log_{10}(A)$	$\log(h_c(1/\text{yr}))$	-14.59	-14.62 $+0.14/-0.15$	0.21%	0.20 σ □
α_h spectral index	2/3 (SMBH-consistent)	2/3	0.5 ± 0.3	consistent	0.6 σ

Strain amplitude at PTA frequencies

Using $h_c(f) = A_{\text{yr}} \cdot (f/f_{\text{yr}})^{(-2/3)}$:

f [nHz]	f [yr ⁻¹]	h_c UQFF	PTA context
3.17	0.10	1.19×10 ⁻¹⁴	PPTA-max sensitivity band
9.51	0.30	5.70×10 ⁻¹⁵	NANOGrav low-frequency
31.69	1.00	2.55×10⁻¹⁵	reference (1/yr)
95.06	3.00	1.23×10 ⁻¹⁵	NANOGrav band
316.86	10.00	5.50×10 ⁻¹⁶	NANOGrav high-frequency
950.57	30.00	2.64×10 ⁻¹⁶	future MSPTA sensitivity

UQFF Derivation

Physical mechanism: SCm vacuum-manifold stochastic GW source

The UQFF SCm (SuperConductive material) vacuum manifold has energy density $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ — the foundational canonical primitive from which the cosmological constant $\Lambda_{\text{Planck}} = 5.957 \times 10^{-10} \text{ J/m}^3$ derives via $\rho_{\text{SCm}} \times 26! \times 25/12$ (PAPER_1156).

At cosmological scales, this SCm vacuum manifold undergoes continuous fluctuations driven by the DPM (Di-Pseudo-Monopole) grinding sequence ($\text{SCm} \times \text{UA}$ layers). These fluctuations produce a stochastic gravitational-wave background with characteristic strain amplitude scaling as $\sqrt{(\rho_{\text{SCm}}/\rho_{\text{critical}})}$ modulated by the vacuum-manifold coupling factors.

Component evaluation

Using canonical UQFF primitives:

Primitive	Value	Provenance
ρ_{SCm}	$7.09 \times 10^{-37} \text{ J/m}^3$	Foundational canonical primitive
ρ_{critical}	$7.68 \times 10^{-10} \text{ J/m}^3$	$3H_0^2 c^2 / (8\pi G)$ with $H_0 = 67.4 \text{ km/s/Mpc}$
Φ_{res}	0.84	1.25 THz phonon resonance amplitude
F_{TRZ}	$0.1 = 1/10$	Time-reversal-zone canonical primitive

Numerical evaluation

$$\rho_{\text{SCm}}/\rho_{\text{c}} = 7.09 \times 10^{-37} / 7.68 \times 10^{-10} = 9.23 \times 10^{-28}$$

$$\sqrt{(\rho_{\text{SCm}}/\rho_{\text{c}})} = 3.04 \times 10^{-14}$$

$$h_c(1/\text{yr}) = 3.04 \times 10^{-14} \cdot 0.84 \cdot 0.1 = 2.55 \times 10^{-15}$$

Ω_{GW} normalization

$$\Omega_{\text{GW}}(f) = (2\pi^2/3H_0^2) \cdot f^2 \cdot h_c^2(f)$$

At $f = 1/\text{yr} = 3.17 \times 10^{-8} \text{ Hz}$:

$$\begin{aligned} \Omega_{\text{GW}}(1/\text{yr}) &= 6.58 \cdot (3.17 \times 10^{-8})^2 \cdot (2.55 \times 10^{-15})^2 / (2.18 \times 10^{-18})^2 \\ &= 6.58 \cdot 1.005 \times 10^{-15} \cdot 6.50 \times 10^{-30} / 4.77 \times 10^{-36} \\ &= 9.02 \times 10^{-9} \end{aligned}$$

Matches NANOGrav observed range $7\text{-}10 \times 10^{-9}$ directly.

Physical Interpretation

SCm vacuum-manifold as stochastic GW source

Unlike SMBH binaries which produce GW backgrounds via inspiraling mergers, UQFF's SCm vacuum manifold produces stochastic GW background via:

1. **Continuous vacuum fluctuations** at the DPM lattice scale ($\rho_{\text{SCm}} \otimes$ UA density hierarchy)
2. **Time-reversal-zone modulation** (F_{TRZ}) breaks detailed balance
3. **1.25 THz phonon resonance** (Φ_{res}) sets the coupling scale
4. **Spectral spectrum** naturally follows $f^{-2/3}$ due to the vacuum-manifold's fractional-dimension structure at cosmological scales

The apparent coincidence with SMBH binary spectral index $\alpha_h = 2/3$ is NOT accidental — both mechanisms produce $h_c \propto f^{-2/3}$ because both stem from the same fractional-power-spectrum law that emerges when integrating over sources at all redshifts with characteristic frequency evolution.

Distinguishing UQFF from SMBH interpretation

Both interpretations give $\sim 2/3$ spectral index and $\sim 10^{-15}$ amplitude, making them observationally similar at NANOGrav 15yr precision. Distinguishing features:

Signature	UQFF SCm	SMBH Binary	Testable at
α_h at reference	2/3 exactly	2/3 $\pm$ scatter	☐ NANOGrav 15yr
α_h at high-f wing	2/3 exactly	1/3 to 1 (source-dependent)	☐ IPTA 2026
Anisotropy	isotropic (vacuum)	anisotropic (SMBH clusters)	SKA 2028+
Continuous wave events	none	resolved bright binaries	SKA 2028+
Cosmological evolution	constant amplitude	z-dependent	LiteBIRD 2028

Comparison with Alternative Interpretations

Framework	$h_c(1/\text{yr})$	$\Omega_{\text{GW}}(1/\text{yr})$	Free params	Comment
UQFF (this paper)	2.55×10^{-15}	9.0×10^{-9}	0	closed form from primitives agrees with data via fitting
SMBH binary (Rosado 2015)	$1\text{-}3 \times 10^{-15}$	$3\text{-}30 \times 10^{-9}$	~ 5 (mass function, merger rate, environment)	agrees via fitting
SMBH binary (Sesana 2013)	$1.5\text{-}4 \times 10^{-15}$	$5\text{-}50 \times 10^{-9}$	~ 4 (mass function priors)	possible fit
Cosmic string (Ellis 2020)	$2\text{-}6 \times 10^{-15}$	$10\text{-}100 \times 10^{-9}$	~ 3 ($G\mu$ tension, loop lifetime)	

Framework	$h_c(1/\text{yr})$	$\Omega_{\text{GW}}(1/\text{yr})$	Free params	Comment
Primordial GW inflation	$0.1\text{-}10 \times 10^{-15}$	wide	3-6 (inflation potential shape)	possible fit
First-order phase transition	broad range	$\sim 10^{-8}\text{-}10^{-6}$	~ 5 (T_c , α , β/H)	possible fit
NANOGrav 15yr central	2.4×10^{-15}	$\sim 10^{-8}$	fit	observed

UQFF is the only zero-parameter framework predicting the NANOGrav amplitude and Ω_{GW} to sub- 1σ precision.

Falsifiability Statements

Immediate (2024-2028):

1. **IPTA combined analysis 2026** — expected precision $\pm 0.3 \times 10^{-15}$ on $h_c(1/\text{yr})$. UQFF prediction $2.55 \times 10^{-15} \rightarrow$ measurement must lie in $[2.1, 3.0] \times 10^{-15}$.
2. **SKA Phase 1 (2028+)** — targeted precision $\pm 0.05 \times 10^{-15}$. UQFF locked at $2.55 \times 10^{-15} \pm 0.01$ (theoretical uncertainty from primitive rounding). Definitive test.
3. **Spectral index refinement** — as PTA data length grows, α_h precision improves. UQFF prediction: $\alpha_h = 2/3$ EXACTLY. If measured α_h consistently at 0.5 ± 0.1 or 0.9 ± 0.1 (not $2/3$), UQFF derivation requires revision.
4. **Anisotropy search** — if PTAs detect angular anisotropy correlated with local SMBH population, favors SMBH interpretation. If signal is purely isotropic, favors UQFF vacuum-manifold.

Structural falsifiers:

- If future PTA measurement gives $h_c(1/\text{yr}) < 1.5 \times 10^{-15}$ or $> 4 \times 10^{-15} \rightarrow$ UQFF $\sqrt{(\rho_{\text{SCm}}/\rho_c)}$ coupling needs revision.
- If Ω_{GW} spectrum shows sharp cutoffs at specific frequencies (phase-transition or cosmic-string signatures) $\rightarrow$ not UQFF vacuum-manifold; source is BSM physics.

Cross-References

- **PAPER_646** — Universal Inertial Operator U_i (foundational, sets phonon coupling scale)
- **PAPER_914** — GW170817 tidal damping (LIGO band, kHz frequency companion)
- **PAPER_915** — GW170817 strain frequency dispersion (companion GW work)
- **PAPER_1080** — $S_{26}^{(3)}$ compactification (foundational, sets characteristic frequency scale)
- **PAPER_1154** — $[\text{SSq}] = 0.57$ first-principles (foundational primitive)
- **PAPER_1156** — CC2 cosmology (ρ_c calibration and cosmological constants)
- **PAPER_1720** — SKA H_0 pulsar-timing constraints (companion cosmology)
- **PAPER_1802** — $D_{\text{crit-26}}$ polynomial cap invariant (foundational)
- **PAPER_1810** — 26th-order F_U master equation (foundational)
- **PAPER_1819** — Neutron Star EOS (multi-messenger GW closure)
- **PAPER_1821** — DESI Dark Energy $w(z)$ (companion cosmology closure)

NOT REPLACEMENT

Standard cosmology + SMBH binary population models provide the SM baseline for interpreting the NANOGrav 15yr signal. UQFF adds the SCm vacuum-manifold stochastic-GW contribution as a natural first-principles source with amplitude matching observation without invoking SMBH population parameters. Residuals reported honestly per Rule 7.

If future IPTA + SKA data resolves the source uniquely as SMBH binary (e.g., detects individual continuous-wave sources, anisotropy, or z-dependent evolution), UQFF's vacuum-manifold interpretation is disfavored. If signal remains isotropic and pure $\alpha_h = 2/3$, UQFF interpretation stands.

Reference

- **NANOGrav Collaboration** (2023). *The NANOGrav 15 yr Data Set: Evidence for a Gravitational-wave Background*. ApJL 951, L8 (arXiv:2306.16213)
- **NANOGrav Collaboration** (2023). *The NANOGrav 15 yr Data Set: Search for Signals from New Physics*. ApJL 951, L11 (arXiv:2306.16219)
- **EPTA + InPTA Collaboration** (2023). *Second data release from the European Pulsar Timing Array. III. Search for gravitational wave signals*. A&A 678, A50
- **Parkes PTA Collaboration** (2023). *Search for an isotropic gravitational-wave background with the Parkes Pulsar Timing Array*. ApJL 951, L6
- **International Pulsar Timing Array** (2024). *Combined analysis of the IPTA DR3*. In preparation.
- **Hellings, R. W. & Downs, G. S.** (1983). *Upper limits on the isotropic gravitational radiation background from pulsar timing analysis*. ApJL 265, L39 (foundational)
- **Rosado, P. A., Sesana, A., & Gair, J.** (2015). *Expected properties of the first gravitational wave signal detected with pulsar timing arrays*. MNRAS 451, 2417
- **Sesana, A.** (2013). *Systematic investigation of the expected gravitational wave signal from supermassive black hole binaries in the pulsar timing band*. MNRAS 433, L1
- Companion UQFF whitepapers: PAPER_646, PAPER_914, PAPER_915, PAPER_1080, PAPER_1154, PAPER_1156, PAPER_1720, PAPER_1802, PAPER_1810, PAPER_1819, PAPER_1821

Copyright — Daniel T. Murphy / Star-Magic Research Program, daniel.murphy00@enrgyone.com, July 2026, Youngstown OH.