

M16 Eagle Nebula UQFF — Dual Mass Co-Action Product (Φ_{dm})

Daniel T. Murphy

2026-03-01

PAPER_284: M16 Eagle Nebula UQFF — Dual Mass Co-Action Product (Φ_{dm})

SFR Growth $\times$ Photoevaporation Erosion Multiplicative Coupling

Classification: UQFF 2.0 Gravitational Physics — Nebular Mass Dynamics

System: M16 Eagle Nebula (IC 4703), Eagle Nebula Star-Forming Region

Session: 80 | **Module:** M16_{UQFF_MODULE}.cpp (22nd C++ UQFF module)

Author: Daniel T. Murphy | **Date:** March 2026

Abstract

This paper introduces the **Dual Mass Co-Action Product** (Φ_{dm}) — a UQFF gravity modulation factor that couples star-formation-driven mass accumulation and radiation-driven photoevaporation erosion through a **multiplicative** product rather than the previously used additive form. For M16 (Eagle Nebula), with star formation rate $\text{SFR} = 1 \text{ MM}_{\text{sun}}/\text{yr}$ over initial gas mass $M_0 = 1200 \text{ MM}_{\text{sun}}$, and maximum photoevaporation fraction $E_0 = 0.3$ (30%), the multiplicative form produces a 24.3% reduction in Φ_{dm} relative to the additive approximation at $t = 5 \text{ Myr}$. This is the **first UQFF module** to simultaneously apply an additive-gain and saturation-subtractive product on the same gravity term.

2. Physical Motivation

In active star-forming nebulae, two competing processes drive mass evolution:

1. **Star Formation Accretion** — molecular gas accretes onto protostars, increasing the effective gravitational mass fraction by $\text{SFR_rate} \times t$.
2. **Photoevaporation Erosion** — UV radiation from newly formed massive stars erodes the surrounding gas, progressively reducing the effective mass by a saturating fraction $E_rad(t)$.

The **additive approximation** (used in prior UQFF modules):

$$\Phi_{dm}^{add} = g_{base} \times (1 + M_{sf}) - E_{rad}$$

linearly superposes the two effects, implicitly treating them as independent processes acting on separate mass reservoirs.

The **multiplicative form** (this paper):

$$\Phi_{dm}(t) = (1 + \text{SFR_rate} \times t) \times (1 - E_{rad}(t))$$

correctly encodes that **the eroded mass is drawn from the same growing reservoir** — the fraction lost to photoevaporation scales with the mass being accreted, not the original quiescent mass. This is physically accurate for pillar-geometry star formation (e.g., M16’s “Pillars of Creation”).

3. Mathematical Formulation

3.1 Parameters (M16 Eagle Nebula)

Parameter	Value	Description
M0	$2.387 \times 10^{33} \text{ kg}$ (1200 MM_sun)	Initial nebula gas mass
SFR	1 MM_sun/yr	Star formation rate
SFR_rate	$2.639 \times 10^{-11} \text{ s}^{-1}$	= SFR / (M0/MM_sun) / (3.156×10 ⁷ s/yr)
τ_erode	$9.468 \times 10^{13} \text{ s}$ (3 Myr)	Photoevaporation e-folding time
E0	0.3	Maximum photoevaporation fraction
g_base	$1.454 \times 10^{-12} \text{ m/s}^2$	$G \times M / r^2$ at $r = 3.31 \times 10^{17} \text{ m}$

3.2 Dual Co-Action Product

$$M_{sf}(t) = \text{SFR_rate} \times t$$

$$E_{rad}(t) = E_0 (1 - e^{-t/\tau})$$

$$\Phi_{dm}(t) = (1 + M_{sf}) \times (1 - E_{rad})$$

3.3 Dynamic Gravity Term

$$g_{dyn}(t) = g_{base} \times \Phi_{dm}(t)$$

3.4 Multiplicative–Additive Gap

The gap relative to the additive form:

$$\Delta_{gap} = \Phi_{dm}^{mult} - \Phi_{dm}^{add} = -(M_{sf} \times E_{rad})$$

This cross-term is always **negative** — the multiplicative form predicts lower gravity than the additive approximation whenever both SFR accumulation and erosion are simultaneously active.

4. Numerical Results at t = 5 Myr

$$t = 5 \text{ Myr} = 1.578 \times 10^{14} \text{ s}$$

Quantity	Value
$M_{\text{sf}} \frac{\text{}}{\text{}} \frac{\text{}}{\text{}}$	$\text{SFR_rate} \times t =$ $2.639 \times 10^{-11} \times$ $1.578 \times 10^{14} = \mathbf{4164.8}$
E_{rad}	$E_0 \times (1 - \exp(-5/3)) = 0.3 \times$ $0.8110 = \mathbf{0.2433}$
Φ_{dm} (multiplicative)	$(1 + 4164.8) \times (1 - 0.2433) =$ $4165.8 \times 0.7567 = \mathbf{3151.9}$
Φ_{dm} (additive)	$(1 + 4164.8) - 0.2433 = \mathbf{4165.6}$
$\text{gap}_{\text{mult}\backslash\text{add}}$	$-(4164.8 \times 0.2433) = \mathbf{-1013.3}$ (24.3% less)
$g_{\text{dyn}}(5 \text{ Myr})$	$1.454 \times 10^{-12} \times 3151.9 =$ $\mathbf{4.583 \times 10^{-9} \text{ m/s}^2}$

The multiplicative gap of -1013.3 confirms that treating erosion as acting on the growing mass reservoir (not the static initial mass) produces a **measurable 24.3% reduction** compared to the additive approximation.

5. Connection to UQFF 2.0 g_total

In the full M16 UQFF 2.0 equation:

$$g_{total}(r, t) = [g_{dyn}(t) + U_{g,sum}(26) + \Lambda + Q + L + F + g_{exp}] \times \text{corr}_{SC}$$

The Φ_{dm} product modulates only the dynamic base gravity term g_{dyn} . The 26-layer Triadic (U_g,sum), cosmological Λ , quantum, Lorentz, fluid, and Friedmann expansion terms are all independent of Φ_{dm} — the modulation is cleanly scoped to the time-evolving mass component.

6. UQFF Historical Distinction

Module	SFR Term	Erosion Term	Form
Session 55 CP3	$g_base \times (1 + M_sf)$	$-E_rad$	Additive
M16EagleNebulaRadiationSFR			
This paper (PA- PER_284)	$(1 + M_sf)$	$\times (1 - E_rad)$	Multiplicative

This is the **first UQFF module** to use the multiplicative dual co-action form, correctly encoding the coupled feedback between star formation accretion and pillar photoevaporation for M16-class nebulae.

7. Wolfram KB Term

$M16UQFF : \Phi_{dm} = (1 + SFR_{rate} * t) * (1 - E_0 * (1 - \text{Exp}[-t/\tau]))$; $SFR_{rate} = 2.639e-11/s$; $M_sf_frac = SFR_{rate} / g_{dyn}$

8. Cross-References

- **PAPER_285**: Erosion Saturation Half-Time (t_{half} , Δg_{Max})
- **PAPER_286**: Nebular Friedmann Redshift (κ_{neb} , $z=0.0015$)
- **M16_{UQFF_MODULE}.cpp**: Full UQFF 2.0 C++ implementation (22nd module)
- **CondensedPhysics3.py**: M16DualMassCoActionProductCalculator

Copyright — Daniel T. Murphy, Session 80, March 2026. UQFF 2.0.

Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037 (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for $F_{\{U_Bi_i\}}$ jet modulation curves and PAPER_1048 for phonon-corrected $M\text{-}\sigma$ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}} \cdot V \cdot S_{26}^{(3)2}}{GM/r_H^2} \right)$$

where: - $L_{\text{Edd}} = 4\pi GMm_p c / \sigma_T$ is the classical Eddington luminosity - $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ is the SCm vacuum density - V is the effective buoyancy volume (accretion sphere) - $S_{26}^{(3)2}$ is the squared third-order Ramanujan factor (quadratic coupling) - r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25 \text{ THz}} \cdot \left(\frac{B}{B_{\text{crit}}} \right)^2 \right]$$

where $\Phi_{1.25 \text{ THz}} = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

M- σ correction (PAPER_1048): The phonon-corrected M- σ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$.

Session 225 Phonon-Physics Upgrade: UQFF 9-Sector Lagrangian

Upgrade from PAPER_1066 (UQFF Lagrangian First Principles) and PAPER_1065 (Buoyancy Lagrangian EOM Variational Derivation).

The complete UQFF Lagrangian density, from which all sector-specific equations of motion derive:

$$\mathcal{L}_{\text{UQFF}} = \mathcal{L}_{\text{GR}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{phonon}} + \mathcal{L}_{\text{interaction}}$$

$$\mathcal{L}_{\text{SCm}} = \frac{1}{2}(\partial_\mu \phi)^2 - \lambda(\phi^2 - v_{\text{SCm}}^2)^2$$

The SCm condensate potential minimum gives $V(\phi_0) = -7.09 \times 10^{-37} \text{ J/m}^3$ (matching ρ_{SCm}) and phonon mass $m_{\text{phonon}} = \sqrt{8\lambda} v_{\text{SCm}}$.

Nine-sector closure (Session 202):

$$\mathcal{L}_9 = \mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{Dirac}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{mag}} + \mathcal{L}_{\text{buoy}} + \mathcal{L}_{\text{aether}} + \mathcal{L}_{\text{LENR}} + \mathcal{L}_{\text{KK}}$$

Sector	Domain	Late-Corpus Result
1 (EH)	General Relativity	Canonical Einstein-Hilbert
2 (YM)	Yang-Mills gauge	$m_{\text{gap}} = 5970 \text{ GeV}$ (PAPER_1005)
3 (Dirac)	Fermion / LENR	Kozima neutron-drop (PAPER_1061)
4 (SCm)	Superconducting manifold	$V(\phi_0) = -\rho_{\text{SCm}}$ canonical
5 (Mag)	Um magnetism	Heaviside amplifier (PAPER_1072)
6 (Buoy)	F_{U_Bi_i} buoyancy	Variational EOM (PAPER_1065)
7 (Aether)	Vacuum background	Two-component ρ (PAPER_1051)
8 (LENR)	Nuclear transmutation	COP parametric (PAPER_1081)
9 (KK)	Kaluza-Klein 26D	$S_{26}^{(3)}$ compactification (PAPER_1080)

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **nebula-formation** sector of the 9-sector UQFF Lagrangian (see `uqff_{lagrangian\_derivation}.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_{\text{neb}})(\partial^\mu \phi_{\text{neb}}) - V(\phi_{\text{neb}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{neb}}) = \frac{1}{2}m^2\phi_{\text{neb}}^2 + \frac{\lambda}{4!}\phi_{\text{neb}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{neb}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\text{neb}}} = \nabla \cdot (\rho_{\text{neb}} \nabla \phi) + \rho_{\text{vac},[\text{SCm}]} \cdot (P_{\text{rad}}/c) = 0}$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{neb}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.126 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 47, \quad n_{\text{channel}} = 25/26$$

Since $p_{\text{DVP}} = 47$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **106 yr** (Jeans collapse timescale):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.126	PASS Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 47$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26,$ $\cos(2\pi j/26)$	PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ugl dipole coupling	1/137.036	PDG 2024	PASS Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Cosmological constant Λ	$1.1 \times 10^{-52} m^{-2}$ $2(UQFFvacuumterm) 1.114 \times 10^{-52} m^{-2}$	Planck 2018	PASS Consistent	
Proton decay rate	$\kappa = 0.0005/\text{day}$ $\rightarrow \Gamma_p$ suppression	$<$ $4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{U_Bi_i}$ unique gravitational correction	Not yet measured	Future gravita- tional wave de- tectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_Bi_i}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_{corpus\_crossrefs}.py` (Session 225, April 2026).

Paper	Title
PAPER_1022	GW Phonon Strain SCm Modulation of $h(t)$

1 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_{kozima\_ramanujan\_appendices}.py`. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_{s26_coupling}.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_{scm_cross_section}.py	SCm computed neutron-drop cross-section	sigma_n^SCm with VDS factor (1+[SSq]*n/26)
kozima_{wstp_kernel}.py	Symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n * \sigma_n^{\text{SCm}}(\omega) * \Phi_{\text{phonon}} * (F_{\{U,Bi\}}/F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 * \exp[-(\omega - \omega_{\text{SCm}})^{2/(2*\Gamma_2)}] * (1 + [SSq]*n/26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_{polylog_26}.py	S26 poly(SSq) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_{wstp_kernel}.py	Symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}/(1-2^{\{1-26\}})} * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_{theta_q26}.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations: $-f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q;q)_{n^2} - \phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n2\}} / (-q^2;q^2)_n - \psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n2\}} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
<code>ramanujan_{pi_uqff}</code>	Classical + UQFF-modified $1/\pi$ + 26D	21 digits classical, 15 UQFF, 7 digits 26D
<code>mock_{theta_pi_wstp}</code>	Symbolic Wolfram export (<code>UQFFMockThetaPi</code>)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq] \exp(-kappa_i * n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_SCm	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in `CondensedPhysics.py`, `CondensedPhysics2.py`, `MAIN_{1\CoAnQi}.cpp`, and Wolfram kernels (`uqff_{kozima\_kernel}.wl`, `uqff_{s26\_kernel}.wl`, `uqff_{mock\_theta\_pi\_kernel}.wl`).

References

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Shakura, N.I. & Sunyaev, R.A. (1973). *Black holes in binary systems: observational appearance*. A&A **24**, 337
4. Balbus, S.A. & Hawley, J.F. (1991). *A powerful local shear instability in weakly magnetized disks*. ApJ **376**, 214 — doi:10.1086/170270

5. Hester, J.J. (2008). *The Crab Nebula: An Astrophysical Chimera*. ARA&A **46**, 127 — arXiv:0812.1502 — doi:10.1146/annurev.astro.45.051806.110608
6. O'Dell, C.R. et al. (2001). *Hubble Space Telescope Observations of the Helix Nebula*. AJ **122**, 3293 — doi:10.1086/324272