

Crab Nebula MUGE – Expanding Supernova Remnant with Pulsar Wind

Daniel T. Murphy

2025-01-01

PAPER_745: Crab Nebula MUGE — Expanding Supernova Remnant with Pulsar Wind

Author: Daniel T. Murphy

Framework: Universal Quantum Field Superconductive Framework (UQFF)

Session: 180 continuation | v5.38

Date: 2025

CP4 Class: #329 — CrabNebulaExpandingMUGECalculator

Abstract

The Crab Nebula (M1) is the remnant of SN 1054, powered by a central pulsar (PSR B0531+21) at the highest rotational energy loss in the known neutron star population. Its expanding ejecta ($r(t) = r_0 + v_r \cdot t$) creates a time-dependent gravitational environment unlike any static system. This paper derives the Crab Nebula MUGE incorporating: the expanding radius $r(t)$, the pulsar wind term F_{wind} , and the nebular magnetic energy M_{mag} . The UQFF framework captures both the mechanical expansion and the magnetospheric energy injection that power the synchrotron nebula.

1. Introduction

The Crab Nebula is a pulsar wind nebula (PWN) — one of the brightest objects in X-rays and gamma-rays. The central pulsar ($P = 33 \text{ ms}$, $\dot{P} = 4.2 \times 10^{-13} \text{ s/s}$) injects $\sim 5 \times 10^{31} \text{ W}$ of spin-down power into the nebulae through a relativistic wind. The expanding ejecta shell moves at $v_r \sim 1500 \text{ km/s}$, with angular size growing observably over human timescales. This expanding geometry requires $r(t)$ to appear explicitly in the MUGE denominator, making the Crab the canonical example of a time-dependent gravitational radius.

Crab Nebula parameters: - Distance = 2.0 kpc - Age $\approx$ 970 yr (SN 1054) - r_0 (initial remnant radius) $\approx 3 \times 10^{15}$ m - $r_{\text{current}} \approx 5.6$ ly = 5.3×10^{16} m - $v_r = 1500$ km/s = 1.5×10^6 m/s - $M_{\text{ejecta}} \approx 4\text{--}5 M_{\text{sun}}$ - $B_{\text{nebula}} \approx 10^{-4}$ T (filamentary magnetic field) - $L_{\text{pulsar}} = 5 \times 10^{31}$ W (spin-down luminosity) - $P_{\text{spin}} = 33$ ms

2. Crab Nebula MUGE

$$\begin{aligned}
g_{\text{Crab}}(r, t) = & (G * M) / r(t)^2 * (1 + H(z) * t) * (1 - B(t) / B_{\text{crit}}) \\
& + (U_g1 + U_g2 + U_g3 + U_g4) \\
& + U_i \\
& + (\text{Lambda} * c^2 / 3) \\
& + (\hbar / \sqrt{\Delta x * \Delta p}) * \text{integral}(\psi * H * \psi dV) * (2\pi / t_{\text{Hubble}}) \\
& + \rho_{\text{ejecta}} * V * g \\
& + F_{\text{wind}}[\text{pulsarwind} \text{---} \text{NEW}] \\
& + M_{\text{mag}}[\text{magneticenergy} \text{---} \text{NEW}]
\end{aligned}$$

3. Time-Dependent Radius $r(t)$

The expanding radius is the defining feature of the Crab MUGE:

$$\begin{aligned}
r(t) &= r_0 + v_r * t \\
r_0 &= \text{initialradiusatSNexplosion}(m) \\
v_r &= \text{expansionvelocity} = 1.5 \times 10^6 \text{ m/s} \\
t &= \text{timesinceSN1054}(s)
\end{aligned}$$

For $t = 970$ yr = 3.06×10^{10} s:

$$\begin{aligned}
r(970 \text{ yr}) &= 3 \times 10^{15} + 1.5 \times 10^6 * 3.06 \times 10^{10} \\
r(970 \text{ yr}) &= 4.6 \times 10^{16} \text{ m} = 4.9 \text{ ly}
\end{aligned}$$

Gravitational deceleration:

$$\begin{aligned}
g_{\text{grav}}(t) &= G * M_{\text{ejecta}} / r(t)^2 \\
g_{\text{grav}}(970 \text{ yr}) &= 6.674 \times 10^{-11} * (4 * 1.989 \times 10^{30}) / (4.6 \times 10^{16})^2 \\
g_{\text{grav}} &= 5 \times 10^{-22} \text{ m/s}^2 (\text{extremelyweak} \text{---} \text{expansiondominates})
\end{aligned}$$

4. F_wind — Pulsar Wind Term

The pulsar wind carries relativistic particles outward, creating an inward reaction force equivalent to:

$$F_{wind} = L_{pulsar} / (4 * \pi * r(t)^2 * c * M_{ejecta})$$

$$L_{pulsar} = 5 \times 10^{31} W$$

$$r(t) = \text{current nebular radius}$$

$$c = 3 \times 10^8 m/s$$

$$F_{wind} = 5 \times 10^{31} / (4 * \pi * (4.6 \times 10^{16})^2 * 3 \times 10^8 * 4 \times 1.989 \times 10^{30})$$

$$F_{wind} = 4.8 \times 10^{-3} m/s^2 (\text{small but physically significant over } 970 \text{ yr})$$

Cumulative momentum injection over nebula lifetime:

$$\Delta p_{wind} = L_{pulsar} * t / c = 5 \times 10^{31} * 3 \times 10^4 / 3 \times 10^8 = 5 \times 10^3 kg * m/s$$

5. M_mag — Magnetic Energy Term

The nebular magnetic field stores and dissipates energy, affecting dynamics:

$$M_{mag} = B^2 / (2 * \mu_0 * r(t) * \rho_{ejecta}) [\text{magnetic pressure divided by density}]$$

$$B = 10^{-4} T (\text{filamentary field})$$

$$\mu_0 = 4\pi \times 10^{-7} H/m$$

$$\rho_{ejecta} = 10^{-22} kg/m^3 (\text{current})$$

$$M_{mag} = (10^{-4})^2 / (2 * 4\pi \times 10^{-7} * 4.6 \times 10^{16} * 10^{-22})$$

$$M_{mag} = 2.7 \times 10^{-17} m/s^2$$

This is the dominant non-DPM-seeded term for the Crab, controlling the synchrotron brightness evolution.

6. UQFF Gravity Components

U_g1 : Pulsarmagneticdipole
 $\mu_dipole_PSR = 3.8 \times 10^{30} J/T (from P,)$
 $B_P SR = 3.8 \times 10^8 T (surface field)$
 U_g1 contributes at pulsar vicinity only
 U_g2 : Superconductive aether field threading nebula
 B_super aligned with pulsar rotation axis
 $U_g2 = 10^5 J/m^3 (strong near pulsar wind shock)$
 U_g3 : External galactic field at 2 kpc
 $U_g3 = G * M_{gal} / r_{gal}^2$
 U_g4 : Galactic center contribution
 $k_4 * \rho_{vac} [SCm] * (M_b h / d_g) * e^{(-\alpha)} * \cos(\pi * t_n)$

7. Temporal Evolution Summary

Time (yr)	r(t) (ly)	g_grav (m/s ²)	M_mag (m/s ²)
0	0.3	2×10^{14}	10^{16}
100	1.6	5×10^{21}	10^{17}
970	4.9	5×10^{22}	2.7×10^{17}
10,000	~50	$\sim 5 \times 10^{27}$	$\sim 10^{19}$

The magnetic term M_mag remains important relative to g_grav throughout nebula evolution.

8. Conclusion

The Crab Nebula MUGE demonstrates that time-dependent radius r(t) is essential for accurately modeling supernova remnant gravity. The F_wind pulsar injection and M_mag magnetic energy terms together dominate over classical DPM-seeded gravity within the nebula. The UQFF successfully models the transition from SN ejecta to mature PWN through its modular environmental forcing framework.

Copyright - Daniel T. Murphy, daniel.murphy00@gmail.com. UQFF Framework.
 PAPER_745, CP4 class #329. Session 180 continuation v5.38.

Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037 (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for $F_{\{U_Bi_i\}}$ jet modulation curves and PAPER_1048 for phonon-corrected $M\text{-}\sigma$ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}} \cdot V \cdot S_{26}^{(3)2}}{GM/r_H^2} \right)$$

where: - $L_{\text{Edd}} = 4\pi GMm_p c / \sigma_T$ is the classical Eddington luminosity - $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ is the SCm vacuum density - V is the effective buoyancy volume (accretion sphere) - $S_{26}^{(3)2}$ is the squared third-order Ramanujan factor (quadratic coupling) - r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25 \text{ THz}} \cdot \left(\frac{B}{B_{\text{crit}}} \right)^2 \right]$$

where $\Phi_{1.25 \text{ THz}} = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

M- σ correction (PAPER_1048): The phonon-corrected M- σ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$.

Session 225 Phonon-Physics Upgrade: UQFF 9-Sector Lagrangian

Upgrade from PAPER_1066 (UQFF Lagrangian First Principles) and PAPER_1065 (Buoyancy Lagrangian EOM Variational Derivation).

The complete UQFF Lagrangian density, from which all sector-specific equations of motion derive:

$$\mathcal{L}_{\text{UQFF}} = \mathcal{L}_{\text{GR}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{phonon}} + \mathcal{L}_{\text{interaction}}$$

$$\mathcal{L}_{\text{SCm}} = \frac{1}{2}(\partial_\mu \phi)^2 - \lambda(\phi^2 - v_{\text{SCm}}^2)^2$$

The SCm condensate potential minimum gives $V(\phi_0) = -7.09 \times 10^{-37} \text{ J/m}^3$ (matching ρ_{SCm}) and phonon mass $m_{\text{phonon}} = \sqrt{8\lambda} v_{\text{SCm}}$.

Nine-sector closure (Session 202):

$$\mathcal{L}_9 = \mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{Dirac}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{mag}} + \mathcal{L}_{\text{buoy}} + \mathcal{L}_{\text{aether}} + \mathcal{L}_{\text{LENR}} + \mathcal{L}_{\text{KK}}$$

Sector	Domain	Late-Corpus Result
1 (EH)	General Relativity	Canonical Einstein-Hilbert
2 (YM)	Yang-Mills gauge	$m_{\text{gap}} = 5970 \text{ GeV}$ (PAPER_1005)
3 (Dirac)	Fermion / LENR	Kozima neutron-drop (PAPER_1061)
4 (SCm)	Superconducting manifold	$V(\phi_0) = -\rho_{\text{SCm}}$ canonical
5 (Mag)	Um magnetism	Heaviside amplifier (PAPER_1072)
6 (Buoy)	$F_{\{U_{Bi_i}\}}$ buoyancy	Variational EOM (PAPER_1065)
7 (Aether)	Vacuum background	Two-component ρ (PAPER_1051)
8 (LENR)	Nuclear transmutation	COP parametric (PAPER_1081)
9 (KK)	Kaluza-Klein 26D	$S_{26}^{(3)}$ compactification (PAPER_1080)

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_{lagrangian\_derivation}.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac},[\text{SCm}]} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac},[\text{SCm}]} \cdot \phi_{\text{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}}/c^2) \phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.065 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 101, \quad n_{\text{channel}} = 18/26$$

Since $p_{\text{DVP}} = 101$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10⁴ yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The tanh saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.065	PASS Threshold-consistent
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\text{DVP}} = 101$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26,$ $\cos(2\pi j/26)$	PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ugl dipole coupling	1/137.036	PDG 2024	PASS Consistent

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Cosmological constant Λ	1.1×10^{-52} m^2 (UQFF vacuum term)	1.114×10^{-52} m^2	Planck 2018	PASS Consistent
Proton decay rate	$\kappa = 0.0005/\text{day}$ $\rightarrow \Gamma_p$ suppression	$<$ $4.17 \times 10^{35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{U_Bi_i}$ unique gravitational correction	Not yet measured	Future gravita- tional wave de- tectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_Bi_i}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_corpus_crossrefs.py` (Session 225, April 2026).

Paper	Title
PAPER_1000	NS Merger F_{U_Bi} Strain Suppression & BCS Gap
PAPER_1011	GW170817 NS Merger $F_{U_Bi_i}$ 66.7% Strain Reduction
PAPER_1012	GW190425 Upgraded $F_{U_Bi_i}$ with S26(3)
PAPER_1022	GW Phonon Strain SCm Modulation of $h(t)$
PAPER_1002	AGN Buoyancy-Corrected Eddington Luminosity
PAPER_1009	3C273 AGN $F_{U_Bi_i}$ Jet Modulation
PAPER_1010	TON618 AGN $F_{U_Bi_i}$ Jet Modulation
PAPER_1037	AGN Buoyancy Jet Calculator — SCm Jet Launching
PAPER_1048	M-Sigma Phonon-Corrected Relation
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression
PAPER_1021	Pulsar Timing Phonon TOA Residual
PAPER_1047	Type Iax Supernova Buoyancy Reversal

Paper	Title
PAPER_1050	MUGE F_{U_Bi_i} Unified 9-System Synthesis
PAPER_1075	3D Volumetric MUGE Gravitational Field Generator

15 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade_{kozima\_ramanujan\_appendices}.py`. For detailed derivations, see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_{s26_coupling}.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_{scm_cross_section}.py	SCm-hyperplated neutron-drop cross-section	sigma_n^SCm with VDS factor (1+[SSq]*n/26)
kozima_{wstp_kernel}.py	symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_neutron^{SCm} = N_n * sigma_n^{SCm}(\omega) * Phi_phonon * (F_{U,Bi}/F_U - 1)$ where $sigma_n^{SCm}(\omega, n) = sigma_0 * exp[-(\omega - \omega_{SCm})^{2/(2*Gamma2)}] * (1 + [SSq]*n/26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_{polylog_{s26}}.py	R26 polylog([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_{wstp_kernel}.py	symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_26(z) = Li_26(z) = eta_26(z)/(1-2^{\{1-26\}}) + 2^{\{1-26\}}/(1-2^{\{1-26\}}) * Li_26(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
<code>mock_{theta_q26}.py</code>	<code>f_26(q)</code> , <code>phi_26(q)</code> , <code>psi_26(q)</code> q-series	Proper q-Pochhammer (a;q)_n

Core equations: $-f_{26}(q) = \text{Sum}_{\{n=0\}^{\wedge}\{25\}} q^{\{n\}2} / (-q;q)_{n^{\wedge}2} - \phi_{26}(q) = \text{Sum}_{\{n=0\}^{\wedge}\{25\}} q^{\{n\}2} / (-q^2;q^2)_n$ - $\psi_{26}(q) = \text{Sum}_{\{n=1\}^{\wedge}\{26\}} q^{\{n\}2} / (q;q^{\wedge}2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
<code>ramanujan_{pi_uqff}.py</code>	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
<code>mock_{theta_pi_wstp9801}.py</code>	Symbolic Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \text{Sum } R_n * (1103+26390n) * W_{26}(n) / C_{26}$ where $W_{26}(n) = \text{Prod}_{\{i=1\}^{\wedge}\{26\}} [1 + [SSq]^{\exp(-kappa_i*n/26)}]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
omega_SCm	$2*\pi \times 1.25 \text{ THz}$	SCm phonon resonance
sigma_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in *CondensedPhysics.py*, *CondensedPhysics2.py*, *MAIN_{1_CoAnQi}.cpp*, and Wolfram kernels (*uqff_{kozima_kernel}.wl*, *uqff_{s26_kernel}.wl*, *uqff_{mock_theta_pi_kernel}.wl*).

References

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Fabian, A.C. (2012). *Observational Evidence of Active Galactic Nuclei Feedback*. ARA&A **50**, 455 — arXiv:1204.4114 — doi:10.1146/annurev-astro-081811-125521
4. McNamara, B.R. & Nulsen, P.E.J. (2007). *Heating Hot Atmospheres with Active Galactic Nuclei*. ARA&A **45**, 117 — arXiv:0709.4098 — doi:10.1146/annurev.astro.45.051806.110625
5. Heckman, T.M. & Best, P.N. (2014). *The Coevolution of Galaxies and Supermassive Black Holes*. ARA&A **52**, 589 — arXiv:1403.4620 — doi:10.1146/annurev-astro-081913-035722
6. Kaspi, V.M. & Beloborodov, A.M. (2017). *Magnetars*. ARA&A **55**, 261 — arXiv:1703.00068 — doi:10.1146/annurev-astro-081915-023329
7. Goldreich, P. & Julian, W.H. (1969). *Pulsar Electrodynamics*. ApJ **157**, 869 — doi:10.1086/150119
8. Riess, A.G. et al. (1998). *Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant*. AJ **116**, 1009 — arXiv:astro-ph/9805200 — doi:10.1086/300499
9. Perlmutter, S. et al. (1999). *Measurements of Omega and Lambda from 42 High-Redshift Supernovae*. ApJ **517**, 565 — arXiv:astro-ph/9812133 — doi:10.1086/307221
10. Janka, H.-T. (2012). *Explosion Mechanisms of Core-Collapse Supernovae*. ARA&A **50**, 531 — arXiv:1206.2503 — doi:10.1146/annurev-astro-081811-125815
11. Lorimer, D.R. & Kramer, M. (2004). *Handbook of Pulsar Astronomy*. Cambridge University Press
12. Hewish, A. et al. (1968). *Observation of a Rapidly Pulsating Radio Source*. Nature **217**, 709 — doi:10.1038/217709a0
13. Manchester, R.N. et al. (2005). *The Australia Telescope National Facility Pulsar Catalogue*. AJ **129**, 1993 — arXiv:astro-ph/0412641 — doi:10.1086/428488
14. Murphy, D. (2026). *Master Universal Gravity Equation (MUGE): DPM-Driven Gravity Framework*. Star-Magic Whitepaper Series — github.com/Daniel8Murphy0007/Star-Magic
15. Lattimer, J.M. & Prakash, M. (2007). *Neutron Star Observations: Prognosis for Equation of State Constraints*. Phys. Rep. **442**, 109 — arXiv:astro-ph/0612440 — doi:10.1016/j.physrep.2007.02.003
16. Demorest, P.B. et al. (2010). *A two-solar-mass neutron star measured using Shapiro delay*. Nature **467**, 1081 — arXiv:1010.5788 —

doi:10.1038/nature09466

17. Cromartie, H.T. et al. (2020). *Relativistic Shapiro delay measurements of an extremely massive millisecond pulsar*. *Nature Astron.* **4**, 72 — arXiv:1904.06759 — doi:10.1038/s41550-019-0880-2
18. Hester, J.J. (2008). *The Crab Nebula: An Astrophysical Chimera*. *ARA&A* **46**, 127 — arXiv:0812.1502 — doi:10.1146/annurev.astro.45.051806.110608
19. O'Dell, C.R. et al. (2001). *Hubble Space Telescope Observations of the Helix Nebula*. *AJ* **122**, 3293 — doi:10.1086/324272