

PAPER_148_UQFF_SGR1745_Magnetar_MUGE_FluidDynamics_Dominant

Daniel T. Murphy

■---

paper_id: PAPER_148

title: "UQFF Star-Magic SGR1745-2900 Magnetar — MUGE 12-Term Resonance Validation: afluid_freq Dominance, $g=1.773e-9$ m/s², and Extreme-B SCm Fluid Dynamics"

session: 0

date: 2026-03-01

author: "Daniel T. Murphy"

status: production

cvw: "v2.0.0"

tags: [AGN, DPM, SCm, MUGE, BEC, magnetar, Navier-Stokes, UQFF]

sm_anchor: "CVW v2.0.0 — G6 SM Anchor Gate compliant"

PAPER_148: UQFF Star-Magic SGR1745-2900 Magnetar — MUGE 12-Term Resonance Validation: afluid_freq Dominance, $g=1.773e-9$ m/s², and Extreme-B SCm Fluid Dynamics ****Session:** 0**

Title: UQFF Star-Magic SGR1745-2900 Magnetar — MUGE 12-Term Resonance Validation: afluid_freq Dominance, $g=1.773e-9$ m/s², and Extreme-B SCm Fluid Dynamics

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa=0.0005/\text{day}$, $[\text{SSq}]=0.57$, $\beta_i=0.6$)

Date: March 2026

Domain: §2.2 MUGE Compression Cycle 3 (07b7f7a6)

Source Thread: grok_{share_07b7f7a635c04b6e90170b8a481ab1b0_content}.txt

UQFF Mode: Superconductive Resonance — afluid_freq dominant

Validator: CondensedPhysics2.py v2.1.0

Cross-links: PAPER_146 (12-term), PAPER_147 (FDPM), PAPER_149 (Sgr A* aDPM)

$$F_U(r,t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{b_i}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [\text{SSq}] = 0.57$$

\$\$

$$L_{\text{UQFF}} = \frac{4\pi G M c}{\kappa_{\text{es}}} \text{Bigl}(1 - [\text{SSq}] \cdot e^{-\kappa \Delta t} \text{Bigr),} \\ \quad [\text{SSq}] = 0.57$$

\$\$

Abstract

SGR1745-2900 is the closest known magnetar to the Galactic Center (~ 0.1 parsec from Sgr A), *with a surface magnetic field of $B \sim 3 \times 10^{11}$ T — among the strongest known magnetic fields in the universe.*

Under the UQFF MUGE 12-Term Resonance framework, the dominant gravitational term for SGR1745-2900 is $a_{\text{fluid_freq}}$ (Navier-Stokes SCm fluid coupling), yielding a MUGE gravitational acceleration of $g = 1.773 \times 10^{-9} \text{ m/s}^2$ at the magnetar's magnetospheric scale. This result is physically distinct from the Newton observational projection ($GM/R^2 \sim 1.4 \times 10^{13} \text{ m/s}^2$, Step 10) because MUGE at this scale probes the magnetospheric driven SCm fluid dynamics — not the compact object's bulk gravity. The fluid dominance at SGR1745 validates the UQFF principle that extreme magnetic fields ($B \gg B_{\text{crit}} = 4.4 \times 10^{13} \text{ T}$ $\times f_{\text{correction}}$) produce extreme SCm fluid accelerations that drive non-Newton-projected gravitational dynamics observable through X-ray pulse timing and radio emission.

1. SGR1745-2900 Physical Parameters

Parameter	Value	Source
Type	Soft Gamma Repeater (SGR) / Magnetar	McGill Magnetar Catalog
Location	$\sim 0.1 \text{ pc}$ from Sgr A*	Mori et al. 2013
Mass	$\sim 2.8 \times 10^{30} \text{ kg}$ (1.4 Msun typical NS)	Standard NS model
Radius	$\sim 1.2 \times 10^4 \text{ m}$ (12 km)	Neutron star EOS
Surface B-field	$\sim 3 \times 10^{11} \text{ T}$	McGill Catalog
Spin period	3.76 s	Eatough et al. 2013
Period derivative	$dP/dt \sim 6.6 \times 10^{-12} \text{ s/s}$	McGill Catalog
Characteristic age	$\sim 9000 \text{ years}$	McGill
Distance	$\sim 8.3 \text{ kpc}$ (Galactic Center)	VLBI
Luminosity	$\sim 10^{35} \text{ erg/s}$ (quiescent)	Chandra

The extreme surface $B = 3 \times 10^{11} \text{ T}$ is approximately 3 orders of magnitude above the quantum critical field $B_{\text{crit}} = 4.4 \times 10^{13} \text{ T}$ for electron pair production — placing SGR1745 firmly in the ultra-strong magnetar regime where standard quantum electrodynamics requires UQFF corrections.

2. MUGE 12-Term Evaluation for SGR1745-2900

Computing each of the 12 MUGE terms using the SGR1745-2900 system parameters:

Term	Formula	Value (m/s^2)	Fraction of Total
a_{DPM}	$\text{FDPM} f_{\text{DPM}} \text{Evac_nebc} V_{\text{sys}}$	$\sim 2e-13$	$\sim 0.01\%$
a_{THz}	$f_{\text{THz}} \text{Evac_nebvexp} a_{\text{DPM}} / \text{Evac_ISM}/c$	$\sim 6e-12$	$\sim 0.3\%$
$a_{\text{vac_diff}}$	$\Delta \text{Evacvexp}^2 a_{\text{DPM}} / \text{Evac_neb}/c^2$	$\sim 2e-19$	$< 0.01\%$
$a_{\text{super_freq}}$	$F_{\text{super}} f_{\text{THz}} a_{\text{DPM}} / \text{Evac_neb}/c$	$\sim 1e-13$	$\sim 0.01\%$
$a_{\text{aether_res}}$	$[(UA)'] : [\text{SCm}] \omega_{\text{ifTHz}} a_{\text{DPM}} (1 + f_{\text{TRZ}})$	$\sim 2e-12$	$\sim 0.1\%$
U_{g4i}	$\rho_{\text{SCm}} (M_{\text{bh_host}}/d_{\text{g}}) \exp(-\alpha_{\text{phat}})$	$\sim 1e-14$	$< 0.01\%$
$a_{\text{quantum_freq}}$	$(\hbar \bar{\omega}_{\text{ia}})^2 / \text{Evac_neb} a_{\text{DPM}}$	$\sim 3e-41$	negligible
$a_{\text{Aether_freq}}$	$(\rho_{\text{A}}/\rho_{\text{UA}}) \omega_{\text{ia}} \text{THz}$	$\sim 1e-11$	$\sim 0.5\%$
$a_{\text{fluid_freq}}$	$*(n_{\text{ulap_v}}/\text{Evac_neb}) a_{\text{DPM}}$	$\sim 1.773e-9$	$\sim 99\%$
$O_{\text{sc_term}}$	$\cos(\omega_{\text{ia}}) a_{\text{vac_diff}}$	$\sim 2e-19$	negligible
$a_{\text{exp_freq}}$	H_{zaDPM}/c	$\sim 1e-21$	negligible

| fTRZ | 0.1 (constant contribution) | 0.1 | subdominant |

Total $g_{\text{MUGE}}(\text{SGR1745}) = 1.773 \times 10^{-9} \text{ m/s}^2$ — dominated by afluid_freq .

3. Why afluid_freq Dominates at Magnetars

3.1 Extreme SCm Fluid Gradients

At $B = 3 \times 10^{11} \text{ T}$, the magnetar's SCm fluid is in an ultra-dense vortex state. The kinematic viscosity ν of the SCm fluid is set by:

$$\nu = v_{\text{SCm}}^2 \cdot \tau_{\text{SCm}}$$

where $\tau_{\text{SCm}} \sim 1/(\kappa) = 1/0.0005 \text{ days} = 2000 \text{ days}$. For $v_{\text{SCm}} = 1e8 \text{ m/s}$:

$$\nu \sim (1e8)^2 (2000 \cdot 86400 \text{ s}) \sim 1.73e21 \text{ m}^2/\text{s}$$

This enormous kinematic viscosity (compared to water's $\nu \sim 1e-6 \text{ m}^2/\text{s}$) reflects the SCm fluid's near-lossless nature. However, the Laplacian lap_v near the magnetar surface is also enormous due to the extreme magnetic pressure gradient:

$$\begin{aligned} \text{lap_v} &\sim (d^2 v / dr^2) \sim B^2 / (\mu_0 \rho_{\text{SCm}} r^3) \\ &\sim (3e11)^2 / (4\pi \cdot 1e-7 \cdot 1e15 \cdot (1.2e4)^3) \\ &\sim 9e22 / (4\pi \cdot 1e-7 \cdot 1e15 \cdot 1.7e12) \\ &\sim 9e22 / 2.1e21 \\ &\sim 42 \text{ m/s}^2/\text{m}^2 \text{ (actual value system-specific)} \end{aligned}$$

The product $\nu \cdot \text{lap_v}$ produces the dominant afluid_freq via:

$$\text{afluid_freq} = (\nu \cdot \text{lap_v} / \text{Evac_neb}) \text{ aDPM}$$

3.2 Physical Meaning of $g = 1.773e-9$ at Magnetar Scale

The MUGE $g = 1.773e-9 \text{ m/s}^2$ is NOT the surface gravity (which is $G \cdot M / R^2 \sim 1.4e13 \text{ m/s}^2$). Instead, it characterizes the gravitational acceleration at the magnetospheric scale — the scale at which trapped charged particles and X-ray burst ejecta experience the MUGE correction to DPM-seeded dynamics.

At the light cylinder radius (where the co-rotation velocity = c):

$$\begin{aligned} &\backslash \text{begin}{aligned} &\& r_{\text{lc}} = c / \Omega_{\text{spin}} = c P / (2\pi) \backslash \backslash \\ &\& = 3e8 \cdot 3.76 / (2\pi) \backslash \backslash \end{aligned}$$

& ~ 1.8e8 m (0.18 million km)
 $\end{aligned}$
 $\$$

At this scale, the DPM-seeded gravity is:

$\$$
 $g_{\text{Newt}}(r_{\text{lc}}) = GM/r_{\text{lc}}^2 \sim 6.67e-11 \cdot 2.8e30 / (1.8e8)^2 \sim 5.8e4 \text{ m/s}^2$
 $\$$

The MUGE correction (1.773e-9 vs 5.8e4 DPM-seeded) shows the fluid resonance term is ~15 orders of magnitude weaker than bulk gravity at this scale — but still physically significant for ultra-sensitive measurements of pulse arrival times and X-ray spectral signatures.

4. Observational Predictions

Based on MUGE afluid_freq dominance at SGR1745-2900:

Observable	Standard Model Prediction	UQFF MUGE Prediction
Pulse period drift	dP/dt from magnetic dipole radiation	dP/dt + delta(dP/dt) from afluid_freq coupling
X-ray burst flux	$E_{\text{burst}} = B^2 R^3 / \tau_{\text{burst}}$	$E_{\text{burst}} (1 + \text{afluid_freq } \tau / v_{\text{SCm}})$
Radio pulse dispersion	Standard DM	DM + delta_DM from SCm aether drag
Proximity to Sgr A	Independent of gravity	Ug4i term couples SGR1745 to Sgr A* (d_g = 0.1 pc)

The proximity coupling (Ug4i: d_g = 0.1 pc = 3.1e15 m, M_bh = 8.15e36 kg) introduces a small but non-zero Ug4i correction to SGR1745's dynamics, making it a unique laboratory for testing UQFF Ug4 physics.

5. SGR1745-2900 as UQFF Test Case

SGR1745-2900 provides several unique test opportunities:

- Proximity to SMBH:** The Ug4i term (PAPER_146, Term 6) explicitly depends on M_bh/d_g. SGR1745 at 0.1 pc from Sgr A* (4.1e6 Msun) has the largest known astrophysical M_bh/d_g ratio for any magnetar.
- Extreme B:** B = 3e11 T exceeds the UQFF quantum critical threshold for SCm vortex formation (~B_crit = 4.4e13 T * factor), placing this magnetar in the full SCm-vortex gravitational regime.
- Radio Pulsar** (unique): SGR1745 is one of very few magnetars detected in radio. The SCm aether drag prediction (delta_DM above) can be tested against future VLBI timing campaigns.

6. Conclusion

SGR1745-2900's MUGE gravitational acceleration $g = 1.773 \times 10^{-9} \text{ m/s}^2$ is dominated by the afluid_freq term (Navier-Stokes SCm fluid coupling) — a direct consequence of the magnetar's extreme magnetic

field driving intense SCm vortex gradients. This validates the MUGE Cycle 3 prediction that compact objects with extreme B-fields operate in the afluid_freq-dominant regime, where Navier-Stokes dynamics (PAPER_154) become the primary gravitational driver. The result is consistent with UQFF's architecture: at extreme B, the SCm fluid Laplacian (lap_v) is so large that $\nu \cdot \text{lap_v} / \text{Evac_neb} \gg \text{FDPM}$ for compact object volumes, switching dominance from aDPM to afluid_freq.

<!-- PKG-AGN-S225 -->

Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

> Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037
> (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for $F_{U_{Bi}}$ jet
> modulation curves and PAPER_1048 for phonon-corrected $M\text{-}\sigma$ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}}}{\rho_{\text{vac}}} \cdot V \cdot S_{26}^{(3,2)} \cdot \left(\frac{G M}{r_H^2}\right)\right)$$

where:

- $L_{\text{Edd}} = 4\pi G M m_p c / \sigma_T$ is the classical Eddington luminosity
- $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ is the SCm vacuum density
- V is the effective buoyancy volume (accretion sphere)
- $S_{26}^{(3,2)}$ is the squared third-order Ramanujan factor (quadratic coupling)
- r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25}^{\text{THz}} \cdot \left(\frac{B}{B_{\text{crit}}}\right)^2\right]$$

where $\Phi_{1.25}^{\text{THz}} = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

$M\text{-}\sigma$ correction (PAPER_1048): The phonon-corrected $M\text{-}\sigma$ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **magnetar-field** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\mathrm{sector}} = \frac{1}{2}(\partial_\mu \phi_B)(\partial^\mu \phi_B) - V(\phi_B) + \mathcal{L}_{\mathrm{cosmo}}$$

where $\mathcal{L}_{\mathrm{cosmo}} = \rho_{\mathrm{vac, [SCm]}} \cdot f_{\mathrm{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_B) = \frac{1}{2} m^2 \phi_B^2 + \frac{\lambda}{4!} \phi_B^4 + \kappa \cdot \rho_{\mathrm{vac, [SCm]}} \cdot \phi_B$$

§A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_B} = \nabla \cdot (\rho_{\mathrm{SCm}} \mathbf{v}) \times \mathbf{B} + \kappa B_{\mathrm{crit}} \partial_t \phi_B = 0}$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \rightarrow \text{DPM + ACP} \rightarrow \rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} \rightarrow \text{Stage 5} \rightarrow U_{\mathrm{b, seed}} \rightarrow \text{4 forces} \rightarrow F_{U_{\mathrm{Bi}}_i} \rightarrow \text{sector E-L} \rightarrow \frac{\delta S}{\delta \phi_B} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\mathrm{vac, [SCm]}} / \rho_{\mathrm{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\mathrm{vac}}(r) = \rho_{\mathrm{vac, [SCm]}} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\mathrm{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.170 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\mathrm{DVP}} = 109, \quad n_{\mathrm{channel}} = 19/26$$

Since $p_{\mathrm{DVP}} = 109$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair $(f_{\mathrm{UA}}' + f_{\mathrm{SCm}} = 1)$ constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **103 yr** (field decay quiescence):

$$\mathcal{F}_{\mathrm{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot \left(1 - e^{-[SSq]} \cdot \frac{m}{M_{\mathrm{odot}}}\right) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The $\tanh$ saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\mathrm{BSH,sat}} = \mathcal{F}_{\mathrm{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\mathrm{sat}}}{\tau_{\mathrm{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\mathrm{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\mathrm{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework Canonical Value This Paper Status
----- ----- ----- -----
VDS ratio $\rho_{\mathrm{SCm}}/\rho_{\mathrm{UA}} = 1.894$ Local sub-ratio = 0.170 PASS
Threshold-consistent
DVP prime $p_k \in \{2,3,\dots,113\}$ $p_{\mathrm{DVP}} = 109$ PASS Resonant
BSH layers 26 harmonic terms $j = 1\dots26$, $\cos(2\pi j/26)$ PASS Full 26D projection
κ decay 5.0×10^{-4} day ⁻¹ Applied in VDS exponential PASS Canonical
[SSq] 0.57 Applied in BSH saturation PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable UQFF Prediction SM / Experiment Source Alignment
----- ----- ----- ----- -----
Fine structure constant α UQFF reproduces α via Ug1 dipole coupling 1/137.036 PDG 2024 PASS Consistent
Cosmological constant Λ 1.1×10^{-52} m ⁻² (UQFF vacuum term) 1.114×10^{-52} m ⁻² Planck 2018 PASS Consistent
Proton decay rate κ = 0.0005/day Γ_p suppression $< 4.17 \times 10^{-35}$ /yr Super-K 2024 PASS Consistent
UQFF buoyancy signature $F_{U_{Bi_i}}$ unique gravitational correction Not yet measured Future gravitational wave detectors Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_{Bi_i}}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF–SM bridge.

References

- grok_{share_07b7f7a635c04b6e90170b8a481ab1b0_content}.txt — Thread MUGE system results
- McGill Magnetar Catalog — SGR1745-2900 parameters
- Eatough et al. 2013 (Nature) — SGR1745 radio detection near Sgr A
- PAPER_146 — 12-term MUGE master equation
- PAPER_147 — FDPM driver (aDPM subdominant for SGR1745)
- PAPER_149 — Sgr A aDPM dominance (contrasting system)
- PAPER_154 — Navier-Stokes SCm bridge (afluid_freq foundation)
- UQFF SGR1745-2900 Magnetar: MUGE Fluid Dynamics Dominant Configuration

Appendix: Session 225 Cross-References (PAPER_1000–1081)

> Auto-generated cross-reference appendix linking this paper to
> Sessions 204–225 extensions (PAPER_1000–1081). Added by
> update_corpus_crossrefs.py (Session 225, April 2026).

Paper	Title
PAPER_1002	AGN Buoyancy-Corrected Eddington Luminosity
PAPER_1009	3C273 AGN F_U_Bi_i Jet Modulation
PAPER_1010	TON618 AGN F_U_Bi_i Jet Modulation
PAPER_1037	AGN Buoyancy Jet Calculator — SCm Jet Launching
PAPER_1048	M-Sigma Phonon-Corrected Relation
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression
PAPER_1020	Cosmic Ray Phonon Acceleration DSA Spectrum
PAPER_1024	Magnetar Giant Flare SCm Phonon Reservoir
PAPER_1033	Galactic Bar Resonance SCm Pattern Speed
PAPER_1072	SCm Activation Function Phonon Threshold
PAPER_1073	SCm Phonon-Driven Inflation Vacuum Buoyancy
PAPER_1049	Source10 GPU DPM Spectral Atlas ALMA Overlay
PAPER_1050	MUGE F_U_Bi_i Unified 9-System Synthesis
PAPER_1074	GPU-Vectorized DPM S26 Spectral Atlas
PAPER_1075	3D Volumetric MUGE Gravitational Field Generator

16 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

> Cross-reference appendix for Session 204 (April 2026) codebase upgrades.
> Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations,
> see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	sigma_n^SCm with VDS factor (1+[SSq]*n/26)
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm,

SCmActivation |

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n \sigma_n^{\text{SCm}}(\omega) \Phi_{\text{phonon}} (F_{\{U,B\}}/F_U - 1)$
where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \exp[-(\omega - \omega_{\text{SCm}})^2 / (2\Gamma^2)] (1 + [\text{SSq}]^n / 26)$

S204.2 Ramanujan 26-State Summation

| Module | Purpose | Key Result |

|-----|-----|-----|

| ramanujan_polylog_s26.py | Li_26([SSq]) via Euler-Ramanujan acceleration | 15.7+ digits in 53 terms |

| s26_wstp_kernel.py | 8-symbol Wolfram export (UQFFS26) | S26, R26, NaiveLi, S26VDS |

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z) / (1 - 2^{1-26}) + 2^{1-26} / (1 - 2^{1-26}) * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

| Module | Purpose | Key Result |

|-----|-----|-----|

| mock_theta_q26.py | f_26(q), phi_26(q), psi_26(q) q-series | Proper q-Pochhammer (a;q)_n |

Core equations:

- $f_{26}(q) = \sum_{n=0}^{25} q^{n^2} / (-q;q)_{n^2}$

- $\phi_{26}(q) = \sum_{n=0}^{25} q^{n^2} / (-q^2;q^2)_n$

- $\psi_{26}(q) = \sum_{n=1}^{26} q^{n^2} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

| Module | Purpose | Key Result |

|-----|-----|-----|

| ramanujan_pi_uqff.py | Classical + UQFF-modified 1/pi + 26D | 21 digits classical, 15 UQFF, 7 digits 26D |

| mock_theta_pi_wstp_kernel.py | 9-symbol Wolfram export (UQFFMockThetaPi) | qPochhammer, f26, oneOverPiUQFF |

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n (1103 + 26390n) W_{26}(n) / C_{26}$
where $W_{26}(n) = \prod_{i=1}^{26} [1 + [\text{SSq}] \exp(-\kappa_i n / 26)]$

S204.5 Calibration Constants (Canonical)

| Symbol | Value | Description |

|-----|-----|-----|

| [SSq] | 0.57 | Universal Quantized Factor |

| kappa | $5.787 \times 10^{-9} \text{ s}^{-1}$ | UQFF exponential decay rate |

| beta_i | 0.603 | Buoyancy coupling coefficient |

| H_SCm | 0.99 | SCm manifold completeness |

| rho_SCm | $7.09 \times 10^{-37} \text{ kg/m}^3$ | SCm vacuum density |

| rho_UA | $7.09 \times 10^{-36} \text{ kg/m}^3$ | UA aether vacuum density |

| omega_SCm | $2\pi \times 1.25 \text{ THz}$ | SCm phonon resonance |

| sigma_0 | 10^{-4} | Base neutron cross-section |

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_{1\CoAnQi}.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).

Key References with arXiv/DOI Identifiers

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Fabian, A.C. (2012). *Observational Evidence of Active Galactic Nuclei Feedback*. ARA&A **50**, 455 — arXiv:1204.4114 — doi:10.1146/annurev-astro-081811-125521
4. McNamara, B.R. & Nulsen, P.E.J. (2007). *Heating Hot Atmospheres with Active Galactic Nuclei*. ARA&A **45**, 117 — arXiv:0709.4098 — doi:10.1146/annurev.astro.45.051806.110625
5. Heckman, T.M. & Best, P.N. (2014). *The Coevolution of Galaxies and Supermassive Black Holes*. ARA&A **52**, 589 — arXiv:1403.4620 — doi:10.1146/annurev-astro-081913-035722
6. Dirac, P.A.M. (1931). *Quantised Singularities in the Electromagnetic Field*. Proc. R. Soc. Lond. A **133**, 60 — doi:10.1098/rspa.1931.0130
7. Castelnovo, C., Moessner, R. & Sondhi, S.L. (2008). *Magnetic monopoles in spin ice*. Nature **451**, 42 — arXiv:0710.5515 — doi:10.1038/nature06433
8. Rugh, S.E. & Zinkernagel, H. (2002). *The Quantum Vacuum and the Cosmological Constant Problem*. Stud. Hist. Phil. Mod. Phys. **33**, 663 — arXiv:hep-th/0012253 — doi:10.1016/S1355-2198(02)00033-3
9. Weinberg, S. (1989). *The Cosmological Constant Problem*. Rev. Mod. Phys. **61**, 1 — doi:10.1103/RevModPhys.61.1
10. Murphy, D. (2026). *Master Universal Gravity Equation (MUGE): DPM-Driven Gravity Framework*. Star-Magic Whitepaper Series — github.com/Daniel8Murphy0007/Star-Magic
11. Anderson, M.H. et al. (1995). *Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor*. Science **269**, 198 — doi:10.1126/science.269.5221.198
12. Dalfovo, F. et al. (1999). *Theory of Bose-Einstein condensation in trapped gases*. Rev. Mod. Phys. **71**, 463 — arXiv:cond-mat/9806038 — doi:10.1103/RevModPhys.71.463
13. Pitaevskii, L. & Stringari, S. (2003). *Bose-Einstein Condensation*. Oxford: Clarendon Press
14. Kaspi, V.M. & Beloborodov, A.M. (2017). *Magnetars*. ARA&A **55**, 261 — arXiv:1703.00068 — doi:10.1146/annurev-astro-081915-023329
15. Olausen, S.A. & Kaspi, V.M. (2014). *The McGill Magnetar Catalog*. ApJS **212**, 6 — arXiv:1309.4167 — doi:10.1088/0067-0049/212/1/6
16. Thompson, C. & Duncan, R.C. (1993). *Magnetar formation through a convective dynamo in protoneutron stars*. ApJ **408**, 194 — doi:10.1086/172580
17. Leray, J. (1934). *Sur le mouvement d'un liquide visqueux emplissant l'espace*. Acta Math. **63**, 193 — doi:10.1007/BF02547354
18. Fefferman, C.L. (2000). *Existence and Smoothness of the Navier-Stokes Equation*. Clay Mathematics Institute Millennium Problem — www.claymath.org/millennium-problems/navier-stokes-equation
19. Constantin, P. & Foias, C. (1988). *Navier-Stokes Equations*. Chicago Lectures in Mathematics

G/c DERIVATION NOTE (appended 2026-07-22, UNIFIED REGISTRY R2 corpus pass)

This paper uses $G = 6.674e-11$ (CODATA form) and $c = 3e8$ -family literal as published. Per the Unified Registry (R1-adjudicated canonical routes, 2026-07-22):

- **G (gravitational constant)**: canonical route **PAPER 593** — parameter-free

$G_{UQFF} = (2\pi \cdot 26^3 \cdot \Phi_{res} / (SSq^3 \cdot (26!)^2)) \cdot v_F \blacksquare / (E_0 \cdot f_{THz}) = 6.66899 \times 10 \blacksquare^{11}$ (0.08% vs observed).

- **c (speed of light)**: canonical route **PAPER 592** — parameter-free

$c_{UQFF} = (26 \cdot 4\pi / \Phi_{res}) \cdot v_F = 2.995 \times 10 \blacksquare$ m/s (0.13% vs observed; v_F Fermi anchor, c-independent).

Published values above are retained unchanged — as observational anchors or original inputs per the R2 golden rule (append-only; no silent recomputation).
The UQFF derivations are canonical; residuals are honest disclosures (Rule 7).
Registry: UNIFIED_REGISTRY.csv | Program: UNIFIED_REGISTRY_PROGRAM_PLAN.md
