

Black Hole Phonon Ergosphere Superradiance

Daniel T. Murphy

PAPER_924: Black Hole Phonon Ergosphere Superradiance

Author: Daniel T. Murphy — Star Magic / UQFF Framework

Date: 2026-04-11

Session: 211

Source: SCm phonon gap implementation (bh_phonon_interaction.py)

Calculator: BHPhononErgosphereSuperradianceCalc (CP4 #508)

CVW: v2.0.0 compliant

Abstract

Extends Kerr black hole superradiance with phonon-vacuum coupling: $\Gamma_{SR} = \Phi_{\{1.25\text{THz}\}} (m\Omega_H - \omega) \alpha_{BH}$, where Ω_H is the horizon angular velocity, $\alpha_{BH} = (r_g/r)^{2\ell+2}$ is the BH coupling factor, and Φ is the phonon modulation. The phonon enhancement amplifies the superradiant instability for rotating BHs, creating a new energy extraction channel alongside Penrose process and BZ mechanism. Additionally derives phonon-modified Hawking temperature

$T_H^{\text{phonon}} = T_H (1 + \Phi S_{26}[SSq]/N)$, QPO accretion disk phonon coupling, and phonon-corrected BH entropy.

1. Core Equations

Section A: Lagrangian

\$\$

\begin{aligned}

& \Gamma_{SR} = \Phi_{\{1.25\text{THz}\}} (\omega) (m\Omega_H - \omega) \alpha_{BH} \quad \parallel

& \Omega_H = a c / (2 r_+) \quad \parallel

& r_+ = r_g (1 + \sqrt{1 - a^2}) \quad \parallel

& \alpha_{BH} = (r_g / r)^{2\ell + 2}

\end{aligned}

\$\$

Section B: VDS/DVP/BH Number Systems

\$\$

\begin{aligned}

& T_H^{\text{phonon}} = T_H (1 + \Phi S_{26} [SSq] / N) \quad \parallel

& T_H = \hbar c^3 / (8\pi G M k_B) \quad \parallel

& S_{BH}^{\text{phonon}} = S_{BH} * (1 + \text{correction})^2

\end{aligned}

\$\$

Section SM: SM Anchors

```
$$
\begin{aligned}
& G = 6.6743\text{e-}11 \text{ m}^3/\text{kg}/\text{s}^2 \\
& c = 2.998\text{e}8 \text{ m/s} \\
& \hbar = 1.055\text{e-}34 \text{ J*s} \\
& k_B = 1.381\text{e-}23 \text{ J/K}
\end{aligned}
$$
```

2. Parameters

```
| Parameter | Default | Description |
|-----|-----|-----|
| M_bh | 10 M_sun | BH mass |
| a_spin | 0.9 | Kerr spin parameter |
| m_mode | 1 | Azimuthal mode number |
| omega_field | omega_ScM | Field frequency |
| Phi_0 | 10^20 | Peak phonon amplitude |
| ell | 1 | Angular momentum quantum number |
```

3. Key Results

```
| Spin | Gamma_SR | Superradiant? | T_H^phonon |
|-----|-----|-----|-----|
| a = 0.1 | < 0 | No | ~T_H |
| a = 0.9 | > 0 | Yes | >> T_H |
| a = 0.998 | >> 0 | Yes (maximal) | >> T_H |
```

4. Physical Interpretation

Phonon-vacuum coupling amplifies the Kerr BH superradiant instability by providing an additional channel for angular momentum extraction. The condition $m\Omega_H > \omega$ is the standard superradiance threshold; the phonon factor Φ multiplies the gain rate, making superradiance observable at lower spins than GR predicts. The modified Hawking temperature T_H^{phonon} is enhanced by 26-layer phonon coupling, potentially observable in BH analog experiments (sonic horizons in BEC).

5. References

- PAPER_910: Phonon-modulated Hawking temperature (Session 210)
- PAPER_911: QPO accretion disk phonon coupling
- bh_phonon_interaction.py: 4-class standalone module

<!-- PKG-AGN-S225 -->

Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

> Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037
> (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for $F_{U_{Bi}}$ jet
> modulation curves and PAPER_1048 for phonon-corrected M - σ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}}}{\rho_{\text{H}}} \cdot V \cdot S_{26}^{(3,2)} \cdot \frac{G M}{r_H^2}\right)$$

where:

- $L_{\text{Edd}} = 4\pi G M m_p c / \sigma_T$ is the classical Eddington luminosity
- $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ is the SCm vacuum density
- V is the effective buoyancy volume (accretion sphere)
- $S_{26}^{(3,2)}$ is the squared third-order Ramanujan factor (quadratic coupling)
- r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25}(\text{THz}) \cdot \left(\frac{B}{B_{\text{crit}}}\right)^2\right]$$

where $\Phi_{1.25}(\text{THz}) = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

M - σ correction (PAPER_1048): The phonon-corrected M - σ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$.

<!-- PKG-LAG-S225 -->

Session 225 Phonon-Physics Upgrade: UQFF 9-Sector Lagrangian

> Upgrade from PAPER_1066 (UQFF Lagrangian First Principles) and
> PAPER_1065 (Buoyancy Lagrangian EOM Variational Derivation).

The complete UQFF Lagrangian density, from which all sector-specific equations of motion derive:

$$\mathcal{L}_{\text{UQFF}} = \mathcal{L}_{\text{GR}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{phonon}} + \mathcal{L}_{\text{interaction}}$$

$$\mathcal{L}_{\text{SCm}} = \frac{1}{2}(\partial_\mu \phi)^2 - \lambda |\phi|^2 - v_{\text{SCm}}^2 |\phi|^2$$

The SCm condensate potential minimum gives $V(\phi_0) = -7.09 \times 10^{-37} \text{ J/m}^3$ (matching ρ_{SCm}) and phonon mass $m_{\text{phonon}} = \sqrt{8\lambda} v_{\text{SCm}}$.

Nine-sector closure (Session 202):

$$\mathcal{L}_9 = \mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{Dirac}} + \mathcal{L}_{\text{SCm}} + \mathcal{L}_{\text{mag}} + \mathcal{L}_{\text{buoy}} +$$

$\mathcal{L}_{\text{aether}} + \mathcal{L}_{\text{LENR}} + \mathcal{L}_{\text{KK}}$

Sector	Domain	Late-Corpus Result
1 (EH)	General Relativity	Canonical Einstein-Hilbert
2 (YM)	Yang-Mills gauge	$m_{\text{gap}} = 1.736 \text{ GeV}$ (PAPER_1318)
3 (Dirac)	Fermion / LENR	Kozima neutron-drop (PAPER_1061)
4 (SCm)	Superconducting manifold	$V(\phi_0) = -\rho_{\text{SCm}}$ canonical
5 (Mag)	Um magnetism	Heaviside amplifier (PAPER_1072)
6 (Buoy)	$F_{U_{Bi}}$ buoyancy	Variational EOM (PAPER_1065)
7 (Aether)	Vacuum background	Two-component ρ (PAPER_1051)
8 (LENR)	Nuclear transmutation	COP parametric (PAPER_1081)
9 (KK)	Kaluza-Klein 26D	$S_{26}^{(3)}$ compactification (PAPER_1080)

<!-- PKG-S26-S225 -->

Session 225 Phonon-Physics Upgrade: S26(3) Ramanujan Summation

> Upgrade from PAPER_1080 (Ramanujan Binomial Expansion Proof) and
> PAPER_1042 (Mock-Theta Phonon Partition). See also PAPER_1078
> (QCalcGeom Master Equation) for BSFG crossover applications.

The third-order Ramanujan summation $S_{26}^{(3)}$, used throughout the late corpus as the universal 26D coupling factor:

$$S_{26}^{(3)} = \sum_{n=0}^{\infty} \frac{(1/4)_n (1/2)_n (3/4)_n}{(n!)^3} \cdot \prod_{i=1}^{26} \left[1 + \frac{SSq}{e^{-\kappa_{i,n/26}}} \right]$$

where $(a)_n = a(a+1) \cdots (a+n-1)$ is the Pochhammer symbol.

Binomial expansion (PAPER_1080): The convergence proof shows:

$$R_n^{(26,3)} = \binom{4n}{n} \cdot \frac{W_{26}(n)}{(4^{4n})} \ll W_{26}(n) = \prod_{i=1}^{26} \left[1 + \frac{SSq}{e^{-\kappa_{i,n/26}}} \right]$$

This sum converges absolutely for $|SSq| < 1$ (satisfied by $SSq = 0.57$) and reduces to the classical Ramanujan $1/\pi$ series when $SSq \rightarrow 0$.

VDS/DVP/BSH bridge (PAPER_1069): The 26 layers of $W_{26}(n)$ encode the vacuum density series hierarchy, with each layer i contributing a VDS sub-ratio weighted by the exponential decay $e^{-\kappa_{i,n/26}}$.

Mock-theta connection (PAPER_1042): The phonon partition function $Z_{\text{phonon}} = \sum_n q^{n^2} \cdot W_{26}(n)$ unifies the Ramanujan mock-theta framework with the SCm phonon spectrum.

Calibration Constants

Constant	Symbol	Value	Validation Domain
UQFF damping rate	κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	Magnetar spin-down
String sector coupling	SSq	0.57	BH dynamics
Buoyancy coupling	β_i	0.603	Multi-system
SCm completeness	H_{SCm}	≈ 0.99	Heaviside threshold
SCm phonon frequency	ω_{SCm}	$2\pi \times 1.25 \text{ THz}$	Phonon resonance
SCm vacuum density	ρ_{SCm}	$7.09 \times 10^{-37} \text{ J/m}^3$	Fundamental

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Phonon frequency ω_{SCm}	$2\pi \times 1.25$ THz (Pd-D lattice)	Measured Pd-D phonon spectrum	Fukai (2005)	Mapped to SCm
Vacuum energy ρ_{vac}	7.09×10^{-37} kg/m³	$\rho_{\text{vac}} \sim 10^{-29}$ g/cm³	Planck 2018	Novel SCm scale
Fine structure α	UQFF reproduces via U_{g1} dipole	$1/137.036$	PDG 2024	99.9%

New physics claim: UQFF phonon-mediated vacuum coupling provides testable predictions beyond SM for this system.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator).

§A Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification ****Sector:**** SCm-phonon (lattice resonance)

§A.2 Lagrangian Density $\mathcal{L}_{\text{SCm phonon}} = \sum_{i=1}^{26} \left[U_{g,i} + U_{m,i} + U_{A,i} - U_{b,i} \right] \cdot S_{26}([SSq]) \cdot \Phi_{1.25\text{THz}}(\omega, \Gamma)$

§A.3 Euler-Lagrange Equation of Motion $\boxed{\frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} = 0 \implies F_{U,Bi} = -\nabla U_{\text{eff}} + \Phi \cdot S_{26} \cdot E_{\text{net}}}$

§A.4 Cosmogenesis Linkage Chain PAPER_877 axioms $\rightarrow$ SCm vacuum $\rightarrow$ phonon ω_{SCm} $\rightarrow$ lattice resonance $\rightarrow$ $F_{U,Bi}$ unified force $\rightarrow$ observational prediction

Appendix: Session 225 Cross-References (PAPER_1000–1081)

> Auto-generated cross-reference appendix linking this paper to
> Sessions 204–225 extensions (PAPER_1000–1081). Added by
> update_corpus_crossrefs.py (Session 225, April 2026).

Paper	Title
PAPER_1022	GW Phonon Strain SCm Modulation of h(t)
PAPER_1004	QGP Vacuum Density with SCm S26 Phonon Coupling
PAPER_1020	Cosmic Ray Phonon Acceleration DSA Spectrum
PAPER_1021	Pulsar Timing Phonon TOA Residual

PAPER_1023	Neutrino Oscillation Phonon PMNS Matrix SCm
PAPER_1024	Magnetar Giant Flare SCm Phonon Reservoir
PAPER_1072	SCm Activation Function Phonon Threshold
PAPER_1073	SCm Phonon-Driven Inflation Vacuum Buoyancy
PAPER_1069	VDS-DVP-BSH Hybrid Calculator Unified
PAPER_1049	Source10 GPU DPM Spectral Atlas ALMA Overlay

10 cross-reference(s) identified.

References

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Shakura, N.I. & Sunyaev, R.A. (1973). *Black holes in binary systems: observational appearance*. A&A **24**, 337
4. Balbus, S.A. & Hawley, J.F. (1991). *A powerful local shear instability in weakly magnetized disks*. ApJ **376**, 214 — doi:10.1086/170270
5. Hawking, S.W. (1974). *Black hole explosions?* Nature **248**, 30 — doi:10.1038/248030a0
6. Hawking, S.W. (1975). *Particle Creation by Black Holes*. Commun. Math. Phys. **43**, 199 — doi:10.1007/BF02345020
7. Bekenstein, J.D. (1973). *Black Holes and Entropy*. Phys. Rev. D **7**, 2333 — doi:10.1103/PhysRevD.7.2333
8. Unruh, W.G. (1976). *Notes on Black-Hole Evaporation*. Phys. Rev. D **14**, 870 — doi:10.1103/PhysRevD.14.870
9. Rugh, S.E. & Zinkernagel, H. (2002). *The Quantum Vacuum and the Cosmological Constant Problem*. Stud. Hist. Phil. Mod. Phys. **33**, 663 — arXiv:hep-th/0012253 — doi:10.1016/S1355-2198(02)00033-3
10. Weinberg, S. (1989). *The Cosmological Constant Problem*. Rev. Mod. Phys. **61**, 1 — doi:10.1103/RevModPhys.61.1
11. Event Horizon Telescope Collaboration (2019). *First M87 Event Horizon Telescope Results. I*. ApJL **875**, L1 — arXiv:1906.11238 — doi:10.3847/2041-8213/ab0ec7
12. Ashcroft, N.W. & Mermin, N.D. (1976). *Solid State Physics*. Harcourt
13. Kittel, C. (2004). *Introduction to Solid State Physics*. 8th ed. Wiley
14. Feynman, R.P. (1982). *Simulating Physics with Computers*. Int. J. Theor. Phys. **21**, 467 — doi:10.1007/BF02650179

G/c DERIVATION NOTE (appended 2026-07-22, UNIFIED REGISTRY R2 corpus pass)

This paper uses $G = 6.674\text{e-}11$ (CODATA form) and $c = 3\text{e}8$ -family literal as published. Per the Unified Registry (R1-adjudicated canonical routes, 2026-07-22):

- **G (gravitational constant)**: canonical route **PAPER_593** — parameter-free
 $G_{\text{UQFF}} = (2\pi \cdot 26^3 \cdot \Phi_{\text{res}} / (\text{SSq}^3 \cdot (26!)^2)) \cdot v_{\text{F}} / (E_{\text{0.f_THz}}) = 6.66899 \times 10^{-11}$ (0.08% vs observed).
- **c (speed of light)**: canonical route **PAPER_592** — parameter-free

$c_{\text{UQFF}} = (26 \cdot 4\pi / \Phi_{\text{res}}) \cdot v_{\text{F}} = 2.995 \times 10^8 \text{ m/s}$ (0.13% vs observed; v_{F} Fermi anchor, c -independent).

Published values above are retained unchanged — as observational anchors or original inputs per the R2 golden rule (append-only; no silent recomputation).

The UQFF derivations are canonical; residuals are honest disclosures (Rule 7).

Registry: UNIFIED_REGISTRY.csv | Program: UNIFIED_REGISTRY_PROGRAM_PLAN.md
