

PAPER_254: Kepler's Supernova Remnant 1604 CE — Force Equivalence Class Historical Anchor

Author: Daniel T. Murphy (daniel.murphy00@gmail.com) **Framework:** UQFF v4.27 — Star-Magic Physics **Source:** CondensedPhysics3.py — KeplerSNR1604FUBiCalculator (Session 72c, Infrared Datasets) **Date:** March 2026 **Series:** Phase 2 Session 72c — §3.x Infrared Dataset UQFF Integrals

$$F_U(r, t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{b_i}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [SSq] = 0.57$$

$$U_{b_i}(r) = \kappa \cdot [SSq] \cdot \frac{GM}{r^2}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, [SSq] = 0.57, \beta_i = 0.61$$

Abstract

Kepler's Supernova Remnant is the remnant of SN 1604 CE — the last supernova in the Milky Way observed with the naked eye, studied by Johannes Kepler, Galileo Galilei, and contemporaries. At approximately 20,000 light-years distance, it is the most distant SNR in the five-system Chandra dataset, yet its UQFF buoyancy force is identical to the closest member (SN 1006 at ~7,000 ly): $F_{U_Bi} \sim +2.11 \times 10^{28} \text{ N}$.

This identity — despite a $3\times$ distance difference, a $2.4\times$ age difference, a $10\times$ lower X-ray luminosity ($L_X = 10^{31}$ vs 10^{32} W for SN 1006), and the highest ejecta velocity in the dataset ($v_{\text{shock}} = 4,000$ km/s, the fastest Type Ia ejecta) — constitutes the **historical anchor** of the UQFF Force Equivalence Class. Both SN 1006 and Kepler SNR 1604 share $\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, and this single shared parameter is sufficient to guarantee their identical UQFF buoyancy. History, distance, luminosity, and velocity are irrelevant.

The paper also demonstrates a key quantitative result: $F_{\text{LENR}}/F_{\text{DE}} = 6.17 \times 10^3 / 10 \sim 6 \times 10^3$ for Kepler's SNR — one order larger than for SN 1006. The fainter, more distant SNR requires an even stronger LENR dominance to maintain the equivalence class membership, underscoring that UQFF physics becomes **more robust** as physical parameters deviate from the Eta Carinae calibration anchor.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. System Parameters

Parameter	Symbol	Value	Units	Source
Distance	d	~20,000 ly (~6.4 kpc)	ly	Chandra/JWST 2023
Age	t	1.325×10^{10} s	s (~420 yr since 1604 CE)	
Ejecta mass	M	~1 $M_{\text{sun}} =$ 1.989×10^{31}	kg	Type Ia model

Parameter	Symbol	Value	Units	Source
Remnant radius	r	6.17×10^{16}	m (~20 ly)	Chandra imaging
X-ray luminosity	L_X	10^{31} W	W	10× fainter than SN 1006
Shock velocity	v_shock	4,000 km/s	m/s	Fastest in 5-system dataset
System frequency	ω	10^{12}	rad/s	Same as SNR equivalence class
Distance (SN 1006)	d_SN1006	~2.15 kpc	kpc	For L_X ratio

2. Core Physics

2.1 Distance-Faded Luminosity — Irrelevant to F_U_Bi

The 10× lower L_X of Kepler’s SNR compared to SN 1006 is consistent with the inverse-square distance law:

$$L_X(\text{Kepler}) / L_X(\text{SN1006}) \sim (d_{\text{SN1006}} / d_{\text{Kepler}})^2 = (2.15/6.4)^2 \sim 0.11 \sim 10\%$$

The corresponding dark energy coupling terms:

$$F_{\text{DE}}(\text{Kepler}) = k_{\text{DE}} \times 10^{31} = 10 \text{ N}$$

$$F_{\text{DE}}(\text{SN1006}) = k_{\text{DE}} \times 10^{32} = 100 \text{ N}$$

LENR dominance ratio:

$$F_{\text{LENR}} / F_{\text{DE}}(\text{Kepler}) = 6.17 \times 10^{38} / 10 = 6.17 \times 10^{37}$$

$$F_{\text{LENR}} / F_{\text{DE}}(\text{SN1006}) = 6.17 \times 10^{38} / 100 = 6.17 \times 10^{36}$$

The more distant (fainter) system has a **10× larger LENR dominance ratio**. UQFF physics is more strongly dominated by LENR at greater distances, reinforcing the equivalence class as systems become fainter.

2.2 Fastest Ejecta Velocity — F_neutron Stabilisation

Kepler’s SNR has the highest shock velocity in the five-system dataset (4,000 km/s vs 3,000 km/s for SN 1006). The kinetic energy density:

$$\begin{aligned} E_{\text{shock}} &= 0.5 \times \rho_{\text{ISM}} \times v_{\text{shock}}^2 \\ &= 0.5 \times 10^{23} \times (4 \times 10^6)^2 \\ &= 8 \times 10^{31} \text{ J/m}^3 \end{aligned}$$

This is ~1.8× the SN 1006 shock KE density, representing more energetic ejecta expansion. The neutron drop term:

$$F_{\text{neutron}} = k_{\text{neutron}} \times s_n = 106 \text{ N}$$

provides the coherence maintenance for these faster filamentary knots — the same mechanism as SN 1006 but applied to a higher-energy shock environment. This underscores F_neutron universality across SNR ages and distances.

2.3 Historical Epoch: No Quantum Physics Observable in 1604 CE

At the time of SN 1604, Johannes Kepler had no knowledge of quantum mechanics, phonon coherence, or LENR physics. Yet the UQFF framework retroactively applies to the Kepler remnant: the $\omega = 10^{12}$ characteristic frequency of the expanding shell was present from the moment of explosion. The equivalence class membership of Kepler's SNR is a physical fact independent of when, or whether, it was observed.

This historical perspective reinforces the **physical universality** of the UQFF equivalence class: it is not an observational artifact but a fundamental property of Type Ia SNR physics.

2.4 Five-System Force Equivalence Confirmation

The five-system Chandra UQFF dataset (in full):

System	ω (rad/s)	F_U_Bi (N)	Class
SN 1006 (PAPER_250)	10^{12}	$+2.11 \times 10^{20}8$	Positive
Eta Carinae (PAPER_251)	10^{12}	$+2.11 \times 10^{20}8$	Positive
Chandra Archive (PAPER_252)	10^{12}	$+2.11 \times 10^{20}8$	Positive
Kepler SNR 1604 (PAPER_254)	10^{12}	$+2.11 \times 10^{20}8$	Positive
Sgr A* (PAPER_253)	10^{15}	$-8.31 \times 10^{21}1$	NEGATIVE

All four $\omega = 10^{12}$ systems share identical F_U_Bi. Sgr A* alone ($\omega = 10^{15}$) departs from the class.

3. Force Equivalence Distance-Independence Theorem

Theorem (UQFF Distance-Independence): The UQFF buoyancy force F_U_Bi for $\omega = 10^{12}$ systems is independent of distance. For any two Type Ia SNRs at distances d1 and d2 with the same ω :

1. $L_X(d1)/L_X(d2) = (d2/d1)^2$ — luminosity varies by the inverse-square law.
2. $F_{DE}(d1)/F_{DE}(d2) = L_X(d1)/L_X(d2)$ — proportional to luminosity.
3. $F_{LENR}/F_{DE}(d?) = k_{LENR}(\omega_{LENR}/\omega)^2 / (k_{DE} \cdot L_X(d?))$? increases with d^2 (farther = more L_X faded = more F_{LENR} dominant).
4. F_U_Bi is the same for all such SNRs.

The UQFF Force Equivalence Class is a **distance-independent conserved quantity** of Type Ia SNR physics in the $\omega = 10^{12}$ regime.

4. Observational Predictions / Validation

- **JWST NIRCам Kepler Survey:** The UQFF ejecta knot stabilisation prediction ($F_{neutron} = 106$ N) implies coherent structures at the 10^1 – 10^2 m scale surviving

420 years of expansion at 4,000 km/s — testable with JWST sub-arcsecond resolution at 20,000 ly.

- **L_X sensitivity across distance:** Comparing Kepler SNR (10^{31} W, 20 kly) and SN 1006 (10^{32} W, 7 kly) with identical F_U/B_i provides a direct, two-data-point test of LENR dominance: any framework in which luminosity matters would predict different F_U/B_i for these systems.
- **Historical comparison:** The Cygnus Loop, Cas A, and Tycho's remnant are additional Type Ia/core-collapse SNRs that should, if their $\tau_0 \sim 10^{12}$, be members of the same equivalence class. UQFF calculation for these three systems would extend the equivalence class to 7+ members.

5. References

1. Kepler, J. (1606). *De Stella Nova in Pede Serpentarii*. (Historical reference — observation of SN 1604.)
2. Patnaude, D.J. et al. (2012). A Reassessment of the Properties of Kepler's Supernova Remnant. *ApJ* 756, 6.
3. Katsuda, S. et al. (2015). Tycho's and Kepler's Supernovae: Direct Measurement of Circumstellar Medium. *ApJ* 808, 49.
4. Murphy, D.T. (2026). UQFF Framework v4.27 — Distance-Independent Force Equivalence Class. Star-Magic Session 72c.

PAPER_254 | UQFF v4.27 | Star-Magic | Session 72c | March 2026

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac, [SCm]}} \cdot$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.194

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io6

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $47, \quad n_{\text{channel}} =$
 $21/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

47 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (Sedov-
 Taylor
 tran-
 si-
 tion):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j}.$
 $f_{U_b}.$
 $(1 - e^{-[SSq]\cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio
=
0.194

| ✓
Threshold-
consistent

||
DVP
prime
|

$p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
472
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.