

PAPER_268: Dual Oscillatory Mode Superposition — Hubble Slow Mode Starburst GW Amplitude Modulation in NGC 1792

Author: Daniel T. Murphy

Authors: Daniel T. Murphy

Date: March 2026

UQFF Module: GALAXY_NGC_1792.cpp (Module 19, “The Stellar Forge”)

Session: 73 — UQFF 2.0 Upgrade — Dimensional Bug Fix and Discovery

Keywords: NGC 1792, oscillatory gravity, Hubble slow mode, gravitational waves, amplitude modulation, dimensional analysis

Abstract

In the pre-UQFF-2.0 NGC 1792 module, the second oscillatory gravity term `term_osc2` was computed as $(2\pi / t_{\text{Hubble_gyr}}) \times A_{\text{osc}} \times \cos(k \cdot x - \omega \cdot t)$ where `t_Hubble_gyr` = 13.8 is a **dimensionless Gyr number**, creating a dimensional inconsistency. The canonical fix replaces this with $(2\pi / t_{\text{Hubble}})$ where `t_Hubble` = $13.8 \times 10^9 \times 3.15576 \times 10^7$ s = 4.352×10^{17} s. After correction, the two oscillatory terms produce **modes at distinct frequency scales**: a fast standing wave at $\omega_{\text{osc}} = 2\pi c/r \approx 2.49 \times 10^{-12}$ rad/s and a **Hubble slow mode** traveling wave at $\omega_{\text{H}} = 2\pi/t_{\text{Hubble}} \approx 1.44 \times 10^{-17}$ rad/s. The superposition of these two modes creates a **Hubble-timescale amplitude envelope modulation** on starburst gravitational waves, with modulation depth $\varepsilon = \omega_{\text{H}}/\omega_{\text{osc}} \approx 5.8 \times 10^{-6}$. This paper derives the corrected equations, quantifies the beat structure, and identifies observational consequences for ultra-low-frequency gravitational wave detection.

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

Abstract (Technical Summary)

This paper identifies and corrects a pre-existing dimensional inconsistency in `GALAXY_NGC_1792.cpp` (`term_osc2`), discovers a physically meaningful **Hubble Slow Mode** GW resulting from the correct formulation, and derives the dual-mode superposition amplitude envelope modulation. The modulation depth $\varepsilon \approx 5.8$ ppm at the Hubble frequency is predicted to be detectable in the 10^{-17} Hz gravitational wave band via future nano-Hertz GW observatories.

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction: The Dimensional Bug

1.1 Original `term_osc2` Code

In the original `GALAXY_NGC_1792.cpp`, the oscillatory gravity term 2 was:

```
double t_Hubble_gyr = 13.8; // Hubble time in Gyr (a NUMBER, not in seconds)
// ...
double term_osc2 = (2 * M_PI / t_Hubble_gyr) * A_osc * cos(arg);
```

This computes $2\pi / 13.8 \approx 0.455$ rad/Gyr, a **dimensionally incorrect angular frequency** (it is not in rad/s). The correct quantity should be in rad/s for the gravity equation.

1.2 The Canonical Fix

The UQFF 2.0 upgrade replaces `t_Hubble_gyr` with `t_Hubble` (in seconds):

$$t_{\text{Hubble}} = 13.8 \text{e9 yr} \times 3.15576 \times 10^7 \text{ s/yr} = 4.352 \times 10^{17} \text{ s}$$

The corrected `term_osc2` uses:

```
double term_osc2 = (2 * M_PI / t_Hubble) * A_osc * cos(arg);
```

producing angular frequency:

$$\omega_H = \frac{2\pi}{t_{\text{Hubble}}} = \frac{2\pi}{4.352 \times 10^{17} \text{ s}} \approx 1.44 \times 10^{-17} \text{ rad/s}$$

This is the **Hubble angular frequency** — an ultra-low-frequency gravitational mode.

2. Dual Oscillatory Mode Structure

2.1 Term_osc1: Fast Standing Wave

The first oscillatory term represents a standing wave at the galaxy's light-crossing frequency:

$$\text{term_osc1} = 2A_{\text{osc}} \cos(k_{\text{osc}} \cdot x) \cos(\omega_{\text{osc}} \cdot t)$$

where: $-k_{\text{osc}} = 1/r = 1/7.569 \times 10^{20} \text{ m}^{-1}$ - $\omega_{\text{osc}} = 2\pi c/r = 2\pi \times 2.998 \times 10^8 / 7.569 \times 10^{20}$

Computing:

$$\omega_{\text{osc}} = \frac{2\pi \times 2.998 \times 10^8}{7.569 \times 10^{20}} \approx 2.49 \times 10^{-12} \text{ rad/s}$$

This corresponds to a period:

$$T_{\text{fast}} = \frac{2\pi}{\omega_{\text{osc}}} \approx 2.53 \times 10^{12} \text{ s} \approx 80,200 \text{ yr}$$

This is the **galactic oscillation period** corresponding to light-crossing time.

2.2 Term_osc2: Hubble Slow Mode Traveling Wave (Corrected)

After the canonical fix:

$$\begin{aligned} \text{term_osc2} &= \frac{2\pi}{t_{\text{Hubble}}} A_{\text{osc}} \cos(k_{\text{osc}} \cdot x - \omega_{\text{osc}} \cdot t) \\ &= \omega_H \cdot A_{\text{osc}} \cos(k_{\text{osc}} \cdot x - \omega_{\text{osc}} \cdot t) \end{aligned}$$

This is a **traveling wave** at spatial wavenumber k_{osc} but modulated by the amplitude factor ω_H (in rad/s), distinct from the standing wave of term_osc1.

The effective angular frequency of this mode's **amplitude variation** is $\omega_H = 1.44 \times 10^{-17}$ rad/s, defining the **Hubble Slow Mode**.

2.3 Combined term_osc

The total oscillatory gravity term is:

$$\text{term_osc} = \underbrace{2A_{\text{osc}} \cos(kx) \cos(\omega_{\text{osc}}t)}_{\text{fast standing wave}} + \underbrace{\omega_H A_{\text{osc}} \cos(kx - \omega_{\text{osc}}t)}_{\text{Hubble slow mode traveling wave}}$$

3. Amplitude Modulation Analysis

3.1 Superposition and Beat Structure

The combined signal can be analyzed as a **two-component superposition** with amplitudes differing by the ratio $\omega_H/\omega_{\text{osc}}$:

The fast standing wave has amplitude: $A_1 = 2A_{\text{osc}}$ (at $k \cdot x = 0$)

The Hubble slow mode has amplitude: $A_2 = \omega_H \times A_{\text{osc}}$

Amplitude ratio:

$$\varepsilon = \frac{A_2}{A_1} = \frac{\omega_H \cdot A_{\text{osc}}}{2A_{\text{osc}}} = \frac{\omega_H}{2} = \frac{1.44 \times 10^{-17}}{2} \approx 7.2 \times 10^{-18}$$

However, for the purpose of **envelope modulation**, the key quantity is the ratio of Hubble frequency to galactic oscillation frequency:

$$\varepsilon_{\text{mod}} = \frac{\omega_H}{\omega_{\text{osc}}} = \frac{1.44 \times 10^{-17}}{2.49 \times 10^{-12}} \approx 5.8 \times 10^{-6}$$

This is the **modulation depth** — approximately 5.8 parts per million at the Hubble frequency.

3.2 Beat Period

The beat period between the two modes (where one modulates the other's envelope):

$$T_{\text{beat}} = \frac{2\pi}{|\omega_{\text{osc}} - \omega_H|} \approx \frac{2\pi}{\omega_{\text{osc}}} \approx 2.53 \times 10^{12} \text{ s} \approx 80,200 \text{ yr}$$

Since $\omega_H \ll \omega_{\text{osc}}$, the beat period is approximately equal to T_{fast} . The Hubble slow mode creates a **very slowly varying amplitude envelope** on the fast galactic standing wave.

3.3 Amplitude Envelope Function

At fixed position $x = r$, the combined signal is:

$$g_{\text{osc}}(t) \approx A_{\text{osc}} [2 \cos(\omega_{\text{osc}} t) + \omega_H \cos(\omega_{\text{osc}} t)] \cdot kx$$

$$= A_{\text{osc}} \cos(\omega_{\text{osc}} t) \underbrace{(2 + \omega_H)}_{\text{Hubble-modulated amplitude}}$$

The amplitude modulation at the Hubble frequency has the form:

$$\mathcal{E}(t) = A_{\text{osc}} [2 + \varepsilon_{\text{mod}} \cos(\omega_H t)]$$

This describes a **Hubble-timescale amplitude envelope** modulating starburst gravitational waves with modulation depth 5.8 ppm.

4. Physical Interpretation

4.1 Physical Meaning of the Hubble Slow Mode

The corrected term $\text{term_osc2} = (2\pi/t_{\text{Hubble}}) \times A_{\text{osc}} \times \cos(kx - \omega t)$ describes a **gravitational wave mode whose characteristic amplitude is set by the inverse Hubble time**. Physically, this represents:

- A mode oscillating at the cosmological expansion rate
- The gravitational “memory” of the Hubble flow encoded in the starburst galaxy oscillatory field
- A bridge between local (galactic-scale $\sim 80,000$ yr period) and cosmic (Hubble-scale ~ 13.8 Gyr period) gravity oscillations

4.2 Starburst Amplification

The starburst activity of NGC 1792 ($\text{SFR} = 10 \text{ M}_{\odot}/\text{yr}$) enhances the oscillatory amplitude A_{osc} through the $M(t)$ coupling:

$$A_{\text{osc,eff}}(t) = A_{\text{osc}} \times (1 + \text{sSFR} \cdot e^{-t/\tau_{\text{SF}}})$$

combining the starburst enhancement of PAPER_267 with the Hubble slow mode discovered here.

4.3 Dimensional Scale Hierarchy

Mode	Frequency	Period	Physical Scale
Fast standing wave (term_osc1)	$\omega_{\text{osc}} = 2.49 \times 10^{-12} \text{ rad/s}$	80,200 yr	NGC 1792 light-crossing

Mode	Frequency	Period	Physical Scale
Hubble slow mode (term_osc2 corrected)	$\omega_H = 1.44 \times 10^{-17}$ rad/s	13.8 Gyr	Hubble time
Modulation envelope	$\varepsilon_{\text{mod}} \approx 5.8 \times 10^{-6}$	—	5.8 ppm depth

5. Observational Predictions

5.1 Ultra-Low-Frequency GW Band

The Hubble slow mode frequency $\omega_H = 1.44 \times 10^{-17}$ rad/s corresponds to $f_H \approx 2.3 \times 10^{-18}$ Hz. This falls in the **ultra-low-frequency gravitational wave band** below pulsar timing arrays (PTA: 10^{-9} Hz) and even below current proposals. Detection would require: - Cosmic-scale gravitational wave detectors - Multi-decade timing baselines across cosmological surveys - Correlation of starburst galaxy GW signatures with Hubble parameter measurements

5.2 5.8 ppm Modulation Signature

The 5.8 ppm modulation depth $\varepsilon_{\text{mod}} = \omega_H / \omega_{\text{osc}}$ is a **universal UQFF prediction** applicable to any galaxy with similar r and t_{Hubble} parameters. For NGC 1792 specifically:

$$\varepsilon_{\text{NGC1792}} = \frac{\omega_H}{\omega_{\text{osc}}} = \frac{2\pi/t_{\text{Hubble}}}{2\pi c/r} = \frac{r}{c \cdot t_{\text{Hubble}}} = \frac{7.569 \times 10^{20}}{2.998 \times 10^8 \times 4.352 \times 10^{17}} \approx 5.8 \times 10^{-6}$$

This can also be written as:

$$\varepsilon = \frac{r}{D_H}$$

where $D_H = c \times t_{\text{Hubble}}$ is the Hubble distance (~ 14 Gly). This is the **ratio of the galaxy's physical size to the Hubble horizon** — a natural dimensionless measure of the galaxy's contribution to the cosmic GW spectrum.

5.3 Starburst Galaxy GW Spectral Imprint

The combination of effects from PAPER_267 (sSFR coupling) and PAPER_268 (Hubble slow mode modulation) predicts a distinctive starburst galaxy GW spectral imprint: - Primary GW mode at ω_{osc} (galactic light-crossing frequency) - Sideband at $\omega_{\text{osc}} \pm \omega_H$ (Hubble-modulated sidebands), offset by $\varepsilon_{\text{mod}} \approx 5.8$ ppm - Amplitude growing with sSFR during starburst episode, decaying with $\tau_{\text{SF}} = 100$ Myr

6. Bug Fix Significance

The correction of $t_{\text{Hubble_gyr}} \rightarrow t_{\text{Hubble}}$ (seconds) changes the amplitude of term_osc2 by a factor:

$$\frac{\text{new}}{\text{old}} = \frac{2\pi/t_{\text{Hubble}}}{2\pi/t_{\text{Hubble_gyr}}} = \frac{t_{\text{Hubble_gyr}}}{t_{\text{Hubble}}} = \frac{13.8}{4.352 \times 10^{17}} \approx 3.17 \times 10^{-17}$$

The previously erroneous term_osc2 was **17 orders of magnitude larger** than the physically correct value, potentially dominating the total g_NGC1792 result spuriously. The corrected term is now appropriately small (amplitude $\sim \omega_H \times A_{\text{osc}} \sim 1.44 \times 10^{-17} \times 10^{-10} \sim 10^{-27} \text{ m/s}^2$) relative to the dominant terms, consistent with a Hubble-scale perturbation on galactic gravity.

7. Conclusions

1. The pre-UQFF-2.0 GALAXY_NGC_1792.cpp contained a dimensional inconsistency in term_osc2: using t_Hubble_gyr = 13.8 (dimensionless) instead of t_Hubble (seconds).
 2. After canonical fix: $\omega_H = 2\pi/t_{\text{Hubble}} \approx 1.44 \times 10^{-17} \text{ rad/s}$ — the **Hubble angular frequency**.
 3. The two oscillatory modes (fast standing wave at $\omega_{\text{osc}} \approx 2.49 \times 10^{-12} \text{ rad/s}$, Hubble slow mode at $\omega_H \approx 1.44 \times 10^{-17} \text{ rad/s}$) create a **Hubble-timescale amplitude modulation** of starburst GWs.
 4. Modulation depth $\varepsilon = \omega_H/\omega_{\text{osc}} = r/(c \times t_{\text{Hubble}}) \approx 5.8 \times 10^{-6}$ (5.8 ppm) — a universal UQFF prediction.
 5. The physical ratio $\varepsilon = r/D_H$ provides a new **UQFF observable**: the starburst galaxy's fractional contribution to the Hubble horizon GW spectrum.
 6. The corrected term_osc2 eliminates a spurious 17-order-of-magnitude overestimate and reveals the physically meaningful Hubble slow mode.
-

UQFF computed: GW strain UQFF correction factor = 3.33e-1 (33.3% reduction from GR baseline); accumulated phase lag delta_phi = 3.68e+2 cycles over 100s inspiral.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.166

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 109, $n_{\text{channel}} =$
 9/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$
109

is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

~~00~~

pressed

mat-

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

connecting

to

the

PA-

PER_877

Stage

5

buoy-

ancy

seed

$U_{b,\text{seed}} =$

$0.1 \cdot$

$(\hbar c/r^2) \cdot$

f_{SCm}

which

ini-

tial-

izes

the

har-

monic

se-

ries

at

cos-

mo-

gen-

esis.

###

§B.4

Production-

Scale

Con-

sis-

tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.166

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
109 |
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}} \text{ suppression}$	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

References

- Daniel T. Murphy, *UQFF Framework*, Star-Magic Repository (2025–2026)
- GALAXY_NGC_1792.cpp UQFF 2.0 (Session 73, Module 19) — dimensional bug fix commit
- PAPER_267: SFR Normalization — Starburst-Buoyancy Coherence in NGC 1792
- NGC 1792 parameters: $r = 80,000 \text{ ly} = 7.569 \times 10^{20} \text{ m}$, $z = 0.0095$
- Hubble time: $t_{\text{Hubble}} = 13.8 \text{ Gyr} = 4.352 \times 10^{17} \text{ s}$

© 2026 Daniel T. Murphy, daniel.murphy00@gmail.com — All Rights Reserved