

PAPER_285: M16 Eagle Nebula UQFF — Erosion Saturation Half-Time and Δg_{Max}

Photoevaporation Asymptotic Saturation: $t_{\text{half}} = \tau \cdot \ln 2 = 2.079 \text{ Myr}$

Classification: UQFF 2.0 Gravitational Physics — Nebular Erosion Dynamics

System: M16 Eagle Nebula (IC 4703), Eagle Nebula Star-Forming Region

Session: 80 | **Module:** M16_UQFF_MODULE.cpp (22nd C++ UQFF module)

Author: Daniel T. Murphy | **Date:** March 2026

Abstract

This paper derives the **Erosion Saturation Half-Time** (t_{half}) and **Maximum Erosion Gravity Amplitude** (Δg_{Max}) for the M16 Eagle Nebula UQFF photoevaporation term. The photoevaporation rate follows an exponential saturation $E_{\text{rad}}(t) = E_0 \times (1 - \exp(-t/\tau))$ with e-folding time $\tau = 3 \text{ Myr}$. The half-erosion time $t_{\text{half}} = \tau \cdot \ln 2 = 6.561 \times 10^{13} \text{ s} = 2.079 \text{ Myr}$ — the time at which the erosion fraction reaches half its asymptotic maximum E_0 . The maximum gravity perturbation is $\Delta g_{\text{Max}} = E_0 \times g_{\text{base}} = 4.36 \times 10^{-13} \text{ m/s}^2$. This is the **first UQFF module** to formally catalogue the photoevaporation half-time and asymptotic erosion concept.

2. Physical Background

UV radiation from massive O-type stars (such as those in the Young Stellar Cluster NGC 6611, embedded in M16) drives photoionisation of surrounding molecular gas — a process called **photoevaporation**. The erosion proceeds not as a linear ramp but as a **saturating exponential**:

$$E_{\text{rad}}(t) = E_0 (1 - e^{-t/\tau})$$

where: - $E_0 = 0.3$ is the **asymptotic maximum fraction** (30% of mass eventually eroded)
- $\tau = 3 \text{ Myr}$ is the **e-folding time** (photoevaporation efficiency timescale) - The saturation arises because dense molecular cores (pillar tips) are progressively shielded by their own column density as surrounding gas is stripped

3. Mathematical Derivation

3.1 Half-Time

The erosion half-time t_{half} is defined as the time when $E_{\text{rad}} = E_0/2$:

$$E_0 (1 - e^{-t_{\text{half}}/\tau}) = \frac{E_0}{2}$$

$$1 - e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$t_{half} = \tau \ln(2)$$

For M16:

$$t_{half} = 9.468 \times 10^{13} \text{ s} \times \ln(2) = 9.468 \times 10^{13} \times 0.6931 = 6.561 \times 10^{13} \text{ s}$$

$$t_{half} = \mathbf{2.079 \text{ Myr}}$$

3.2 Maximum Gravity Amplitude

The maximum erosion-induced gravity perturbation (asymptotic limit as $t \rightarrow \infty$):

$$\Delta g_{Max} = E_0 \times g_{base}$$

For M16:

$$\Delta g_{Max} = 0.3 \times 1.454 \times 10^{-12} = \mathbf{4.36 \times 10^{-13} \text{ m/s}^2}$$

3.3 Peak Erosion Rate

The instantaneous erosion rate at $t = 0$ (maximum rate, before saturation):

$$\left. \frac{dE_{rad}}{dt} \right|_{t=0} = \frac{E_0}{\tau}$$

The corresponding gravity change rate:

$$\left. \frac{dg_{erode}}{dt} \right|_{t=0} = \frac{E_0}{\tau} \times g_{base} = \frac{0.3}{9.468 \times 10^{13}} \times 1.454 \times 10^{-12} = 4.61 \times 10^{-27} \text{ m/s}^2/\text{s}$$

4. Saturation Profile

Time	t (s)	E_rad / E ₀	E_rad	g_erode (m/s ²)
0 Myr	0	0%	0	0
t_half = 2.079 Myr	6.561×10 ¹³	50%	0.150	2.18×10⁻¹³
τ = 3 Myr	9.468×10 ¹³	63.2%	0.190	2.76×10 ⁻¹³
5 Myr	1.578×10 ¹⁴	81.1%	0.243	3.54×10 ⁻¹³
∞ (asymptote)	→ ∞	100%	0.300	4.36×10⁻¹³

Key insight: At $\tau = 3$ Myr (the e-folding time), erosion has consumed only 63.2% of its capacity, NOT 100%. Half-erosion occurs earlier at 2.079 Myr. The pillar structure of M16 means the ~5700 ly “Pillars of Creation” are still observed today because erosion saturates — it cannot fully strip the densest pillar cores within observable timescales.

5. UQFF 2.0 Context

In the full M16 g_{total} equation, the erosion half-time governs the **temporal shape of $g_{\text{dyn}}(t)$** :

$$g_{\text{dyn}}(t) = g_{\text{base}} \times (1 + M_{\text{sf}}) \times (1 - E_0(1 - e^{-t/\tau}))$$

The transition from rapid to slow erosion occurs at $t_{\text{half}} = 2.079$ Myr. For the UQFF simulation (t stepping from 0 to t_{max}), the half-time provides a natural **inflection point** in the dynamic gravity trajectory — before t_{half} , erosion is dominant; after t_{half} , star formation accumulation dominates (since M_{sf} grows linearly while E_{rad} asymptotes).

Crossover Time

The era when SFR growth exactly compensates erosion ($d\Phi_{\text{dm}}/dt = 0$ — maximum Φ_{dm}):

$$\frac{d\Phi_{\text{dm}}}{dt} = \text{SFR}_{\text{rate}} \times (1 - E_{\text{rad}}) - (1 + M_{\text{sf}}) \times \frac{E_0}{\tau} e^{-t/\tau} = 0$$

This crossover defines when the Eagle Nebula achieves maximum effective gravitational influence, after which continued SFR growth dominates erosion.

6. Wolfram KB Term

M16UQFF: $t_{\text{half}}=\tau*\text{Log}[2]=6.561\text{e}13\text{s}=2.079\text{Myr}$; $\text{DeltaGMax}=E_0*g_{\text{base}}=4.36\text{e}-13 \text{ m/s}^2$ [PAPER_285]

7. Cross-References

- **PAPER_284**: Dual Mass Co-Action Product ($\Phi_{\text{dm}} = (1+M_{\text{sf}})\times(1-E_{\text{rad}})$)
 - **PAPER_286**: Nebular Friedmann Redshift (κ_{neb} , $z=0.0015$)
 - **M16_UQFF_MODULE.cpp**: Full UQFF 2.0 C++ implementation (22nd module)
 - **CondensedPhysics3.py**: M16ErosionSaturationHalfTimeCalculator
-

Copyright — Daniel T. Murphy, Session 80, March 2026. UQFF 2.0.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac, [SCm]}} \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.172

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io5

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $53, n_{\text{channel}} =$
 $26/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

53 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁶
yr
 (Jeans
 col-
 lapse
 timescale):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j}.$
 $f_{U_b}.$
 $(1 - e^{-[SSq]\cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.172

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$
|

$p_{\text{DVP}} =$
531
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.