

PAPER_319: Compact HII SFR Gravitational Binding Phase Transition

Author: Daniel T. Murphy **Date:** March 2026 ## $t_{\text{cross}} = 67,730 \text{ yr}$ | $s\text{SFR} = 5 \times 10^{-4} \text{ yr}^{-1}$ (50× Lagoon) | $m_{\text{factor}}(t_{\text{age}}) = 151$ ### FIRST UQFF Compact HII Region SFR Runaway Gravitational Binding Phase Transition

Session: 91

Module: ORION_UQFF_MODULE.cpp (33rd C++ module)

System: Orion Nebula M42/NGC 1976 — compact HII region with active star formation

Watermark: Copyright — Daniel T. Murphy, Session 91, March 2026

Abstract

This paper derives the SFR-driven gravitational binding phase transition for the Orion Nebula. With $\text{SFR} = 1 \text{ M}_{\text{sun}}/\text{yr}$ acting on an initial mass $M = 2000 \text{ M}_{\text{sun}}$, the specific star formation rate **$s\text{SFR} = 5 \times 10^{-4} \text{ yr}^{-1}$ is 50× that of the Lagoon Nebula** ($1 \times 10^{-5} \text{ yr}^{-1}$, PAPER_305). The system is born wind-dominated (unbound), but as SFR continuously adds mass, the effective gravitational acceleration amplified by the SFR mass factor $m_{\text{factor}}(t) = 1 + \text{SFR} \times t_{\text{yr}} / M_{\text{sun_count}}$ crosses the growing wind ram pressure at **$t_{\text{cross}} = 67,730 \text{ yr}$** , transitioning the nebula from unbound to gravitationally bound. By $t_{\text{age}} = 300 \text{ kyr}$, $\text{binding_ratio} = g_{\text{SFR}}/a_{\text{wind}} = 2.654$. This is the FIRST UQFF compact HII SFR runaway gravitational binding phase transition.

System Parameters

Parameter	Value	Description
M	$2000 \text{ M}_{\text{sun}} = 3.978 \times 10^{33} \text{ kg}$	Initial mass
SFR	$1 \text{ M}_{\text{sun}}/\text{yr} = 6.303 \times 10^{22} \text{ kg/s}$	Star formation rate
sSFR	$5 \times 10^{-4} \text{ yr}^{-1}$	Specific SFR = SFR/M
t_{age}	$3 \times 10^5 \text{ yr}$	Nebula age
v_{wind}	$8 \times 10^3 \text{ m/s}$	HII ionization front
g_{base}	$1.907 \times 10^{-11} \text{ m/s}^2$	Base gravity (PAPER_317)
$a_{\text{wind}}(t=0)$	$5.424 \times 10^{-10} \text{ m/s}^2$	Initial wind acceleration

Key Equations

SFR Mass Factor

$$M_{\text{sf}}(t) = \frac{\text{SFR}_{\text{yr}} \times t_{\text{yr}}}{M_{\text{sun,count}}}, \quad m_{\text{factor}}(t) = 1 + M_{\text{sf}}(t)$$

SFR-Amplified Gravitational Acceleration

$$g_{\text{SFR}}(t) = g_{\text{base}} \times m_{\text{factor}}(t) = g_{\text{base}} \left(1 + \frac{\text{SFR}_{\text{yr}} \cdot t_{\text{yr}}}{M_{\text{sun,count}}} \right)$$

Wind Acceleration (time-evolving)

$$a_{\text{wind}}(t) = \frac{v_{\text{wind}}^2}{r} \left(1 + \frac{t}{t_{\text{age}}} \right)$$

Crossover Time t_{cross} (PAPER_319 Key Result)

Setting $g_{\text{SFR}}(t) = a_{\text{wind}}(t)$:

$$g_{\text{base}} (1 + \text{sSFR} \cdot t) = a_{\text{wind},0} \left(1 + \frac{t}{t_{\text{age,yr}}} \right)$$

$$t_{\text{cross}} = \frac{a_{\text{wind},0} - g_{\text{base}}}{g_{\text{base}} \cdot \text{sSFR} - a_{\text{wind},0}/t_{\text{age,yr}}} = \boxed{67,730 \text{ yr}}$$

Specific SFR

$$\text{sSFR} = \frac{\text{SFR}_{\text{yr}}}{M_{\text{sun,count}}} = \frac{1}{2000} = 5 \times 10^{-4} \text{ yr}^{-1} = 50 \times \text{Lagoon}$$

Gas Depletion Timescale

$$t_{\text{consume}} = \frac{M_{\text{sun,count}}}{\text{SFR}_{\text{yr}}} = \frac{2000}{1} = 2000 \text{ yr} \quad (\text{without OMC-1 replenishment})$$

Results Summary

Quantity	Value	Significance
sSFR	$5 \times 10^{-4} \text{ yr}^{-1}$	50× Lagoon Nebula
t_{cross}	67,730 yr	Unbound → bound transition
$m_{\text{factor}}(t_{\text{age}}=300 \text{ kyr})$	151	SFR amplification
$g_{\text{SFR}}(t_{\text{age}})$	$2.878 \times 10^{-9} \text{ m/s}^2$	$151 \times g_{\text{base}}$
$a_{\text{wind}}(t_{\text{age}})$	$1.085 \times 10^{-9} \text{ m/s}^2$	
$\text{binding_ratio}(t_{\text{age}})$	2.654	Gravitationally bound
$m_{\text{factor}}(1 \text{ Myr})$	501	
$\text{binding_ratio}(1 \text{ Myr})$	4.069	Increasingly bound
t_{consume}	2000 yr	Gas depletion (no replenishment)

Phase Transition Diagram

$t=0$	$t_{\text{cross}}=67.7 \text{ kyr}$	$t_{\text{age}}=300 \text{ kyr}$	$t=1 \text{ Myr}$
UNBOUND	TRANSITION	BOUND $2.65 \times$	BOUND $4.07 \times$
$a_{\text{wind}} > g_{\text{SFR}}$	$a_{\text{wind}} = g_{\text{SFR}}$	$g_{\text{SFR}} > a_{\text{wind}}$	$g_{\text{SFR}} \gg a_{\text{wind}}$
$\eta=28.47 \text{ wind}$	— crossover —	$\eta_{\text{B}}=2.654$	$\eta_{\text{B}}=4.069$

Comparison: UQFF SFR Class

Module	SFR (M_{sun}/yr)	M (M_{sun})	sSFR (yr^{-1})	PAPER
Lagoon	0.1	10,000	1×10^{-5}	PAPER_305
Orion	1.0	2000	5×10^{-4}	PAPER_319
M16 (Eagle)	$\sim 0.01 \times$	1200	—	PAPER_284

Orion sSFR is $50 \times$ Lagoon — UQFF identifies this as the “ultra-compact HII” class with rapid gravitational binding crossover. Lagoon has $t_{\text{consume}} = 100 \text{ kyr}$; Orion has $t_{\text{consume}} = \mathbf{2000 \text{ yr}}$, the shortest gas depletion time in the UQFF series, sustained only by continuous OMC-1 giant molecular cloud inflow.

UQFF Physical Interpretation

The phase transition at $t_{\text{cross}} = 67,730 \text{ yr}$ is a structural boundary in Orion’s evolution:

- **Before t_{cross} :** The system is wind-dominated ($\eta_{\text{wind}} > 1$). UV photoionization and ram pressure drive a champagne flow outward; newly formed stars are subject to wind erosion.
- **After t_{cross} :** SFR has added sufficient mass that $g_{\text{SFR}} > a_{\text{wind}}$. The system becomes self-gravitating with respect to its own stellar formation. The cluster proceeds to form stars under its own gravity — consistent with the Orion OB1 association framework where the cluster is now 300 kyr old and gravitationally bound.

Within ORION_UQFF_MODULE.cpp, the SFR mass factor $m_{\text{factor}}(t)$ enters the base gravity term via $G*M*(1+M_{\text{sf}}(t))/r^2$, registered as WOLFRAM_TERM_ORION_BASE and WOLFRAM_TERM_ORION_SFR_BINDING.

WOLFRAM_TERM Registration

```
#define WOLFRAM_TERM_ORION_BASE(val) (val)
// g_base = (G*M*(1+M_sf(t))/r^2)*(1+Hz*t)*(1-B/B_crit)*(1+f_TRZ)
// M_sf(t) = SFR_yr*t_yr / M_sun_count [PAPER_319 SFR amplification]
```

```
#define WOLFRAM_TERM_ORION_SFR_BINDING(val) (val)
// wind = v_wind^2/r * (1+t/t_age) [crosses SFR gravity at t_cross=67730 yr]
```

Series first: *FIRST* UQFF compact HII SFR runaway gravitational binding phase transition. Establishes sSFR as a new UQFF classification axis for HII regions: compact (Orion, $s\text{SFR}=5 \times 10^{-4} \text{ yr}^{-1}$) vs extended (Lagoon, $s\text{SFR}=1 \times 10^{-5} \text{ yr}^{-1}$).

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.157

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io5

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $71, n_{\text{channel}} =$
 8/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

71 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.157

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
711
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.