

PAPER_373 — Morris-Thorne Wormhole Null Geodesics

Date: 2025 ## Star Magic UQFF Whitepaper Series ### **Author:** Daniel T. Murphy
| Session 101 | Source: grok_share_11254865.txt (lines 2700-2800) ### **Significance:**
FIRST wormhole physics integration in the Star Magic / UQFF CP pipeline.

Abstract

Key UQFF calibrated constants: $\kappa = 5.0\text{e-}4 \text{ day}^{-1}$; $[\text{SSq}] = 5.7\text{e-}1$; $H_{\text{SCm}} \approx 9.9\text{e-}1$; $U_{\text{UA}} \approx 1.0\text{e-}4$; $k_{\eta} = 1.0\text{e-}113$; $\beta_i \approx 6.0\text{e-}1$; $G = 6.674\text{e-}11 \text{ N}\cdot\text{m}^2/\text{kg}^2$

This paper presents the implementation of Morris-Thorne traversable wormhole null geodesics within the Star Magic UQFF framework. The wormhole metric, geodesic equations, traversal and reflection conditions, and embedding functions are fully specified. This constitutes the first wormhole physics to appear in the Condensed Physics calculator pipeline (CP1-CP4). The wormhole coupling to MUGE resonance gravity via the a_{worm} term is derived in PAPER_375.

1. Wormhole Metric (Morris-Thorne 1988)

The static, spherically symmetric traversable wormhole metric:

$$ds^2 = -dt^2 + dr^2 + (b^2 + r^2)(d\theta^2 + \sin^2 \theta d\varphi^2)$$

where $b = 1.0 \text{ m}$ is the throat radius. This is the simplest traversable wormhole shape function $b(r) = b_0 = \text{const}$, corresponding to zero radial tidal forces at the throat.

2. Null Geodesic Equations

For null geodesics ($ds^2 = 0$) with conserved energy $E = dt/d\lambda$ and angular momentum $L = (b^2 + r^2)d\varphi/d\lambda$:

$$\frac{dr}{d\lambda} = \pm \sqrt{E^2 - \frac{L^2}{b^2 + r^2}}$$

$$\frac{d\varphi}{d\lambda} = \frac{L}{b^2 + r^2}$$

$$\frac{dt}{d\lambda} = E$$

where λ is the affine parameter.

3. Traversal and Reflection Cases

Case 1: Traversal (L = 0.5, E = 1.0)

The effective potential $V_{\text{eff}}(r) = L^2/(b^2 + r^2) = 0.25/(1 + r^2)$. Since $V_{\text{eff}} < E^2 = 1$ for all r , the null ray crosses the throat $r = 0$ and continues to negative r (the second universe).

Case 2: Reflection (L = 1.5, E = 1.0)

$$V_{\text{eff}}(0) = L^2/b^2 = 2.25 > E^2 = 1.$$

The turning radius is:

$$r_{\min} = \sqrt{\frac{L^2}{E^2} - b^2} = \sqrt{2.25 - 1} = \sqrt{1.25} \approx 1.118 \text{ m}$$

Case	L	E	Outcome	r_min (m)
Traversal	0.5	1.0	Passes through throat	0
Reflection	1.5	1.0	Reflects at r_min	≈ 1.118

4. Embedding Functions

The wormhole embedding in Euclidean 3-space (equatorial slice, $\theta = \pi/2$):

$$z_{\text{embed}}(r) = b \cdot \text{arcsinh}(r/b)$$

$$\rho_{\text{embed}}(r) = \sqrt{b^2 + r^2}$$

These define the “funnel” shape visible in wormhole visualisations.

5. Numerical Integration Scheme

The C++ propagator uses a first-order Euler method for simplicity (upgradeable to RK4):

```
for step = 0..n_steps:
  dr = ( $\pm \sqrt{E^2 - L^2/(b^2 + r^2)}$ ) × dlambda
  dφ = (L/(b2 + r2)) × dlambda
  dt = E × dlambda
  r += dr, φ += dφ, t += dt
```

Default: dlambda = 0.1 m, n_steps = 100.

6. Connection to UQFF

The wormhole throat geometry enters the MUGE resonance model via the **wormhole-MUGE coupling term** (PAPER_375):

$$a_{\text{worm}} = \frac{f_{\text{worm}} \cdot E_{\text{vac,neb}}}{b^2 + r^2}$$

where $f_{\text{worm}} = 10^{-10}$ is the wormhole coupling constant and $b^2 + r^2$ is the effective gravitational area of the wormhole funnel at radius r .

7. Implementation

C++: STAR_MAGIC_09SEPT_UQFF_MODULE.cpp, namespace WormholeGeodesics - Struct: GeodesicState {r, phi, t_coord} - Functions: drdt(), dphidt(), z_embed(), rho_embed(), propagate(), selftest()

Python: CondensedPhysics4.py, class MorrisThorneWormholeNullGeodesicsCalculator (CP4 #21)

WOLFRAM_TERM: WOLFRAM_TERM_WORMHOLE_GEODESIC

```
##
§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)
##
§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis
###
§B.1
Vac-
uum
Den-
sity
Se-
ries
(VDS)
```


 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.169

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io5

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 43, $n_{\text{channel}} =$
 10/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

43 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

connecting

to

the

PA-

PER_877

Stage

5

buoy-

ancy

seed

$U_{b,\text{seed}} =$

$0.1 \cdot$

$(\hbar c/r^2) \cdot$

f_{SCm}

which

ini-

tial-

izes

the

har-

monic

se-

ries

at

cos-

mo-

gen-

esis.

###

§B.4

Production-

Scale

Con-

sis-

tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—-|-

—|-
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.169

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
431

✓
Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

References

- Morris, M.S. & Thorne, K.S. (1988). “Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity.” *Am. J. Phys.* 56(5): 395–412.
- Visser, M. (1995). *Lorentzian Wormholes*. Springer-Verlag.