

PAPER_438 — Galaxy NGC 2525: Per-System MUGE with M_SN(t) Supernova Mass-Loss and g_BH Proximity

Author: Daniel T. Murphy **Date:** 2025

Source: grok_share_68eb34022.txt — Document 10: “Master Universal Gravity Equation_Galaxy NGC 2525 Evolution_03May2025.docx” (lines 3085–3429) **Session:** 119 **CP4 Class:** NGC2525PerSystemMUGE_SNMassLoss_BHProximity_Calculator (#93)

Abstract

This paper presents a UQFF analysis of Galaxy NGC 2525: Per-System MUGE with M_SN(t) Supernova Mass-Loss and g_BH Proximity, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

1. Overview

PAPER_438 delivers the **complete per-system MUGE** for NGC 2525 — a barred spiral galaxy at $z \approx 0.016$ ($d \approx 70$ Mpc) famous as the host of SN 2018gv (Type Ia supernova observed in Hubble Year 29 anniversary images). The total galaxy mass is $M \approx 10^{10} M_\odot$, with a central supermassive black hole $M_{\text{BH}} \approx 2.25 \times 10^7 M_\odot$ at $r_{\text{BH}} = 1.496 \times 10^{11}$ m (1 AU — representing the SMBH influence sphere inner boundary).

Novel claim (Q1): First UQFF MUGE incorporating **SN Ia mass loss as a negative gravitational term:** $T_{\text{SN}} = -GM_{\text{SN}}(t)/r^2$ where $M_{\text{SN}}(t) = 1.4 M_\odot e^{-t/\tau_{\text{SN}}}$ with $\tau_{\text{SN}} = 1$ yr, plus a simultaneous SMBH proximity term $g_{\text{BH}} = GM_{\text{BH}}/r_{\text{BH}}^2$ — establishing the first MUGE where supernova mass ejection directly reduces the effective gravitational field of the parent galaxy.

2. System Parameters

Parameter	Symbol	Value
Total galaxy mass	M	$(10^{10} + 2.25 \times 10^7) M_\odot \approx 10^{10} M_\odot$
Galaxy radius	r	9.2 kpc $= 2.836 \times 10^{20}$ m
Galaxy redshift	z	0.016
Hubble at z	$H(z)$	$\approx 2.19 \times 10^{-18} \text{ s}^{-1}$
SMBH mass	M_{BH}	$2.25 \times 10^7 M_\odot =$ $4.475 \times 10^{37} \text{ kg}$
SMBH influence radius	r_{BH}	$1.496 \times 10^{11} \text{ m}$ (1 AU)
SN Ia ejecta mass	M_{SN0}	$1.4 M_\odot = 2.785 \times 10^{30}$ kg (Chandrasekhar)
SN decay timescale	τ_{SN}	1 yr $= 3.156 \times 10^7 \text{ s}$
Magnetic field	B	10^{-5} T

3. Time-Dependent Functions

SN mass loss:

$$M_{\text{SN}}(t) = 1.4 M_{\odot} e^{-t/\tau_{\text{SN}}} = 2.785 \times 10^{30} e^{-t/3.156 \times 10^7} \text{ kg}$$

At $t = 0$: $M_{\text{SN}} = 1.4 M_{\odot}$ (SN peak — Chandrasekhar-mass progenitor)

At $t = 1 \text{ yr}$: $M_{\text{SN}} = 0.515 M_{\odot}$ (ejecta dispersed at 1/e)

At $t = 10 \text{ yr}$: $M_{\text{SN}} \rightarrow 0$ (remnant phase)

4. Complete 10-Term MUGE

$$g_{\text{N2525}}(r, t) = T_1 + T_{\text{BH}} + T_2 + T_3 + T_4 + T_5 + T_6 + T_7 + T_8 + T_{\text{SN}}$$

T1 — Newtonian + H(z)t + B:

$$T_1 = \frac{GM}{r^2}(1 + H(z)t) \left(1 - \frac{B}{B_{\text{crit}}}\right) \approx \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{40}}{(2.836 \times 10^{20})^2} \approx 1.65 \times 10^{-11} \text{ m/s}^2$$

T_{BH} — SMBH proximity gravity:

$$T_{\text{BH}} = \frac{GM_{\text{BH}}}{r_{\text{BH}}^2} = \frac{6.674 \times 10^{-11} \times 4.475 \times 10^{37}}{(1.496 \times 10^{11})^2} = \frac{2.984 \times 10^{27}}{2.238 \times 10^{22}} \approx 1.334 \times 10^5 \text{ m/s}^2$$

T2 — UQFF Ug with f_{TRZ}:

$$T_2 = 2 \times 1.65 \times 10^{-11} \times 1.1 \approx 3.63 \times 10^{-11} \text{ m/s}^2$$

T3 — Λ : $\sim 3.3 \times 10^{-36} \text{ m/s}^2$ (negligible)

T4 — Quantum: negligible

T5 — Scaled EM: $\sim 10^{-24} \text{ m/s}^2$ (negligible)

T6 — Fluid: minor

T7 — Oscillatory spiral arm modes: minor (oscillatory density waves)

T8 — DM perturbation:

$$T_8 \approx \frac{(M + 0.1M)\delta\rho/\rho + 3GM/r^3}{M} \sim 10^{-11} \text{ m/s}^2$$

T_{SN} — SN Ia mass loss (negative term):

$$T_{\text{SN}}(t) = -\frac{GM_{\text{SN}}(t)}{r^2} = -\frac{6.674 \times 10^{-11} \times 2.785 \times 10^{30}}{(2.836 \times 10^{20})^2} \times e^{-t/\tau_{\text{SN}}} \approx -2.31 \times 10^{-21} e^{-t/\tau} \text{ m/s}^2$$

The SN term is $\sim 10^{10}$ times smaller than the galaxy self-gravity — negligible at galaxy scale but **observable at the SN remnant radius** ($r \sim 1 \text{ pc}$ where $T_{\text{SN}} \gg T_1$).

5. Canonical Numerical Result

At $t = 0$, $r = r_{\text{galaxy}} = 2.836 \times 10^{20}$ m:

Term	Value (m/s ²)	Notes
T_{BH}	$+1.334 \times 10^5$	Dominant (at $r_{\text{BH}} = 1$ AU)
T_2 UQFF Ug	$+3.63 \times 10^{-11}$	Primary at galaxy scale
T_1 Newtonian	$+1.65 \times 10^{-11}$	Standard
T_8 DM	$+ \sim 10^{-11}$	Minor
T_{SN} (galaxy scale)	-2.31×10^{-21}	Negligible at galaxy r

Note: T_{BH} evaluated at r_{BH} represents the SMBH inner sphere — appropriate for the Gaia/VLBI proper motion frame near the nucleus.

$$g_{\text{N2525}}^{\text{galaxy}} \approx 3.63 \times 10^{-11} \text{ m/s}^2 \quad [\text{UQFF Ug dominant at disk scale}]$$

6. Uniqueness vs Prior Papers

Prior Paper	System	Overlap	New in PAPER_438
PAPER_383	NGC 2525 tail Δ_{N2525}	2-line summary	Full 10-term + SN term derivation
PAPER_422	System 10: NGC 2525 tail	Brief	Complete numerical evaluation
None	SN Ia $T_{\text{SN}} = -GM_{\text{SN}}/r^2$	N/A	First UQFF SN mass-loss term

7. Comparison to Standard Model

SM galactic dynamics ignore individual SN mass loss ($1.4M_{\odot}$ vs $M_{\text{gal}} = 10^{10} M_{\odot} = 0$ ppm). UQFF preserves this term as a matter of completeness and precision — it becomes **significant at** $r \sim 0.1$ pc (SN remnant interior), predicting a velocity structure different from standard Sedov-Taylor blast wave models by the additional $GM_{\text{SN}}(t)/r_{\text{SN}}^2$ gravitational retardation term.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.112

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io5

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $67, \quad n_{\text{channel}} =$
 $23/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

67 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁶
M_{BH}/M_⊙
yr
 (quasi-
 normal
 mode
 ring-
 down):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—-|-

—|-
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.112
| ✓

Threshold-
consistent
||

DVP
prime
|

$p_k \in$
 $\{2,3,...,113\}$
|
 $p_{\text{DVP}} =$
671
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Thomson σ_T (QED synchrotron)	UQFF U_m scattering kernel: $\sigma_T = 6.6524e-29 \text{ m}^2$	$\sigma_T = 6.6524e-29 \text{ m}^2$ (PDG QED exact)	PDG 2024	100% (exact QED input)
NGC 2525 SN Host luminosity Optical + X-ray GR Schwarzschild limit κ vacuum rate vs X-ray variability	UQFF MUGE g_total $\rightarrow L_X$ via Stefan-Boltzmann + buoyancy flux: $L_X \approx g_{\text{total}} \times M_{\text{env}}$ UQFF g_total must satisfy $g \leq c^2/(2r_s)$ at event horizon UQFF $\kappa = 0.0005/\text{day} \rightarrow$ timescale $\tau_{\text{UQFF}} = 2000 \text{ days}$	$L_X \text{ SN } M_V \sim -19 \text{ mag}$ $r_s = 2GM/c^2$ (GR exact) Observed X-ray variability τ_{obs} (instrument monitoring)	HST + Chandra PDG 2024 / GR HST + Chandra	✓ Consistent order of magnitude ✓ UQFF respects GR horizon Testable UQFF variability timescale

New physics claim: UQFF MUGE generates gravity enhancement factors ($g_{\text{total}}/g_{\text{Newt}} > 1$) for NGC 2525 SN Host through vacuum buoyancy coupling — a mechanism absent from GR+SM. The enhancement factor and X-ray luminosity are linked via the UQFF buoyancy flux, providing a testable prediction for future HST + Chandra monitoring observations.

Cite *PAPER_642* (*UQFFSMParameterBridgeMasterComparisonCalculator*) for full UQFF-SM bridge.

8. Testable Predictions

Q5 Prediction 1: $M_{\text{SN}}(t) = 1.4M_{\odot}e^{-t/1 \text{ yr}}$ predicts that by day 365 after SN 2018gv maximum, the SN Ia ejecta mass contributing to the gravitational term has decayed to $0.515M_{\odot}$ — testable by comparing the UQFF gravity retardation model to the observed deceleration of the SN 2018gv photosphere expansion (Hubble spectroscopic monitoring).

Q5 Prediction 2: The SMBH proximity term $T_{\text{BH}} = GM_{\text{BH}}/r_{\text{BH}}^2 \approx 1.334 \times 10^5 \text{ m/s}^2$ at $r_{\text{BH}} = 1 \text{ AU}$ predicts a stellar velocity dispersion at $r < r_{\text{BH}}$ scaling as $v \propto \sqrt{T_{\text{BH}} \times r_{\text{BH}}} \approx 4.2 \times 10^3 \text{ m/s} = 4.2 \text{ km/s}$ — consistent with the NGC 2525 SMBH mass scaling relation ($\sigma\text{-}M_{\text{BH}}$) for a $10^7 M_{\odot}$ black hole.

Q5 Prediction 3: The UQFF $f_{\text{TRZ}} = 0.1$ factor predicts a 10% periodic oscillation in the galaxy rotation curve at $\omega_{\text{TRZ}} = v_{\text{gas}}/(r) \sim 10^{-16} \text{ rad/s}$ — a very slow pattern speed observable as a $\sim 10\%$ arm-to-interarm density contrast in deep HI 21-cm mapping of NGC 2525.