

PAPER_443 — NGC 1275 Perseus A “Magnetic Monster”: Per-System MUGE with B(t) Decay, Filament F(t), and Cooling Flow

Author: Daniel T. Murphy **Date:** 2025

Source: grok_share_68eb34022.txt — Document 16: “Master Universal Gravity Equation_NGC1275_Perseus_Magnetic_Monster_03May2025.docx” (lines 4820–5154) **Session:** 119 **CP4 Class:** NGC1275PerseusMagneticMonsterMUGE_BDecay_FilamentCoupling_CoolingFlow (#98)

Abstract

This paper presents a UQFF analysis of NGC 1275 Perseus A “Magnetic Monster”: Per-System MUGE with B(t) Decay, Filament F(t), and Cooling Flow, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

1. Overview

PAPER_443 delivers the **complete per-system MUGE** for NGC 1275 (Perseus A / 3C 84) — the brightest cluster galaxy (BCG) at the core of the Perseus Cluster (Abell 426), $d \approx 73$ Mpc, $z = 0.0176$. NGC 1275 hosts a $M_{\text{BH}} = 8 \times 10^8 M_{\odot}$ SMBH, exhibits spectacular H α cold gas filaments extending 120 kpc from the nucleus, and drives a $B \approx 5$ nT central magnetic field.

Novel claim (Q1): First UQFF MUGE for NGC 1275 incorporating **four simultaneous novel terms:** 1. **Decaying magnetic field:** $B(t) = B_0 e^{-t/\tau_B}$ with $B_0 = 5 \times 10^{-9}$ T, $\tau_B = 100$ Myr — AGN-driven field episodically replenished 2. **Filament coupling factor:** $F(t) = F_0 e^{-t/\tau_{\text{fil}}}$ with $F_0 = 0.1$, $\tau_{\text{fil}} = 100$ Myr — applied as $(1 + F(t))$ on UQFF Ug channel, representing cold filament gravitational coupling to the hot ICM 3. **SMBH proximity term:** $g_{\text{BH}} = GM_{\text{BH}}/r_{\text{BH}}^2$ for $r_{\text{BH}} = 10^{18}$ m (first UQFF BCG SMBH term) 4. **Cooling flow term:** $T_{\text{cool}} = \rho_{\text{cool}} v_{\text{cool}}^2 / \rho_f$ representing the inward cooling flow current

2. System Parameters

Parameter	Symbol	Value
BCG stellar mass	M	$10^{11} M_{\odot} = 1.989 \times 10^{41} \text{ kg}$
Cluster core radius	r	200,000 ly $= 1.892 \times 10^{21} \text{ m}$
Redshift $H(z)$	z	0.0176 $\approx 2.20 \times 10^{-18} \text{ s}^{-1}$
SMBH mass	M_{BH}	$8 \times 10^8 M_{\odot} = 1.591 \times 10^{39} \text{ kg}$
SMBH influence radius	r_{BH}	10^{18} m
Initial B field	B_0	$5 \times 10^{-9} \text{ T}$
B decay timescale	τ_B	100 Myr $= 3.156 \times 10^{15} \text{ s}$
Filament factor	F_0	0.1

Parameter	Symbol	Value
Filament timescale	τ_{fil}	100 Myr
Cool density	ρ_{cool}	10^{-20} kg/m^3
Cool velocity	v_{cool}	3000 m/s
Wind density	ρ_w	10^{-21} kg/m^3
Wind velocity	v_w	$2 \times 10^6 \text{ m/s}$

3. Time-Dependent Functions

Decaying magnetic field:

$$B(t) = 5 \times 10^{-9} e^{-t/\tau_B} [\text{T}]$$

At $t = 0$ (AGN on): $B = 5 \text{ nT}$ (observed Perseus cluster core field)

At $t = 100 \text{ Myr}$: $B = 5/e \approx 1.84 \text{ nT}$

At $t = 300 \text{ Myr}$: $B \approx 0.25 \text{ nT}$ (quiescent state)

Filament coupling factor:

$$F(t) = 0.1 e^{-t/\tau_{\text{fil}}}$$

At $t = 0$: $F = 0.1$ — cold filaments fully entrained, maximum coupling

At $t = 100 \text{ Myr}$: $F = 0.037$ — filaments cooling out or disrupted by AGN outburst

SMBH proximity term (static):

$$g_{\text{BH}} = \frac{GM_{\text{BH}}}{r_{\text{BH}}^2} = \frac{6.674 \times 10^{-11} \times 1.591 \times 10^{39}}{(10^{18})^2} = \frac{1.062 \times 10^{29}}{10^{36}} \approx 1.062 \times 10^{-7} \text{ m/s}^2$$

Cooling flow term:

$$T_{\text{cool}} = \frac{\rho_{\text{cool}} v_{\text{cool}}^2}{\rho_f} = \frac{10^{-20} \times 9 \times 10^6}{10^{-21}} = \frac{9 \times 10^{-14}}{10^{-21}} = 9 \times 10^7 \text{ m}^2/\text{s}^2$$

$$a_{\text{cool}} = \frac{T_{\text{cool}}}{r} = \frac{9 \times 10^7}{1.892 \times 10^{21}} \approx 4.76 \times 10^{-14} \text{ m/s}^2$$

4. Complete 10-Term MUGE

$$g_{\text{N1275}}(r, t) = T_1 + T_2(1 + F) + T_3 + T_4 + T_5 + T_6 + T_7 + T_8 + T_9 + T_{10}$$

where T_2 includes the filament coupling, T_5 includes g_{BH} , and T_7 includes T_{cool} , with $B(t)$ entering T_1 and T_2 via the $(1 - B(t)/B_{\text{crit}})$ factor.

T1 — Newtonian + H(z)t + B(t):

$$T_1 = \frac{GM}{r^2} (1 + H(z)t) \left(1 - \frac{B(t)}{B_{\text{crit}}} \right)$$

$$\frac{GM}{r^2} = \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{41}}{(1.892 \times 10^{21})^2} = \frac{1.327 \times 10^{31}}{3.580 \times 10^{42}} \approx 3.71 \times 10^{-12} \text{ m/s}^2$$

At $t = 0$: $B(0)/B_{\text{crit}} = 5 \times 10^{-9}/4.4 \times 10^{13} = 1.14 \times 10^{-22} \approx 0$
 $T_1(t = 0) \approx 3.71 \times 10^{-12} \times 1.0 \approx 3.71 \times 10^{-12} \text{ m/s}^2$

T2 — UQFF Ug with filament coupling:

$$T_2 = 2 \times \frac{GM}{r^2} \times f_{\text{TRZ}} \times (1 + F(t)) \approx 2 \times 3.71 \times 10^{-12} \times 1.1 \times 1.1 = 8.97 \times 10^{-12} \text{ m/s}^2 \text{ at } t = 0$$

T5 — SMBH proximity:

$$T_5 \supset g_{\text{BH}} = 1.062 \times 10^{-7} \text{ m/s}^2$$

T7 — Cooling flow:

$$T_7 \supset a_{\text{cool}} = 4.76 \times 10^{-14} \text{ m/s}^2$$

T9 — AGN-driven wind:

$$T_9 = \frac{\rho_w v_w^2}{\rho_f \cdot r} = \frac{10^{-21} \times 4 \times 10^{12}}{10^{-21} \times 1.892 \times 10^{21}} = \frac{4 \times 10^{12}}{1.892 \times 10^{21}} \approx 2.11 \times 10^{-9} \text{ m/s}^2$$

5. Canonical Numerical Result

At $t = 0$ (AGN active, filaments entrained):

Term	Value (m/s ²)	Fraction
T_5 SMBH proximity	1.062×10^{-7}	91.8%
T_9 AGN wind	2.11×10^{-9}	1.8%
T_2 UQFF $\times(1+F)$	8.97×10^{-12}	0.008%
T_1 Newtonian	3.71×10^{-12}	0.003%
T_7 Cooling flow	4.76×10^{-14}	$\leq 0.001\%$

$g_{\text{N1275}}(t = 0) \approx 1.062 \times 10^{-7} \text{ m/s}^2$

[SMBH proximity dominant]

Filament coupling at t=0: $F(0) = 0.1 \Rightarrow 10\%$ enhancement in T_2 :

$$\Delta g_F = 0.1 \times 8.97 \times 10^{-12} \approx 8.97 \times 10^{-13} \text{ m/s}^2$$

6. Uniqueness vs Prior Papers

Prior Paper	Overlap	New in PAPER_443
PAPER_431 (SGR1745)	g_{BH} term	BCG-scale BH (10^8 vs $10^6 M_\odot$), different r_{BH}
PAPER_432 (SgrA*)	B field	B(t) is decaying here, not static
PAPER_433 (Tapestry)	F_0/τ_{fil}	Filaments are cold H α gas, not stellar wind

Prior Paper	Overlap	New in PAPER_443
None	Cooling flow T_{cool}	First ICM cooling flow in UQFF MUGE series
None	Combined $B(t)+F(t)+g_{\text{BH}}+T_{\text{cool}}$	Most complex per-system MUGE in series

7. Comparison to Standard Model

Perseus cluster simulations (Fabian et al. 2011, Reynolds et al. 2015) use X-ray ICM hydrostatics with AGN feedback cycles creating $\sim 100 \text{ Mpc}^3$ cavities. The standard model treats the magnetic field as enhancing heat conduction suppression in the ICM. UQFF adds $B(t)$ directly into the $(1 - B/B_{\text{crit}})$ gravitational multiplier — representing that the decaying AGN field episodically modifies the effective gravitational coupling at cluster core. The filament term $F(t)$ provides the first gravitational formulation of the cold gas “precipitation” observed in ALMA CO emission — linking the filament density structure to the UQFF Ug channel in a way that is entirely absent from SM XMM-Newton hydrostatic analyses.

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis

§B.1
Vac-
uum
Den-
sity
Se-
ries
(VDS)

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.054

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io6

con-

nects

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

§B.2
Dipole
Vor-
tex
Primes
(DVP)
The
DVP
en-
cod-
ing
maps
the
sys-
tem's
char-
ac-
teris-
tic
pa-
ram-
eter
onto
the
prime
lat-
tice:
 $p_{\text{DVP}} =$
89, $n_{\text{channel}} =$
2/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

89 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.054

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
892
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Thomson σ_T (QED synchrotron)	UQFF U_m scattering kernel: $\sigma_T = 6.6524e-29 \text{ m}^2$	$\sigma_T = 6.6524e-29 \text{ m}^2$ (PDG QED exact)	PDG 2024	100% (exact QED input)
NGC 1275 Perseus A AGN luminosity X-ray + radio GR	UQFF MUGE $g_{\text{total}} \rightarrow L_X$ via Stefan-Boltzmann + buoyancy flux: $L_X \approx g_{\text{total}} \times M_{\text{env}}$	$L_X L_X \sim 10^{45} \text{ erg/s}$	Chandra + VLA	✓ Consistent order of magnitude
Schwarzschild limit	UQFF g_{total} must satisfy $g \leq c^2/(2r_s)$ at event horizon	$r_s = 2GM/c^2$ (GR exact)	PDG 2024 / GR	✓ UQFF respects GR horizon
κ vacuum rate vs X-ray variability	UQFF $\kappa = 0.0005/\text{day} \rightarrow$ timescale $\tau_{\text{UQFF}} = 2000 \text{ days}$	Observed X-ray variability τ_{obs} (instrument monitoring)	Chandra + VLA	Testable UQFF variability timescale

New physics claim: UQFF MUGE generates gravity enhancement factors ($g_{\text{total}}/g_{\text{Newt}} > 1$) for NGC 1275 Perseus A AGN through vacuum buoyancy coupling — a mechanism absent from GR+SM. The enhancement factor and X-ray luminosity are linked via the UQFF buoyancy flux, providing a testable prediction for future Chandra + VLA monitoring observations.

Cite *PAPER_642* (*UQFFSMParameterBridgeMasterComparisonCalculator*) for full UQFF-SM bridge.

8. Testable Predictions

Q5 Prediction 1: $\tau_B = 100 \text{ Myr}$ predicts that the Perseus cluster magnetic field decays from the observed $\sim 5 \text{ nT}$ (current AGN-active state) to $\sim 0.25 \text{ nT}$ within $\sim 300 \text{ Myr}$ if AGN feedback ceases. UQFF predicts an associated 10% reduction in g_{BH} -anchored UQFF U_g at the cluster core — detectable as a measurable shift in the $H\alpha$ filament velocity dispersion from current $\sigma_v \sim 100 \text{ km/s}$ to $\sim 90 \text{ km/s}$ during a future AGN quiescent phase, accessible via VLT/MUSE integral-field spectroscopy.

Q5 Prediction 2: $F_0 = 0.1$, $\tau_{\text{fil}} = 100 \text{ Myr}$ predicts the cold filaments contribute a 10% gravitational enhancement to the UQFF U_g at $t = 0$, declining to 3.7% at 100 Myr. During the current active AGN phase, ALMA CO(2-1) kinematics should show a $\Delta v \approx 0.1 \times v_{U_g}$ excess velocity gradient from filament-gravity coupling — approximately $\sim 2 \text{ km/s}$ per kpc at 120 kpc distance, testable with sub-arcsecond ALMA maps.

Q5 Prediction 3: $T_{\text{cool}} = 4.76 \times 10^{-14} \text{ m/s}^2$ (cooling flow acceleration) predicts that the net inflow momentum flux of the cool ICM gas at 200 kpc is $F_{\text{cool}} = \rho_{\text{cool}} \times a_{\text{cool}} \times V_{\text{core}} \approx 10^{-20} \times 4.76 \times 10^{-14} \times (3 \times 10^{21})^3 \approx 1.3 \times 10^{28} \text{ N}$ — equivalent to a $\sim 10^{35} \text{ W}$ power input to the BCG gravitational field, consistent with the $\dot{M}_{\text{cool}} \approx 100 - 200 M_{\odot}/\text{yr}$ cooling rates derived from Hitomi/XRISM X-ray spectroscopy for Perseus A.