

# PAPER\_465 — NGC 1316 “Hubble Spies Cosmic Dust Bunnies”: MUGE UQFF Merger History, AGN Jets, Star Cluster Disruption

**Date:** 2025

**Whitepaper Series:** Star-Magic UQFF Phase 2 — Merger-Active Elliptical Galaxy  
**Session:** 120 (C++ module encoded) / Whitepapers created Session 122 **Source:** grok\_share\_dc707f5d3.txt (Doc 47 — NGC1316UQFFModule, “MUGE Hubble spies cosmic dust bunnies”) **Classification:** FIRST MUGE UQFF for NGC 1316; FIRST merger-tidal  $F_{\text{env}}$  term with spiral disruption; FIRST UQFF dust-fluid coupling for post-merger elliptical **Author:** Daniel T. Murphy **CP4 Class:** Pending (dc707f5d3 batch) **C++ Module:** NGC1316UQFFModule.h / NGC1316UQFFModule.cpp

---

## Abstract

This paper presents the complete NGC 1316 (Fornax A) gravitational evolution model under the Master Universal Gravity Equation (MUGE) + UQFF integration. NGC 1316 is a prominent post-merger elliptical galaxy in the Fornax Cluster with prominent radio lobes (Fornax A), AGN jets, and extensive dust-lane structure formed from cannibalizing a spiral companion. The UQFF model incorporates merger history via an evolving  $M_{\text{spiral}}(t)$  mass term, tidal cluster disruption ( $M_{\text{cluster}} = 1 \times 10^6 M_{\odot}$ ), dust-lane fluid coupling ( $\rho_{\text{dust}} = 1 \times 10^{-21} \text{ kg/m}^3$ ), AGN jet energy via Ug4 reactive decay, and dark matter halo. Result:  $g_{\text{NGC1316}} \approx 2 \times 10^{37} \text{ m/s}^2$  (DM/fluid dominant; repulsive terms advance framework).

---

## 2. Core Physics — PAPER\_465

### 2.1 System Parameters

| Parameter            | Value                                                                  | Notes                              |
|----------------------|------------------------------------------------------------------------|------------------------------------|
| M (total)            | $5 \times 10^{11} M_{\odot}$ ( $\sim 9.95 \times 10^{41} \text{ kg}$ ) | Post-merger mass                   |
| $M_{\text{spiral}}$  | $1 \times 10^{10} M_{\odot}$                                           | Cannibalized spiral companion      |
| $M_{\text{BH}}$      | $1 \times 10^8 M_{\odot}$                                              | Central AGN black hole             |
| $M_{\text{cluster}}$ | $1 \times 10^6 M_{\odot}$                                              | Star cluster undergoing disruption |
| r                    | 46 kpc ( $\sim 1.42 \times 10^{21} \text{ m}$ )                        | Effective radius                   |
| $d_{\text{spiral}}$  | 50 kpc                                                                 | Original separation from spiral    |
| $\rho_{\text{dust}}$ | $1 \times 10^{-21} \text{ kg/m}^3$                                     | Dust-lane fluid density            |
| B                    | $1 \times 10^{-4} \text{ T}$                                           | Enhanced AGN magnetic field        |
| z                    | 0.005                                                                  | Fornax Cluster redshift            |
| $M_{\text{DM}}$      | $\sim 0.85 \times M$                                                   | Dark matter fraction               |

### 2.2 Merger-Modified Gravitational Equation

$$g_{\text{NGC1316}}(r, t) = \frac{GM_{\text{sf}}(t)}{r(t)^2} (1 + H_z t) \left(1 - \frac{B}{B_{\text{crit}}}\right) (1 + F_{\text{env}}(t)) \\ + \sum U_{gi} + \frac{\Lambda c^2}{3} + U_i + g_{\text{quantum}} + g_{\text{fluid}} + g_{\text{DM}}$$

## 2.3 Merger-Driven $F_{\text{env}}(t)$

The tidal + cluster disruption environmental force:

$$F_{\text{tidal}} = \frac{GM_{\text{spiral}}}{d_{\text{spiral}}^2}$$

$$F_{\text{cluster}} = \frac{GM_{\text{cluster}}}{r_{\text{cluster}}^2}$$

$$F_{\text{env}}(t) = F_{\text{tidal}} + F_{\text{cluster}}$$

The cannibalized spiral contributes a persistent tidal debris field, while ongoing star cluster disruption deposits angular momentum into the dust lanes.

## 2.4 Merger Mass Evolution

$$M_{\text{merge}}(t) = M_{\text{spiral}}(1 - e^{-t/\tau_{\text{merge}}})$$

This term tracks the gradual gravitational capture and absorption of the spiral companion debris into the NGC 1316 potential well.

## 2.5 Ug Sub-terms

- **Ug1** (AGN magnetic dipole): Enhanced by  $B = 1 \times 10^{-4}$  T from radio lobe activity —  $10 \times$  typical galaxy field
- **Ug2** (superconductor boundary):  $U_{g2} = B_{\text{super}}^2 / (2\mu_0)$
- **Ug3'** (external spiral gravitational pull):  $U'_{g3} = GM_{\text{spiral}} / d_{\text{spiral}}^2$
- **Ug4** (AGN reactive energy):  $U_{g4} = k_4 \cdot E_{\text{react}}(t)$ ,  $E_{\text{react}} = 10^{46} e^{-0.0005t}$  — models jet energy deposition over cosmic timescales

## 2.6 Dust-Lane Fluid Term

$$g_{\text{fluid}} = \rho_{\text{dust}} \cdot V \cdot g_{\text{base}}$$

The dust-lane density ( $\rho_{\text{dust}} = 1 \times 10^{-21}$  kg/m<sup>3</sup>) is significantly lower than typical ISM gas, reflecting the post-merger dispersed dust structure visible in Hubble ACS images.

## 3. Equation Summary

$$g_{\text{NGC1316}}(r, t) = \frac{GM_{\text{sf}}(t)}{r(t)^2} (1 + H_z t) (1 - B/B_{\text{crit}}) (1 + F_{\text{tidal}} + F_{\text{cluster}}) + \sum U_{gi} + \frac{\Lambda c^2}{3} + U_i + g_{\text{quar}}$$

**Computed Result:**  $g_{\text{NGC1316}} \approx 2 \times 10^{37}$  m/s<sup>2</sup> — DM/fluid dominant; repulsive Ug2 and  $\Lambda$  terms provide merger equilibrium; AGN Ug4 advances the reactive decay framework.

## 4. Physical Interpretation

- **Post-merger AGN enhancement:**  $B = 1 \times 10^{-4}$  T (vs.  $B = 1 \times 10^{-5}$  T for normal spirals) directly amplifies U<sub>g1</sub> magnetic dipole term — modeling the Fornax A radio lobe power.
  - **Dust-lane dynamics:**  $\rho_{\text{dust}}$  term in the fluid equation captures the post-merger dust structure as a pseudo-fluid modifying the gravitational field profile.
  - **Cluster disruption:**  $F_{\text{cluster}}$  in  $F_{\text{env}}$  represents ongoing star cluster tidal stripping — a UQFF innovation modeling galactic cannibalism.
- 

## 5. C++ Module Reference

**Module:** NGC1316UQFFModule (root-level, Session 120 from grok\_share\_dc707f5d3.txt)

**Key method:** computeG(double t) — returns total  $g_{\text{NGC1316}}$  in  $\text{m/s}^2$

**Unique feature:** computeMmerge(double t) — merger mass evolution term **Integration point:** MAIN\_1\_CoAnQi.cpp multi-system validation

---

---

##  
§A.  
Cosmogenesis-  
Linked  
La-  
grangian  
(PA-  
PER\_877  
Sym-  
bolic  
Ex-  
port)  
##  
§B.  
VDS/DVP/BSH  
Deep  
Syn-  
the-  
sis  
###  
§B.1  
Vac-  
uum  
Den-  
sity  
Se-  
ries  
(VDS)

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)  
 The  
 canon-  
 ical  
 VDS  
 ratio  
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$   
 1.894  
 gov-  
 erns  
 the  
 double-  
 exponential  
 vac-  
 uum  
 con-  
 den-  
 sate  
 pro-  
 file:  
 $\rho_{\text{vac}}(r) =$   
 $\rho_{\text{vac},[\text{SCm}]}$   
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

---

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER\_877

Sym-

bolic

Ex-

port)

---

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.129

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

$\pi$

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io5

con-

nects

---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

###  
 §B.2  
 Dipole  
 Vor-  
 tex  
 Primes  
 (DVP)  
 The  
 DVP  
 en-  
 cod-  
 ing  
 maps  
 the  
 sys-  
 tem's  
 char-  
 ac-  
 teris-  
 tic  
 pa-  
 ram-  
 eter  
 onto  
 the  
 prime  
 lat-  
 tice:  
 $p_{\text{DVP}} =$   
 $53, n_{\text{channel}} =$   
 $24/26$

---

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER\_877

Sym-

bolic

Ex-

port)

---

Since

$p_{\text{DVP}} =$

53 is

**res-**

**o-**

**nant**

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

###  
 §B.3  
 Buoy-  
 ancy  
 Sat-  
 ura-  
 tion  
 Har-  
 mon-  
 ics  
 (BSH)  
 The  
 BSH  
 satu-  
 ra-  
 tion  
 timescale  
 for  
 this  
 sec-  
 tor  
 is  
**10<sup>4</sup>**  
**yr**  
 (spin-  
 down  
 equi-  
 lib-  
 rium):  
 $\mathcal{F}_{\text{BSH}} =$   
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$   
 $f_{U_b} \cdot$   
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$   
 $\cos\big(\frac{2\pi j}{26}\big)$



---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

The  
 tanh  
 satu-  
 ra-  
 tion  
 en-  
 ve-  
 lope  
 pre-  
 vents  
 un-  
 phys-  
 ical  
 di-  
 ver-  
 gence:  
 $\mathcal{F}_{\text{BSH,sat}} =$   
 $\mathcal{F}_{\text{BSH}}$   
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER\_877

Sym-

bolic

Ex-

port)

connecting

to

the

PA-

PER\_877

Stage

5

buoy-

ancy

seed

$U_{b,\text{seed}} =$

$0.1 \cdot$

$(\hbar c/r^2) \cdot$

$f_{\text{SCm}}$

which

ini-

tial-

izes

the

har-

monic

se-

ries

at

cos-

mo-

gen-

esis.

###

§B.4

Production-

Scale

Con-

sis-

tency

##  
§A.  
Cosmogenesis-  
Linked  
La-  
grangian  
(PA-  
PER\_877  
Sym-  
bolic  
Ex-  
port)

|  
Frame-  
work  
|  
Canon-  
ical  
Value

|  
This  
Pa-  
per |  
Sta-  
tus |  
|—

—  
|—  
—  
—|—  
—

—|—  
—||

VDS  
ratio  
|  
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$   
1.894

| Lo-  
cal  
sub-  
ratio

=  
0.129

| ✓  
Threshold-  
consistent

||  
DVP  
prime

|  
 $p_k \in$   
 $\{2,3,...,113\}$   
|

$p_{\text{DVP}} =$   
531  
✓

Res-

## §SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

| Observable                                                                 | UQFF Prediction                                                                                                                    | SM / Experiment                                                          | Source              | Alignment                                  |
|----------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------|---------------------|--------------------------------------------|
| Thomson $\sigma_T$<br>(QED<br>synchrotron)                                 | UQFF $U_m$<br>scattering kernel:<br>$\sigma_T = 6.6524 \times 10^{-29}$<br>$m^2$                                                   | $\sigma_T = 6.6524 \times 10^{-29}$<br>$m^2$ (PDG QED exact)             | PDG<br>2024         | 100% (exact<br>QED input)                  |
| Active<br>Galactic<br>Nucleus /<br>SMBH<br>luminosity<br>X-ray 2–10<br>keV | UQFF MUGE $g_{total}$<br>$\rightarrow L_X$ via<br>Stefan-Boltzmann +<br>buoyancy flux: $L_X$<br>$\approx g_{total} \times M_{env}$ | $L_X L_X \sim 10^{43} - 10^{46}$<br>erg/s                                | Chandra/XMM         | Consistent<br>order of<br>magnitude        |
| GR<br>Schwarzschild<br>limit                                               | UQFF $g_{total}$ must<br>satisfy $g \leq c^2/(2r_s)$<br>at event horizon                                                           | $r_s = 2GM/c^2$ (GR<br>exact)                                            | PDG<br>2024 /<br>GR | ✓ UQFF<br>respects GR<br>horizon           |
| $\kappa$ vacuum rate<br>vs X-ray<br>variability                            | UQFF $\kappa =$<br>0.0005/day $\rightarrow$<br>timescale $\tau_{UQFF} =$<br>2000 days                                              | Observed X-ray<br>variability $\tau_{obs}$<br>(instrument<br>monitoring) | Chandra/XMM         | Stable<br>UQFF<br>variability<br>timescale |

**New physics claim:** UQFF MUGE generates gravity enhancement factors ( $g_{total}/g_{Newt} > 1$ ) for Active Galactic Nucleus / SMBH through vacuum buoyancy coupling — a mechanism absent from GR+SM. The enhancement factor and X-ray luminosity are linked via the UQFF buoyancy flux, providing a testable prediction for future Chandra/XMM monitoring observations.

*Cite PAPER\_642 (UQFFSMParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.*

**QS=5** — Full UQFF integration: merger  $F_{env}$ , AGN Ug4, dust-fluid coupling, post-merger elliptical parameters. *Copyright — Daniel T. Murphy. Encoded Oct 10, 2025.*