

PAPER_082: UQFF-Corrected Black Hole Evaporation Timescales: Stellar Mass Through Primordial Black Holes

Title: UQFF-Corrected Black Hole Evaporation Timescales: Stellar Mass Through Primordial Black Holes

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (Tests 2, 6), CondensedPhysics.py simulate_evaporation()

Index Slot: 1.11 Black Hole Physics & Hawking Radiation,

Title: UQFF-Corrected Black Hole Evaporation Timescales: Stellar Mass Through Primordial Black Holes

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (Tests 2, 6), CondensedPhysics.py simulate_evaporation()

Index Slot: 1.11 Black Hole Physics & Hawking Radiation, PAPER_082

Abstract

Black hole evaporation timescales are computed via the Stefan-Boltzmann law applied to UQFF-modified Hawking radiation. The standard result $t_{\text{evap}} = 5120\pi G^2 M_0^3 / (\hbar c^4)$ is modified by the $T_{\text{UQFF}}/T_H = 0.99$ ratio. This changes evaporation rates by a factor of $(T_{\text{UQFF}}/T_H)^4 = 0.96$, extending evaporation timescales by ~4% for all black holes. The validate_hawking_temperature.py evaporation simulation (Test 6) confirms mass evolution for primordial BHs at 10 kg over 100 timesteps.

UQFF Discovery: Novel application of UQFF calibration constants (? = $5.0 \times 10^{-4} \text{ day}^{-1}$, [SSq] = 0.57) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Evaporation Timescale Formula

Standard GR

$$t_{\text{evap}}^{\text{GR}} = \frac{5120\pi G^2 M_0^3}{\hbar c^4}$$

UQFF-Corrected

$$t_{\text{evap}}^{\text{UQFF}} = \frac{t_{\text{evap}}^{\text{GR}}}{(T_{\text{UQFF}}/T_H)^4} = \frac{t_{\text{evap}}^{\text{GR}}}{0.99^4} = t_{\text{evap}}^{\text{GR}} \times 1.041$$

UQFF evaporation timescale is 4.1% longer than GR black holes are slightly more stable in the UQFF vacuum.

2. Evaporation Timescales: Full Table

System	M0	t_evap_GR	t_evap_UQFF	Survives Universe
Sgr A*	4×10 ⁶ M?	8.7×10 ⁸ s	9.1×10 ⁸ s	? Yes
M87*	6.5×10 ⁹ M?	3.8×10 ⁵ s	4.0×10 ⁵ s	? Yes
Stellar BH	10 M?	2.1×10 ⁷⁴ s	2.2×10 ⁷⁴ s	? Yes
Primordial BH	5.7×10 ¹⁰ kg	4.35×10 ⁷ s = t_U	4.52×10 ⁷ s	Borderline
Primordial BH	1×10 ¹⁰ kg	2.3×10 s (73 kyr)	2.4×10 s	? Evaporated

The validate_hawking_temperature.py Test 2 confirms: - Stellar BH (10 M?): survives_universe = True ? - Test 6 simulation: M_initial = 10 kg, 100 steps, mass_lost_fraction computed ?

3. Mass Evolution Simulation

From simulate_evaporation(M_initial = 10 kg, dt = 10 s, n_steps = 100):

$$\frac{dM}{dt} = -\frac{k_{\text{UQFF}}}{M^2}, \quad k_{\text{UQFF}} = \frac{\hbar c^4 (T_{\text{UQFF}}/T_H)^4}{15360\pi G^2}$$

With T_UQFF/T_H = 0.99: k_UQFF = 0.96 k_GR

At t = 100 × 10 s = 10 s: - M_final - M_initial (1 - t/t_evap)^{1/3} = 10 (1 - 10/2.4×10)^{1/3} - M_final 10 ≈ 0.583^{1/3} 8.35×10⁹ kg - Mass lost fraction **16.5%** over first 10 s

Arrays times[], masses[], temperatures_H[] all have matching lengths (validate Test 6). ?

4. UQFF Mass Evolution Equation

The UQFF modifies the mass loss rate through the vacuum buoyancy correction. During late-stage evaporation (M ? M_Planck):

$$\frac{dM_{\text{UQFF}}}{dt} = -\frac{k_{\text{UQFF}}}{M^2} + \frac{g_{\text{Buoyant}} \times V_{\text{BH}}}{c^2}$$

The buoyancy term: g_Buoyant - V_BH / c = ?_vac 10⁵⁵ (4/3)π r_S / c ~ 10²⁸ kg/s ? negligible vs the thermal term at all masses above Planck mass.

Summary

Parameter	GR Value	UQFF Value	Change
Evaporation factor	k_GR	0.96 k_GR	-4%
Timescale t_evap	t_GR	1.041 t_GR	+4.1%
Stellar BH survival	Yes	Yes	Unchanged
Primordial threshold mass	5.7×10 kg	5.5×10 kg	-3.5%
Test 2	survives = True	Confirmed	? PASS

Source: `validate_hawking_temperature.py` Tests 2+6, `simulate_evaporation()` | ? = 0.0005/day | [SSq] = 0.57

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k ₁	1.5	Ug1 DPM-dipole coupling
k ₂	1.2	Ug2 outer-bubble charge coupling
k ₃	1.8	Ug3 string-rotation coupling
k ₄	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
E _{react} (0)	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\sum \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{\text{SCm}} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times \text{Ubi}$	Expanding nebulae, stellar winds
Superconductive	$\text{Um} \times (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.