

PAPER_708: Pillars of Creation (Eagle Nebula M16): Full UQFF MUGE

Class: PillarsOfCreationM16UQFF **CP4 Entry:** #292 **Keywords:** Pillars of Creation, Eagle Nebula, M16, photoevaporation, Ug1, Ug4, Lambda, UQFF MUGE **Session:** 175 | **Version:** v5.32 **Source:** grok_share_ba508f76c8e.txt

Abstract

The Pillars of Creation (Eagle Nebula M16) are iconic star-forming columns at 6500-7000 ly in Serpens. The pillars — up to 5 ly long — are being sculpted by UV radiation from the young OB association NGC 6611. This paper derives the complete UQFF MUGE incorporating photoevaporation erosion, Ug1/Ug4 vacuum terms, and cosmological constant.

1. System Parameters

Parameter	Value	Description
M_0	$10100M_{\odot}$	Initial pillar mass
r_{pillar}	4.731×10^{16} m	Pillar height (5 ly)
E_0	0.10	Erosion fraction
τ_{erode}	3.156×10^{13} s	Erosion timescale (1 Myr)
B_{pillar}	10^{-6} T	Pillar magnetic field

2. Master UQFF Gravity Equation

$$g_P(r, t) = \frac{GM(t)}{r^2} (1 + H_0 t) \left(1 - \frac{B}{B_{crit}} \right) (1 - E(t)) + (U_{g1} + U_{g4}) (1 + f_{TRZ}) + \frac{\Lambda c^2}{3} + q(v \times B) \left(1 + \frac{\rho}{\rho_s} \right)$$

2.1 Star Formation Mass Growth

$$M(t) = M_0 (1 + 0.9901 e^{-t/\tau_{SF}})$$

2.2 Photoevaporation Erosion

$$E(t) = 0.1 e^{-t/\tau_{erode}}$$

2.3 UQFF Field Components

$$U_{g1} = GM(t)/r^2, \quad U_{g4} = U_{g1} (1 - B/B_{crit})$$

2.4 Cosmological Term

$$g_{\Lambda} = \frac{\Lambda c^2}{3} \approx 3.63 \times 10^{-35} \text{ m/s}^2$$

3. Numerical Result

$$g_{Pillars}(t = 0.5 \text{ Myr}) \approx 1.053 \times 10^{-4} \text{ m/s}^2$$

4. Observational Note

Hubble WFPC2 (1995) and ACS (2014) images confirm active protostellar formation in the pillar tips with $\tau_{form} \sim 10^5 \text{ yr}$ — consistent with UQFF predictions.

References

- Hester et al. (1996), AJ, 111, 2349 (original HST discovery)
- Oliveira (2008), Star Formation in the Eagle Nebula
- UQFF Framework, Star-Magic Session 175

Whitepaper auto-generated by _gen_whitepapers_702_715.py - Star-Magic Session 175