

PAPER TITLE: Sigma Inventory of cluster Closures under v5.78

Star-Magic UQFF
PAPER_NNNN / Session SESSION
DATE

Abstract

This paper aggregates N closures from `cluster description` and reports the sigma-deviation inventory under UQFF v5.78. Of N closures, $n_{exact} > areexact(CE), < n_1sigma > within1\sigma$ of the observational target, $n_2sigma > within2\sigma$, and $n_{open} > remainasOPEN_PREDICTIONSsentinels$.

1 Method

For each closure i with UQFF prediction P_i and observation $O_i \pm \sigma_i$,

$$z_i = \frac{P_i - O_i}{\sigma_i}, \quad \chi^2 = \sum_i z_i^2, \quad \text{dof} = N - 1. \quad (1)$$

A closure is *exact* (CE) if $|z_i| < 10^{-6}$ and the predicted form is algebraically derivable from the canonical constants without fitting.

2 Inventory

#	Closure label	Predicted	Observed	z_i	Tag
1	<code>closure₁ ></code>	<code>P_i</code>	<code>O_i</code>	<code>z_i</code>	<code>iCE / 1S / 2S / OPEN_i</code>
2	<code>closure₂ ></code>	<code>P_i</code>	<code>O_i</code>	<code>z_i</code>	<code>iCE / 1S / 2S / OPEN_i</code>

3 Aggregate Statistics

N closures	N
Exact (CE)	$n_{exact} >$
Within 1σ	$n_1sigma >$
Within 2σ	$n_2sigma >$
Beyond 2σ	$n_{outlier} >$
Open predictions (9999)	$n_{open} >$
χ^2 / dof	$\chi^2_i / \text{dof}_i = \text{ratio}_i$

4 Outlier Analysis

For every closure beyond 2σ , give a one-paragraph diagnostic: which canonical constant or equation snippet from the `.txt` source dumps reconciles the gap, and which prior PAPER- i hosts the calibration.

5 Cross-references

- Master ledger: `master_closures.csv` (rows tagged `<cluster>`).
- Tracker source: `_uqff_program.py`.
- Open-prediction sentinels: `LEDGER_REVIEW_QUEUE.csv` OPEN_PLACEHOLDER.

References

[1] DOI/arXiv per observational anchor.