

PAPER_562: Buoyancy-Stratified Factorial Geometry — Bohr-Sommerfeld Aether Quantization

Unified Quantum Field Framework (UQFF)

Daniel T. Murphy • UQFF Research Consortium • 2026

Source: PAPER_562_BSFG_BohrSommerfeld_Aether_Quantization.md

PAPER_562: Buoyancy-Stratified Factorial Geometry — Bohr-Sommerfeld Aether Quantization

Author: Daniel T. Murphy — Star Magic / UQFF Framework **Session:** 149 | **Source:** Composed from CP4 #150, #43 (Sessions 148, 107–110) **CP4 Class:** [BSFGBBohrSommerfeldAetherQuantizationCalculator](#) (#157) **Date:** 2026-03-27

***Context note:** CP4 #150 (PAPER_555) derived the BSFG geodesic equation $v^2_{\text{orbit}} = GM/r + r c^2 \epsilon'/2$, showing the Aether contributes a velocity correction to circular orbits. This paper applies the Bohr-Sommerfeld quantization condition $J = n\hbar$ to compute the fractional action correction $\delta J/J$, the quantum of Aether action $\hbar_{\text{eta}}$, and the BSFG crossover radius r_{cross} where Aether and Newtonian orbital effects are equal.*

§1 Abstract

Applying Bohr-Sommerfeld quantization to the BSFG effective potential:

$$U_{\text{BSFG}}(r) = -\frac{GM}{r} + \frac{\eta c^2 C_{\text{num}} \cos(\pi t_n)}{2r^3}$$

we derive the fractional orbital action correction:

$$\frac{\delta J}{J} \approx \frac{v^2_{\text{aether}}}{2v^2_{\text{newton}}} = \frac{r c^2 \epsilon'}{2GM}$$

The Aether contribution dominates for $r < r_{\text{cross}}$, where:

$$r_{\text{cross}} = \left(\frac{\eta c^2 |\cos(\pi t_n)| C_{\text{num}}}{GM} \right)^{1/2} \approx 0.36 \text{ AU} \quad (t_n = 0)$$

Inside 0.36 AU from the Sun, BSFG orbital corrections are not small perturbations — they fundamentally alter the orbit. The quantum of Aether action:

$$\hbar_{\eta} = \eta \cdot h_{\text{Planck}} = 10^{-22} \times 6.626 \times 10^{-34} = 6.63 \times 10^{-56} \text{ J} \cdot \text{m}^3 \cdot \text{s/J}$$

provides the coupling between the BSFG metric perturbation and quantum orbital action.

§2 BSFG Effective Potential

From the geodesic equation (CP4 #150):

$$\frac{d^2 r}{d\lambda^2} = -\Gamma^r_{00} \left(\frac{dt}{d\lambda} \right)^2 = \frac{GM}{r^2} + \frac{c^2 \epsilon'}{2r^3}$$

The total (Newtonian + Aether) orbital effective potential per unit mass:

$$U_{\text{BSFG}}(r) = -\frac{GM}{r} + \frac{\eta c^2 C_{\text{num}} \cos(\pi t_n)}{2r^3}$$

For circular orbits, setting $dU/dr = 0$ (neglecting centrifugal for the action correction):

$$v_{\text{orbit}}^2 = \frac{GM}{r} + \frac{r c^2 \epsilon'(r)}{2}$$

where $\epsilon'(r) = -3\eta \cos(\pi t_n) C_{\text{num}} / r^4$ from CP4 #149.

§3 Bohr-Sommerfeld Action Integral

Step 1. The classical action for a circular orbit of radius r (angular momentum sector):

$$J = m v_{\text{orbit}} r = m \sqrt{GM/r + r^2 c^2 \epsilon'/2} \cdot r^{1/2} \approx m \sqrt{GM/r} \left(1 + \frac{r c^2 \epsilon'}{4GM} \right)$$

Step 2. The Newtonian Bohr-Sommerfeld action $J_0 = m \sqrt{GM/r}$; the BSFG correction:

$$\frac{\Delta J}{J} \approx \frac{v_{\text{aether}}^2}{2v_{\text{newton}}^2} = \frac{r c^2 \epsilon'/2}{2(GM/r)} = \frac{r^2 c^2 \epsilon'}{4GM}$$

Step 3. Substituting $\epsilon' = -3\eta \cos(\pi t_n) C_{\text{num}} / r^4$:

$$\boxed{\frac{\Delta J}{J}} = \frac{-3\eta \cos(\pi t_n) c^2 C_{\text{num}}}{4GM r^2}$$

Step 4. Values:

Radius	$\Delta J/J$ (at $t_n = 0$)
$r = R_{\odot} = 6.96 \times 10^8 \text{ m}$	$\approx -4.5 \times 10^4$ (Aether completely dominates)
$r = r_{\text{cross}} \approx 5.4 \times 10^{10} \text{ m}$	-0.5 (equal contributions)
$r = 1 \text{ AU} = 1.496 \times 10^{11} \text{ m}$	≈ -0.10 (10% correction)
$r = 10 \text{ AU}$	$\approx -9.7 \times 10^{-4}$
$r = 100 \text{ AU}$	$\approx -9.7 \times 10^{-6}$

Note: The large $\Delta J/J$ at sub-AU scales does not mean the theory is unphysical — it means Keplerian perturbation theory breaks down there, and the full BSFG orbit must be solved numerically. The proplyd confinement from PAPER_550 ($r_q \approx 0.1 \text{ AU}$) occurs precisely in this strong-field regime.

§4 Crossover Radius

Step 5. The BSFG crossover radius r_{cross} where $|v_{\text{aether}}^2| = v_{\text{newton}}^2$:

$$\eta^2 |C_{\text{num}}| |\cos(\pi t_n)| / r^2 = GM \implies r_{\text{cross}} = \sqrt{\frac{\eta^2 |C_{\text{num}}| |\cos(\pi t_n)|}{GM}}$$

At $t_n = 0$:

$$r_{\text{cross}} = \sqrt{\frac{10^{-22} \times 9 \times 10^{16} \times 4.27 \times 10^{46}}{6.674 \times 10^{-11} \times 1.989 \times 10^{30}}} = \sqrt{\frac{3.843 \times 10^{41}}{1.327 \times 10^{20}}} \approx 5.38 \times 10^{10} \text{ m} \approx 0.360 \text{ AU}$$

The Solar System planets Mercury (0.39 AU) and beyond lie in the Newtonian-dominant regime. Venus and Mercury's perihelion corrections require the full BSFG geodesic numerics.

§5 Aether Quantum of Action

Step 6. The Planck constant h has units J·s. The Aether coupling η has units m^3/J . Their product:

$$h_{\eta} \equiv \eta \times h_{\text{Planck}} = 10^{-22} \times 6.626 \times 10^{-34} \text{ m}^3 \cdot \text{s}$$

This quantity h_{η} represents the **minimum BSFG perturbation on a quantum orbital action** — how much the Aether-metric coupling shifts one quantum of angular momentum $\hbar$ when evaluated over a volume η .

Step 7. BSFG shift in the Keplerian quantum number $n = J_{\text{spec}}/\hbar = \sqrt{GMr}/\hbar$:

$$\Delta n_{\rm BSFG} = \frac{\Delta J}{J} \cdot n_{\rm Kepler} = \frac{-3\eta \, c^2 \, C_{\rm num}}{4GMr^2} \cos(\pi t_n) \sqrt{GMr} \hbar$$

At $r = 1 \, \rm AU$: $n_{\rm Kepler} \approx 2.23 \times 10^{74}$ and $\Delta n \approx -2.2 \times 10^{73}$.

§6 Physical Meaning

The BSFG Bohr-Sommerfeld analysis reveals three distinct orbital regimes:

Region	Dominant term	Physical consequence
$r < r_{\rm cross} \approx 0.36 \, \rm AU$	Aether	Non-Keplerian: proplyd confinement, DVP resonances
$r \approx r_{\rm cross}$	Both equal	BSFG transition zone — orbit switching
$r > r_{\rm cross}$	Newtonian	Classical Keplerian + small BSFG correction $\sim r^{-2}$

The BSFG crossover radius $r_{\rm cross} \approx 0.36 \, \rm AU$ is located between Mercury and Venus — consistent with the known anomalous perihelion precession corrections needed in the inner Solar System.

§7 References

- CP4 #150 — [BSFGGeodesicMetricCompatibilityCalculator](#) — PAPER_555 ($v^2_{\rm orbit} = GM/r + rc^2\varepsilon'/2$)
- CP4 #43 — Aether coupling $\eta = 10^{-22}$, PAPER_392
- CP4 #149 — [BSFGRiemannCurvatureAetherMetricCalculator](#) — PAPER_554 ($\varepsilon'(r)$)
- CP4 #147 — [Um26DPolyQuantizationDPMConfinementCalculator](#) — PAPER_550 (proplyd r_q in BSFG strong-field zone)