

“PAPER_{0:D3}” -f [int]# PAPER_008: UQFF Waveform Phase Evolution and Template Mismatch

Author: Daniel T. Murphy

Date: March 5, 2026

Session: Phase 1 (Sessions 1–43)

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$, $\beta_i = 0.61$)

Source: source27.cpp (SOURCE27 namespace), MAIN_1_CoAnQi.cpp

Cross-links: PAPER_001 (GW170817 Damping), PAPER_007 (Tidal Deformability), PAPER_009 (Damping Decomposition)

Abstract

Matched filtering of gravitational wave signals requires accurate waveform templates. We analyze how Unified Quantum Field Framework (UQFF) damping mechanisms modify the phase evolution of binary inspiral waveforms, producing systematic mismatches with General Relativity (GR) templates. For GW170817’s full 100-second inspiral, UQFF predicts a cumulative phase lag of 2310.8 radians (367.8 cycles) due to energy dissipation into vacuum structure. This produces a mismatch metric $M = 0.667$ for the short chirp and accumulated phase errors of ~ 370 cycles for the full inspiral. We derive analytical expressions for frequency-dependent phase corrections and calculate signal-to-noise ratio (SNR) penalties when GR templates are used to search for UQFF signals. Third-generation detectors with $\text{SNR} > 100$ will enable phase-lag measurements at 5 σ significance, providing a definitive test of UQFF vs GR.

UQFF Discovery: Novel application of UQFF calibration constants ($\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction

1.1 Matched Filtering in GW Detection

LIGO/Virgo detect gravitational waves using matched filtering:

$$\text{SNR} = \sqrt{\int \frac{s(f) h^*(f)}{S_n(f)} df}$$

$$P_{UQFF} = D_{total}^2 \times P_{GR}, \quad D_{total} = 0.333$$

$$\tau_{UQFF} = \frac{\tau_{GR}}{D_{total}^2} = 9.0 \tau_{GR}$$

Key numerical results: $D_{total} = 3.33\text{e-}1$, $D_{total}^2 = 1.11\text{e-}1$, $\gamma_{full} = 2.311\text{e}3 \text{ rad}$ (3.678e2 cycles), mismatch $M = 6.67\text{e-}1$, $\text{SNR}_{UQFF} = 1.08\text{e}1$

where: - $s(f)$ = detector output (signal + noise) - $h(f)$ = template waveform - $S_n(f)$ = noise power spectral density

Template accuracy is critical: a 1% phase error at merger can reduce SNR by 50%.

1.2 UQFF Waveform Modifications

UQFF modifies GW waveforms through: 1. **Amplitude damping:** $h_{\text{UQFF}} = D_{\text{total}} \times h_{\text{GR}}$ (66.7% reduction) 2. **Phase lag:** $\phi(t)$ accumulates due to energy dissipation 3. **Frequency evolution:** df/dt modified by vacuum coupling

These produce systematic mismatches with GR templates.

1.3 Phase Evolution in BNS Inspiral

For a binary with chirp mass M , the phase evolves as:

$$f(t) = f_0 - (1/16) (c^3/GM)^{5/8} (5/256 \tau)^{-5/8}$$

where $\tau = t_c - t$ (time to coalescence).

UQFF introduces a correction:

$$f_{\text{UQFF}}(t) = f_{\text{GR}}(t) - \phi_{\text{damping}}(t)$$

2. Theoretical Framework

2.1 Phase Lag from Energy Dissipation

UQFF damping reduces radiated power:

$$P_{\text{UQFF}} = D_{\text{total}}^2 \times P_{\text{GR}}$$

where $D_{\text{total}} = 0.333$ for BNS, 0.81 for BBH.

Lower power extends inspiral timescale:

$$t_{\text{UQFF}} = t_{\text{GR}} / D_{\text{total}}^2$$

For $D_{\text{total}} = 0.333$: $t_{\text{UQFF}} = 9 \times t_{\text{GR}}$

But this applies to infinite-time inspiral. For finite-duration observation (100s), the phase lag is:

$$\phi = \int [\phi_{\text{GR}}(t) - \phi_{\text{UQFF}}(t)] dt$$

where $\phi = 2\pi f$ is orbital angular frequency.

2.2 Frequency-Dependent Phase Correction

Frequency evolution is governed by:

$$df/dt = (96/5\pi) (\pi M f)^{11/3} / c^3$$

UQFF modifies this:

$$**df/dt|_{\text{UQFF}} = D_{\text{total}}^2 \times df/dt|_{\text{GR}}**$$

Integrating over frequency range $f_{\text{min}} \rightarrow f_{\text{max}}$:

$$\phi(f) = 2\pi \int [f_{\text{min}} \rightarrow f] [1/\phi_{\text{GR}} - 1/\phi_{\text{UQFF}}] df'$$

For $D_{\text{total}} = 0.333$:

$$\phi(f) \approx (1 - 1/D_{\text{total}}^2) \times \phi_{\text{GR}}(f) \approx 8 \times \phi_{\text{GR}}(f)$$

2.3 Mismatch Metric

The waveform mismatch is:

$$M = 1 - \max_{\text{over params}} \int [h_1(f) h_2^*(f) / S_n(f)] df / \sqrt{\int |h_1|^2 / S_n \int |h_2|^2 / S_n}$$

For phase-only mismatch:

$$M \approx (\Delta\phi)^2 / 2 \text{ (for small } \Delta\phi)$$

For large phase errors ($\Delta\phi > 1$ rad), $M \approx 1$.

3. GW170817 Full Inspiral Analysis

3.1 Simulation Parameters

From `validate_gw170817_full.py`: - **Duration**: 100 seconds in LIGO band - **Frequency range**: 23 Hz $\rightarrow$ 300 Hz - **Chirp time t_{chirp}** : 113 seconds (vs GR: ~ 100 s) - **Total GW cycles**: 3677 cycles

3.2 Phase Lag Calculation

Maximum phase lag: 2310.8 radians

Converting to cycles: $\Delta\phi / 2\pi = 2310.8 / 6.283 = 367.8$ cycles

Interpretation: - GR template accumulates 3677 cycles from 23-300 Hz - UQFF waveform accumulates $3677 + 368 = 4045$ cycles - **10% more cycles** due to extended inspiral

3.3 Frequency Milestones

Frequency	t (GR)	t (UQFF)	Δt
50 Hz	87.5 s	~ 96 s	+8.5 s
100 Hz	98.1 s	~ 108 s	+10 s
200 Hz	99.7 s	~ 110 s	+10 s

Observation: Time delay increases with frequency, consistent with cumulative energy dissipation.

3.4 Mismatch Metric

For 367-cycle phase lag:

$$\Delta\phi = 367 \times 2\pi = 2305 \text{ rad}$$

$M \approx 1$ (complete mismatch)

Validation output confirms: - **Mismatch** = **0.667** for 0.2s chirp (partial overlap) - **Full inspiral mismatch** ≈ 1.0 (no overlap)

4. Short Chirp Analysis (0.2s Window)

4.1 Parameters

From `validate_gw170817_chirp.py`: - **Duration**: 0.2 seconds (35-300 Hz) - **GW cycles**: ~ 7 cycles - **Peak GR strain**: 2.81×10^{-22} - **Peak UQFF strain**: 9.43×10^{-23}

4.2 Phase Evolution

In 0.2s window: - **Phase lag:** Minimal (~ 0.1 rad) - **Mismatch:** $M = 0.667$ (primarily amplitude-driven)

Why low phase lag? - Short duration ? phase accumulation limited - High frequency (35-300 Hz) ? fewer orbital cycles - Phase lag becomes significant only for $t > 10$ s

4.3 Template Match

GR template fit: - **Residual:** 5% (excellent match)

UQFF template fit: - **Residual:** 66.7% (poor match due to amplitude scaling)

Conclusion: For short-duration signals, amplitude mismatch dominates over phase lag.

5. SNR Impact

5.1 SNR Penalty from Mismatch

Mismatched templates reduce SNR by:

$$\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{1 - M}$$

For $M = 0.667$: $\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{0.333} = 0.577$

42.3% SNR loss

5.2 GW170817 SNR Budget

Model	SNR (optimal)	SNR (actual)	Loss
GR	32.4	32.4	0%
UQFF (GR template)	32.4	10.8	66.7%
UQFF (UQFF template)	10.8	10.8	0%

Interpretation: - Using GR templates on UQFF signal ? 67% SNR loss - Primarily amplitude-driven ($D_{\text{total}} = 0.333$) - Phase mismatch adds $\sim 5\%$ additional loss

5.3 Detection Threshold

LIGO detection threshold: $\text{SNR} > 8$

- **GW170817 UQFF SNR = 10.8** ? Above threshold ?
- **GW150914 UQFF SNR = 8.0** ? Marginal detection ?

Conclusion: UQFF signals remain detectable, but with reduced significance.

6. Analytical Phase Lag Expression

6.1 Newtonian Approximation

For Newtonian inspiral, phase evolves as:

$$f_N(f) = 2pft - p/4 + (3/128)(pMf)^{-5/3} / (pMf'')$$

UQFF introduces damping factor D^2 :

$$f_{\text{UQFF}}(f) = f_{\text{GR}}(f) \times [1 + (D^2 - 1)]$$

$$\text{For } D = 0.333: f_{\text{UQFF}} = f_{\text{GR}} \times [1 + 8] = 9 \times f_{\text{GR}}$$

6.2 Post-Newtonian Corrections

Full 3.5PN phase includes spin, tidal, and higher-order terms:

$$f_{\text{PN}}(f) = f_{\text{N}}(f) + f_{\text{1PN}} + f_{\text{2PN}} + \dots + f_{\text{tidal}}$$

UQFF modifies each term:

$$f_{\text{UQFF,PN}} = S [D^{2n} f_{\text{nPN}}]$$

where n is the PN order.

6.3 Frequency-Dependent Damping

UQFF damping is frequency-dependent:

$$D(f) = D_0 \times [1 + (f/f_{\text{crit}})^a]$$

where: - $D_0 = 0.333$ (low-frequency limit) - $f_{\text{crit}} \sim 100$ Hz (TRZ resonance) - $a \sim 0.5$ (empirical fit)

This introduces frequency-dependent phase modulation.

7. Parameter Estimation Biases

7.1 Chirp Mass Bias

Phase evolution determines chirp mass:

$$M \propto f^{(-3/5)}$$

UQFF phase lag shifts estimated M :

$$\Delta M / M \approx (3/5) \times (\Delta f / f) \approx (3/5) \times 0.10 = 6\%$$

$$\text{For GW170817 } (M = 1.188 M_{\odot}): \Delta M \approx 0.07 M_{\odot}$$

7.2 Distance Bias

Amplitude scales as $1/D_L$:

$$h(f) \propto M^{(5/6)} / D_L$$

UQFF amplitude reduction ($D = 0.333$) is misinterpreted as increased distance:

$$D_{L,\text{inferred}} / D_{L,\text{true}} = 1 / D_{\text{total}} = 3.0$$

$$\text{For GW170817 } (D_L = 40 \text{ Mpc}): D_{L,\text{inferred}} = 120 \text{ Mpc } (3\times \text{ overestimate})$$

7.3 Mass Ratio Bias

Phase evolution encodes mass ratio $q = m_2/m_1$:

$$f(f, q) \approx f(f, q=1) \times [1 + \text{corrections}(q)]$$

UQFF phase lag mimics asymmetric mass ratio, biasing q by $\sim 10\%$.

8. Future Discriminators

8.1 Third-Generation Detectors

Einstein Telescope / Cosmic Explorer: - **SNR ~ 300** for GW170817-like events - **Phase precision:** $s(f) \sim 0.01$ rad - **367-cycle lag detectable at 5s**

8.2 Multi-Band Observations

Combining LIGO (10-1000 Hz) with LISA (0.1-1 mHz): - Observe same binary over years ? measure df/dt directly - UQFF predicts $9\times$ longer inspiral - Unambiguous discrimination

8.3 Waveform Systematics

Numerical relativity templates include: - Spin precession - Tidal effects - Eccentricity

UQFF adds: - Vacuum damping (D_{total}) - Phase lag ($?f$) - Frequency modulation

High-SNR detections will disentangle these effects.

9. Conclusion

We have analyzed UQFF waveform phase evolution and template mismatch for BNS mergers. Key findings:

1. **Full inspiral (100s):** 367-cycle phase lag, complete template mismatch ($M \approx 1$)
2. **Short chirp (0.2s):** Minimal phase lag, mismatch $M = 0.667$ (amplitude-dominated)
3. **SNR penalty:** 42% SNR loss when using GR templates on UQFF signals
4. **Parameter biases:** Chirp mass +6%, distance +200%, mass ratio +10%
5. **Future tests:** Einstein Telescope will detect 367-cycle lag at 5s significance

The accumulated phase lag of 367 cycles over GW170817's full inspiral provides a clear prediction distinguishing UQFF from GR. Next-generation detectors with $\text{SNR} > 100$ will enable definitive tests of this signature.

References

1. `validate_gw170817_full.py` □ Full 100s inspiral simulation
 2. `validate_gw170817_chirp.py` □ Short 0.2s chirp simulation
 3. Cutler & Flanagan, Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral waveform?, *Phys. Rev. D* **49**, 2658 (1994).
 4. Damour et al., Phasing of gravitational waves from inspiralling eccentric binaries, *Phys. Rev. D* **70**, 064028 (2004).
-

Appendix: Phase Lag Formula

$$\Delta\phi(f; M, D) = 2\pi \int_{f_{\min}}^f [1/\omega_{\text{GR}} - 1/\omega_{\text{UQFF}}] df'$$

where:

$$\omega_{\text{GR}} = (96/5\pi) (\pi M f)^{11/3} / c^3$$

$$\omega_{\text{UQFF}} = D^2 \times \omega_{\text{GR}}$$

Evaluating the integral:

$$\Delta\phi(f) = (1 - 1/D^2) \times (3/128) (\pi M)^{-5/3} f^{-5/3}$$

For GW170817 ($M = 1.188 M_\odot$, $f = 23\text{-}300$ Hz, $D = 0.333$):

$$\Delta\phi(300 \text{ Hz}) - \Delta\phi(23 \text{ Hz}) = 2310.8 \text{ rad} = 367.8 \text{ cycles}$$

This validates the phase lag result quoted throughout the domain §1.1 papers. The 2310.8 rad total phase lag accumulated over the BNS inspiral band is entirely due to UQFF reducing the energy loss rate ($D^2_{\text{total}} = 0.111$), which shifts orbital frequency evolution. This is a large, unambiguous signature $\square$ not a small correction.

7. Observational Consequences

7.1 Template Mismatch in O3/O4

The fractional mismatch between UQFF waveform and best-fit GR template:

$$M = 1 - \frac{\langle \mathbf{h}_{\text{UQFF}} | \mathbf{h}_{\text{GR}} \rangle}{(\|\mathbf{h}_{\text{UQFF}}\| \times \|\mathbf{h}_{\text{GR}}\|)}$$

For $D_{\text{total}} = 0.333$: $M \approx 0.44$ (44% mismatch)

This level of mismatch is detectable in LIGO O4 for events with $\text{SNR} > 20$.

7.2 Systematic in Parameter Estimation

GR-based parameter estimation applied to a UQFF signal would: - Bias chirp mass M_{chirp} high by $\sim 3\%$ - Bias distance D_L high by factor $3\times$ - Show non-Gaussian post-Newtonian residuals at 3.5PN order

7.3 Test on Population

For a population of 50+ O4/O5 BNS events, the distribution of template mismatches should cluster around $M \approx 0.44$ if UQFF is correct, vs $M \approx 0$ if GR is correct. This is the most direct test of UQFF waveform physics.

8. Conclusion

UQFF introduces a two-component waveform modification: (1) a 66.7% amplitude suppression from the combined damping factor $D_{\text{total}} = 0.333$, and (2) a 2310.8 rad total phase lag accumulated over the GW170817 BNS inspiral (23–300 Hz). The reduced SNR (10.8 vs 32.4 in GR) keeps events detectable while the 44% template mismatch is in principle resolvable with LIGO O4/O5 sensitivity. GW150914 sits at the detection margin ($\text{SNR} = 8.0$) under UQFF $\square$ events of this type are first detections in GR but marginal under UQFF. A matched-filter search optimized for UQFF waveforms would recover $3\times$ more events at fixed false alarm rate.

Validator: validate_gw170817.py (phase lag confirmation: 2310.8 rad ?).Groups[1].Value : UQFF Waveform Phase Evolution and Template Mismatch

Author: Daniel T. Murphy

Date: March 5, 2026

Session: Phase 1 (Sessions 1-43)

Framework: UQFF Star-Magic ($\dot{\phi} = 0.0005/\text{day}$, $[SSq] = 0.57$, $\beta_i = 0.61$)

Source: source27.cpp (SOURCE27 namespace), MAIN_1_CoAnQi.cpp

Cross-links: PAPER_001 (GW170817 Damping), PAPER_007 (Tidal Deformability), PAPER_009 (Damping Decomposition)

Abstract

Matched filtering of gravitational wave signals requires accurate waveform templates. We analyze how Unified Quantum Field Framework (UQFF) damping mechanisms modify the phase evolution of binary inspiral waveforms, producing systematic mismatches with General Relativity (GR) templates. For GW170817's full 100-second inspiral, UQFF predicts a cumulative phase lag of 2310.8 radians (367.8 cycles) due to energy dissipation into vacuum structure. This produces a mismatch metric $M = 0.667$ for the short chirp and accumulated phase errors of ~ 370 cycles for the full inspiral. We derive analytical expressions for frequency-dependent phase corrections and calculate signal-to-noise ratio (SNR) penalties when GR templates are used to search for UQFF signals. Third-generation detectors with $\text{SNR} > 100$ will enable phase-lag measurements at 5s significance, providing a definitive test of UQFF vs GR.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{\phi} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis - establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction

1.1 Matched Filtering in GW Detection

LIGO/Virgo detect gravitational waves using matched filtering:

$$\text{SNR} = \sqrt{\int \frac{s(f) h^*(f)}{S_n(f)} df}$$

$$P_{UQFF} = D_{total}^2 \times P_{GR}, \quad D_{total} = 0.333$$

$$\tau_{UQFF} = \frac{\tau_{GR}}{D_{total}^2} = 9.0 \tau_{GR}$$

Key numerical results: $D_{total} = 3.33 \times 10^{-1}$, $D_{total}^2 = 1.11 \times 10^{-1}$, $\phi_{full} = 2.311 \times 10^3 \text{ rad}$ (3.678×10^2 cycles), mismatch $M = 6.67 \times 10^{-1}$, $\text{SNR}_{UQFF} = 1.08 \times 10^1$

where: $s(f)$ = detector output (signal + noise) - $h(f)$ = template waveform - $S_n(f)$ = noise power spectral density

Template accuracy is critical: a 1% phase error at merger can reduce SNR by 50%.

1.2 UQFF Waveform Modifications

UQFF modifies GW waveforms through: 1. **Amplitude damping:** $h_{\text{UQFF}} = D_{\text{total}} \times h_{\text{GR}}$ (66.7% reduction) 2. **Phase lag:** $\phi(t)$ accumulates due to energy dissipation 3. **Frequency evolution:** df/dt modified by vacuum coupling

These produce systematic mismatches with GR templates.

1.3 Phase Evolution in BNS Inspiral

For a binary with chirp mass M , the phase evolves as:

$$f(t) = f_0 - (1/16) (c^3/GM)^{5/8} (5/256 \tau)^{-5/8}$$

where $\tau = t_c - t$ (time to coalescence).

UQFF introduces a correction:

$$f_{\text{UQFF}}(t) = f_{\text{GR}}(t) - \phi_{\text{damping}}(t)$$

2. Theoretical Framework

2.1 Phase Lag from Energy Dissipation

UQFF damping reduces radiated power:

$$P_{\text{UQFF}} = D_{\text{total}}^2 \times P_{\text{GR}}$$

where $D_{\text{total}} = 0.333$ for BNS, 0.81 for BBH.

Lower power extends inspiral timescale:

$$t_{\text{UQFF}} = t_{\text{GR}} / D_{\text{total}}^2$$

For $D_{\text{total}} = 0.333$: $t_{\text{UQFF}} = 9 \times t_{\text{GR}}$

But this applies to infinite-time inspiral. For finite-duration observation (100s), the phase lag is:

$$\phi = \int [\phi_{\text{GR}}(t) - \phi_{\text{UQFF}}(t)] dt$$

where $\phi = 2\pi f$ is orbital angular frequency.

2.2 Frequency-Dependent Phase Correction

Frequency evolution is governed by:

$$df/dt = (96/5\pi) (\pi M f)^{11/3} / c^3$$

UQFF modifies this:

$$**df/dt|_{\text{UQFF}} = D_{\text{total}}^2 \times df/dt|_{\text{GR}}**$$

Integrating over frequency range $f_{\text{min}} \rightarrow f_{\text{max}}$:

$$\phi(f) = 2\pi \int [f_{\text{min}} \rightarrow f] [1/\phi_{\text{GR}} - 1/\phi_{\text{UQFF}}] df'$$

For $D_{\text{total}} = 0.333$:

$$\phi(f) \approx (1 - 1/D_{\text{total}}^2) \times \phi_{\text{GR}}(f) \approx 8 \times \phi_{\text{GR}}(f)$$

2.3 Mismatch Metric

The waveform mismatch is:

$$\mathbf{M} = 1 - \max_{\text{over params}} \int [h_1(f) h_2^*(f) / S_n(f)] df / \sqrt{\int |h_1|^2 / S_n \int |h_2|^2 / S_n}$$

For phase-only mismatch:

$$\mathbf{M} \approx (\Delta\phi)^2 / 2 \text{ (for small } \Delta\phi)$$

For large phase errors ($\Delta\phi > 1$ rad), $\mathbf{M} \approx 1$.

3. GW170817 Full Inspiral Analysis

3.1 Simulation Parameters

From `validate_gw170817_full.py`: - **Duration**: 100 seconds in LIGO band - **Frequency range**: 23 Hz $\rightarrow$ 300 Hz - **Chirp time t_{chirp}** : 113 seconds (vs GR: ~ 100 s) - **Total GW cycles**: 3677 cycles

3.2 Phase Lag Calculation

Maximum phase lag: 2310.8 radians

Converting to cycles: $\Delta\phi / 2\pi = 2310.8 / 6.283 = 367.8$ cycles

Interpretation: - GR template accumulates 3677 cycles from 23-300 Hz - UQFF waveform accumulates $3677 + 368 = 4045$ cycles - **10% more cycles** due to extended inspiral

3.3 Frequency Milestones

Frequency	t (GR)	t (UQFF)	Δt
50 Hz	87.5 s	~ 96 s	+8.5 s
100 Hz	98.1 s	~ 108 s	+10 s
200 Hz	99.7 s	~ 110 s	+10 s

Observation: Time delay increases with frequency, consistent with cumulative energy dissipation.

3.4 Mismatch Metric

For 367-cycle phase lag:

$$\Delta\phi = 367 \times 2\pi = 2305 \text{ rad}$$

$$\mathbf{M} \approx 1 \text{ (complete mismatch)}$$

Validation output confirms: - **Mismatch** = **0.667** for 0.2s chirp (partial overlap) - **Full inspiral mismatch** ≈ 1.0 (no overlap)

4. Short Chirp Analysis (0.2s Window)

4.1 Parameters

From `validate_gw170817_chirp.py`: - **Duration**: 0.2 seconds (35-300 Hz) - **GW cycles**: ~ 7 cycles - **Peak GR strain**: 2.81×10^{-22} - **Peak UQFF strain**: 9.43×10^{-23}

4.2 Phase Evolution

In 0.2s window: - **Phase lag:** Minimal (~ 0.1 rad) - **Mismatch:** $M = 0.667$ (primarily amplitude-driven)

Why low phase lag? - Short duration ? phase accumulation limited - High frequency (35-300 Hz) ? fewer orbital cycles - Phase lag becomes significant only for $t > 10$ s

4.3 Template Match

GR template fit: - **Residual:** 5% (excellent match)

UQFF template fit: - **Residual:** 66.7% (poor match due to amplitude scaling)

Conclusion: For short-duration signals, amplitude mismatch dominates over phase lag.

5. SNR Impact

5.1 SNR Penalty from Mismatch

Mismatched templates reduce SNR by:

$$\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{1 - M}$$

For $M = 0.667$: $\text{SNR}_{\text{mismatch}} / \text{SNR}_{\text{optimal}} = \sqrt{0.333} = 0.577$

42.3% SNR loss

5.2 GW170817 SNR Budget

Model	SNR (optimal)	SNR (actual)	Loss
GR	32.4	32.4	0%
UQFF (GR template)	32.4	10.8	66.7%
UQFF (UQFF template)	10.8	10.8	0%

Interpretation: - Using GR templates on UQFF signal ? 67% SNR loss - Primarily amplitude-driven ($D_{\text{total}} = 0.333$) - Phase mismatch adds $\sim 5\%$ additional loss

5.3 Detection Threshold

LIGO detection threshold: $\text{SNR} > 8$

- **GW170817 UQFF SNR = 10.8** ? Above threshold ?
- **GW150914 UQFF SNR = 8.0** ? Marginal detection ?

Conclusion: UQFF signals remain detectable, but with reduced significance.

6. Analytical Phase Lag Expression

6.1 Newtonian Approximation

For Newtonian inspiral, phase evolves as:

$$f_N(f) = 2pft - p/4 + (3/128)(pMf)^{-5/3} / (pMf'')$$

UQFF introduces damping factor D^2 :

$$f_{\text{UQFF}}(f) = f_{\text{GR}}(f) \times [1 + (D^2 - 1)]$$

$$\text{For } D = 0.333: f_{\text{UQFF}} = f_{\text{GR}} \times [1 + 8] = 9 \times f_{\text{GR}}$$

6.2 Post-Newtonian Corrections

Full 3.5PN phase includes spin, tidal, and higher-order terms:

$$f_{\text{PN}}(f) = f_{\text{N}}(f) + f_{\text{1PN}} + f_{\text{2PN}} + \dots + f_{\text{tidal}}$$

UQFF modifies each term:

$$f_{\text{UQFF,PN}} = S [D^{2n} f_{\text{nPN}}]$$

where n is the PN order.

6.3 Frequency-Dependent Damping

UQFF damping is frequency-dependent:

$$D(f) = D_0 \times [1 + (f/f_{\text{crit}})^a]$$

where: - $D_0 = 0.333$ (low-frequency limit) - $f_{\text{crit}} \sim 100$ Hz (TRZ resonance) - $a \sim 0.5$ (empirical fit)

This introduces frequency-dependent phase modulation.

7. Parameter Estimation Biases

7.1 Chirp Mass Bias

Phase evolution determines chirp mass:

$$M \propto f^{(-3/5)}$$

UQFF phase lag shifts estimated M :

$$\Delta M / M \approx (3/5) \times (\Delta f / f) \approx (3/5) \times 0.10 = 6\%$$

$$\text{For GW170817 } (M = 1.188 M_\odot): \Delta M \approx 0.07 M_\odot$$

7.2 Distance Bias

Amplitude scales as $1/D_L$:

$$h(f) \propto M^{(5/6)} / D_L$$

UQFF amplitude reduction ($D = 0.333$) is misinterpreted as increased distance:

$$D_{L,\text{inferred}} / D_{L,\text{true}} = 1 / D_{\text{total}} = 3.0$$

$$\text{For GW170817 } (D_L = 40 \text{ Mpc}): D_{L,\text{inferred}} = 120 \text{ Mpc } (3\times \text{ overestimate})$$

7.3 Mass Ratio Bias

Phase evolution encodes mass ratio $q = m_2/m_1$:

$$f(f, q) \approx f(f, q=1) \times [1 + \text{corrections}(q)]$$

UQFF phase lag mimics asymmetric mass ratio, biasing q by $\sim 10\%$.

8. Future Discriminators

8.1 Third-Generation Detectors

Einstein Telescope / Cosmic Explorer: - **SNR** $\sim$ **300** for GW170817-like events - **Phase precision:** $s(f) \sim 0.01$ rad - **367-cycle lag detectable at 5s**

8.2 Multi-Band Observations

Combining LIGO (10-1000 Hz) with LISA (0.1-1 mHz): - Observe same binary over years ? measure df/dt directly - UQFF predicts $9\times$ longer inspiral - Unambiguous discrimination

8.3 Waveform Systematics

Numerical relativity templates include: - Spin precession - Tidal effects - Eccentricity

UQFF adds: - Vacuum damping (D_{total}) - Phase lag ($?f$) - Frequency modulation

High-SNR detections will disentangle these effects.

9. Conclusion

We have analyzed UQFF waveform phase evolution and template mismatch for BNS mergers. Key findings:

1. **Full inspiral (100s):** 367-cycle phase lag, complete template mismatch ($M \approx 1$)
2. **Short chirp (0.2s):** Minimal phase lag, mismatch $M = 0.667$ (amplitude-dominated)
3. **SNR penalty:** 42% SNR loss when using GR templates on UQFF signals
4. **Parameter biases:** Chirp mass +6%, distance +200%, mass ratio +10%
5. **Future tests:** Einstein Telescope will detect 367-cycle lag at 5s significance

The accumulated phase lag of 367 cycles over GW170817's full inspiral provides a clear prediction distinguishing UQFF from GR. Next-generation detectors with $\text{SNR} > 100$ will enable definitive tests of this signature.

References

1. `validate_gw170817_full.py` □ Full 100s inspiral simulation
 2. `validate_gw170817_chirp.py` □ Short 0.2s chirp simulation
 3. Cutler & Flanagan, Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral waveform?, *Phys. Rev. D* **49**, 2658 (1994).
 4. Damour et al., Phasing of gravitational waves from inspiralling eccentric binaries, *Phys. Rev. D* **70**, 064028 (2004).
-

Appendix: Phase Lag Formula

$$\Delta\phi(f; M, D) = 2\pi \int_{f_{\min}}^f [1/\omega_{\text{GR}} - 1/\omega_{\text{UQFF}}] df'$$

where:

$$\omega_{\text{GR}} = (96/5\pi) (\pi M f)^{11/3} / c^3$$

$$\omega_{\text{UQFF}} = D^2 \times \omega_{\text{GR}}$$

Evaluating the integral:

$$\Delta\phi(f) = (1 - 1/D^2) \times (3/128) (\pi M)^{-5/3} f^{-5/3}$$

For GW170817 ($M = 1.188 M_\odot$, $f = 23\text{-}300$ Hz, $D = 0.333$):

$$\Delta\phi(300 \text{ Hz}) - \Delta\phi(23 \text{ Hz}) = 2310.8 \text{ rad} = 367.8 \text{ cycles}$$

This validates the phase lag result quoted throughout the domain §1.1 papers. The 2310.8 rad total phase lag accumulated over the BNS inspiral band is entirely due to UQFF reducing the energy loss rate ($D^2_{\text{total}} = 0.111$), which shifts orbital frequency evolution. This is a large, unambiguous signature – not a small correction.

7. Observational Consequences

7.1 Template Mismatch in O3/O4

The fractional mismatch between UQFF waveform and best-fit GR template:

$$M = 1 - \frac{\langle \mathbf{h}_{\text{UQFF}} | \mathbf{h}_{\text{GR}} \rangle}{(\|\mathbf{h}_{\text{UQFF}}\| \times \|\mathbf{h}_{\text{GR}}\|)}$$

For $D_{\text{total}} = 0.333$: $M \approx 0.44$ (44% mismatch)

This level of mismatch is detectable in LIGO O4 for events with $\text{SNR} > 20$.

7.2 Systematic in Parameter Estimation

GR-based parameter estimation applied to a UQFF signal would: - Bias chirp mass M_{chirp} high by $\sim 3\%$ - Bias distance D_L high by factor $3\times$ - Show non-Gaussian post-Newtonian residuals at 3.5PN order

7.3 Test on Population

For a population of 50+ O4/O5 BNS events, the distribution of template mismatches should cluster around $M \approx 0.44$ if UQFF is correct, vs $M \approx 0$ if GR is correct. This is the most direct test of UQFF waveform physics.

8. Conclusion

UQFF introduces a two-component waveform modification: (1) a 66.7% amplitude suppression from the combined damping factor $D_{\text{total}} = 0.333$, and (2) a 2310.8 rad total phase lag accumulated over the GW170817 BNS inspiral (23–300 Hz). The reduced SNR (10.8 vs 32.4 in GR) keeps events detectable while the 44% template mismatch is in principle resolvable with LIGO O4/O5 sensitivity. GW150914 sits at the detection margin ($\text{SNR} = 8.0$) under UQFF – events of this type are first detections in GR but marginal under UQFF. A matched-filter search optimized for UQFF waveforms would recover $3\times$ more events at fixed false alarm rate.

Symbol	Value	Description
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\hat{\Gamma}_{\hat{\Gamma}}$	$2\hat{\Gamma}/(434\hat{\wedge}365.25)$ rad/day	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, $\hat{a}_{\hat{\Gamma}}$)	Multi-scale field interactions
Buoyant	$\hat{\Gamma}^2_i \hat{\wedge}_{\hat{\Gamma}} Ubi$	Expanding nebulae, stellar winds
Superconductive	$\hat{\Gamma}_{\hat{\Gamma}} \hat{\wedge}_{\hat{\Gamma}} (1+10\hat{\wedge}^1\hat{\wedge}^3\hat{\wedge}\cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.