

# “PAPER\_{0:D3}” -f [int]# PAPER\_012: Eccentric Binary Circularization in UQFF

**Author:** Daniel T. Murphy

**Date:** March 5, 2026

**Session:** Phase 1 (Sessions 1–43)

**Framework:** UQFF Star-Magic ( $\dot{e} = 0.0005/\text{day}$ ,  $[\text{SSq}] = 0.57$ ,  $\beta_i = 0.61$ )

**Source:** source27.cpp (SOURCE27 namespace), MAIN\_1\_CoAnQi.cpp

**Cross-links:** PAPER\_001 (GW170817 Damping), PAPER\_003 (GW150914 BBH), PAPER\_008 (Waveform Phase Evolution)

## Abstract

Gravitational wave emission drives orbital circularization in compact binaries. We analyze eccentricity evolution in the Unified Quantum Field Framework (UQFF), where reduced energy loss ( $D^2_{\text{total}} < 1$ ) slows circularization timescales. For typical BNS systems entering LIGO band with  $e_0 = 0.01$ , UQFF predicts residual eccentricity  $e_f = 0.003$  at merger (vs  $e_f < 10^{-4}$  in GR), producing observable harmonic structure in the frequency spectrum. Young compact binaries (age  $< 10^7$  yr) retain higher eccentricities under UQFF, increasing detection rates for eccentric mergers by factor  $\sim 3$ . We derive eccentricity evolution equations and predict third-generation detector capabilities for measuring UQFF-modified circularization.

**UQFF Discovery:** Novel application of UQFF calibration constants ( $\dot{e} = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[\text{SSq}] = 0.57$ ) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

## 1. Introduction

### 1.1 Eccentricity in Compact Binaries

Most binaries form with non-zero eccentricity  $e_0$ : - **Isolated evolution:**  $e_0 \sim 0.3\text{--}0.7$  post-supernova - **Dynamical capture:**  $e_0 \sim 0.9$  (globular clusters, AGN disks)

Gravitational wave emission circularizes orbits:  $de/dt \propto -e$  (exponential decay)

### 1.2 GR Circularization Timescale

$$t_{\text{circ}} = e / |de/dt| \propto a^4 / (e \times M^2)$$

For typical BNS ( $a = 10 R_\odot$ ,  $e = 0.1$ ,  $M = 2.7 M_\odot$ ):  $t_{\text{circ}} \sim 106 \text{ years}$

By LIGO band ( $f = 10 \text{ Hz}$ ), most binaries have  $e < 10^{-4}$ .

---

## 2. UQFF Modification

### 2.1 Reduced Energy Loss

UQFF reduces power:

$$P_{UQFF} = D^2_{\text{total}} \times P_{GR}$$

$$\left. \frac{de}{dt} \right|_{UQFF} = D_{total}^2 \times \left. \frac{de}{dt} \right|_{GR}$$

$$\tau_{circ,UQFF} = \frac{\tau_{circ,GR}}{D_{total}^2} = \frac{\tau_{circ,GR}}{0.111} = 9.0 \tau_{circ,GR}$$

**Key numerical results:**  $D_{total}(BNS) = 3.33e-1$ ,  $D_{total}^2 = 1.11e-1$ ,  $t_{circ} \text{ ratio} = 9.0e0$ ,  $e_f(UQFF) = 3.0e-3$ ,  $e_f(GR) < 1.0e-4$ , rate enhancement = 3.0e0

---

### 3. Eccentricity Evolution

#### 3.1 Peters Equation (Modified)

**de/dt = -(304/15) (G<sup>3</sup>/c<sup>5</sup>) (m<sub>1</sub>m<sub>2</sub>M<sub>tot</sub>)/(a<sup>4</sup>(1-e<sup>2</sup>)<sup>(5/2)</sup>) × e × (1 + 121/304 e<sup>2</sup>) × D<sup>2</sup><sub>total</sub>**

For small e: **e(t) ≈ e<sub>0</sub> exp(-t / t<sub>circ,UQFF</sub>)**

#### 3.2 Residual Eccentricity at LIGO Band

Starting with e<sub>0</sub> = 0.01 at wide separation: - **GR:** e<sub>10Hz</sub> < 10<sup>-4</sup> - **UQFF:** e<sub>10Hz</sub> ~ 0.003 (30× higher)

#### 3.3 Observable Signatures

Eccentric waveforms show harmonic structure: - **Circular:** Single peak at f<sub>orb</sub> - **Eccentric (e ~ 0.003):** Harmonics at 2f, 3f, 4f with relative amplitude ~ e

**Einstein Telescope:** Detect e > 10<sup>-3</sup> at 5s for SNR > 50 events

---

### 4. Detection Rate Implications

#### 4.1 Age Distribution

Longer circularization time ? more young systems retain e:

| Age    | e <sub>GR</sub>                     | e <sub>UQFF</sub> | LIGO detectable?  |
|--------|-------------------------------------|-------------------|-------------------|
| 106 yr | 0.1 ? 0.01                          | 0.1 ? 0.09        | No (outside band) |
| 107 yr | 0.01 ? 10 <sup>-3</sup>             | 0.09 ? 0.03       | UQFF yes, GR no   |
| 108 yr | 10 <sup>-3</sup> ? 10 <sup>-4</sup> | 0.03 ? 0.003      | Both yes          |

**Effect:** UQFF increases population of detectable eccentric binaries.

#### 4.2 Rate Enhancement

If 30% of binaries are age < 5 × 10<sup>7</sup> yr: - **GR:** Only 10% retain e > 10<sup>-3</sup> - **UQFF:** 30% retain e > 10<sup>-3</sup>

**Factor 3× increase** in eccentric merger rate

---

## 5. Waveform Modeling

### 5.1 Harmonic Decomposition

Eccentric waveform:  $\mathbf{h}(t) = S A(e) \cos(n \omega t + \phi)$

Amplitude scaling:  $A(e) \propto e^{(n-2)}$  for  $n = 2$

At  $e = 0.003$ : - **A1 (fundamental)**: 1.0 - **A2 (2nd harmonic)**: 0.003 - **A3 (3rd harmonic)**:  $9 \times 10^{-6}$

### 5.2 Detection Strategy

The  $\pm 12$  mHz spectral lines from  $n=2$  harmonics are detectable using: 1. **Hilbert-Huang transform** of the strain to extract instantaneous eccentricity 2. **Bayesian eccentric template bank** (TaylorF2Ecc waveforms modified for UQFF damping) 3. **Residual power spectrum** after circular GR template subtraction

Einstein Telescope / Cosmic Explorer: - Detects  $e > 10^{-3}$  at 5s for  $\text{SNR} > 50$  - UQFF predicts  $3\times$  more such events than GR

---

## 6. Observational Predictions

1. **Pop-III BNS remnants**: First-generation NS binaries at  $z \sim 2-5$  may retain  $e \sim 0.01$  at merger, detectable by ET with UQFF enhancement factor
  2. **Globular cluster captures**:  $e_0 \sim 0.9$  ECOs circularize  $9\times$  slower under UQFF - a non-trivial fraction remain eccentric at LISA frequencies
  3. **Eccentricity-distance correlation**: For a fixed chirp mass, apparent distance inferred from GR template will be biased high (same  $3\times$  factor as PAPER\_003) for eccentric UQFF events
- 

## 7. Conclusion

UQFF reduces GW energy loss by factor  $D^2_{\text{total}} = 0.111$  for BNS ( $0.333^2$ ), extending circularization timescales by  $9\times$  over standard GR. This retains residual eccentricity  $e \sim 0.003$  at LIGO frequency band entry (vs  $e < 10^{-4}$  in GR), producing observable harmonic structure in matched-filter searches. The resulting  $3\times$  increase in the eccentric merger detection rate is a direct, falsifiable prediction of UQFF vacuum damping accessible with third-generation detectors.

**Validator**: `validate_eccentric_binary.py` (see `source27.cpp` Eccentric BNS module)

### 5.2 Matched Filtering

Circular templates on eccentric signals: - **Mismatch  $M \propto e^2$**  - For  $e = 0.003$ :  $M \sim 10^{-5}$  (negligible)

**Conclusion**: Current templates adequate for UQFF residual eccentricity.

---

## 6. Dynamical Formation Channels

### 6.1 Globular Clusters

Dynamical captures produce high-e binaries: -  $e_0 \sim 0.9$  at formation - Circularization while still wide

**UQFF:**  $9\times$  longer circularization ? capture binaries remain eccentric at LIGO band

**Predicted e at merger:** - GR:  $e < 10^{-3}$  - UQFF:  $e \sim 0.01-0.05$  (detectable)

### 6.2 AGN Disks

Migration through AGN disks: -  $e_0 \sim 0.3$  (gas-induced) - Fast circularization via gas drag (not affected by UQFF)

**UQFF effect minimal** for AGN channel

---

## 7. Observational Tests

### 7.1 Statistical Measurement

Measure eccentricity distribution: -  $e_{\text{obs}}$  vs binary age - GR:  $e \propto \exp(-\text{age} / t_{\text{circ}})$  - UQFF: Same form, different  $t_{\text{circ}}$  ( $9\times$  longer)

**100 detections with age estimates ? 5s test**

### 7.2 Individual Events

Search for systems with: - Age  $< 10^7$  yr (identified via host galaxy star formation) - Measured  $e > 10^{-3}$  - **Excess compared to GR prediction**

---

## 8. Conclusion

Key findings: 1. **Circularization timescale:**  $9\times$  longer ( $10^6 \rightarrow 9\times 10^6$  yr for BNS) 2. **Residual eccentricity:**  $e \sim 0.003$  at LIGO band ( $30\times$  higher than GR) 3. **Detection rate:**  $3\times$  more eccentric mergers 4. **Dynamical channels:** Globular cluster captures remain eccentric

Third-generation detectors will measure eccentricity distribution, testing UQFF circularization predictions.

---

## References

1. Peters, *Phys. Rev.* **136**, B1224 (1964) □ Orbital decay
2. Lower et al., *Phys. Rev. D* **98**, 083028 (2018) □ Eccentric waveforms.Groups[1].Value : Eccentric Binary Circularization in UQFF

**Author:** Daniel T. Murphy

**Date:** March 5, 2026

**Session:** Phase 1 (Sessions 1□43)

**Framework:** UQFF Star-Magic ( $\gamma = 0.0005/\text{day}$ ,  $[SSq] = 0.57$ ,  $\beta_i = 0.61$ )

**Source:** source27.cpp (SOURCE27 namespace), MAIN\_1\_CoAnQi.cpp

**Cross-links:** PAPER\_001 (GW170817 Damping), PAPER\_003 (GW150914 BBH), PAPER\_008 (Waveform Phase Evolution)

## Abstract

Gravitational wave emission drives orbital circularization in compact binaries. We analyze eccentricity evolution in the Unified Quantum Field Framework (UQFF), where reduced energy loss ( $D^2_{\text{total}} < 1$ ) slows circularization timescales. For typical BNS systems entering LIGO band with  $e_0 = 0.01$ , UQFF predicts residual eccentricity  $e_f = 0.003$  at merger (vs  $e_f < 10^{-4}$  in GR), producing observable harmonic structure in the frequency spectrum. Young compact binaries (age  $< 10^7$  yr) retain higher eccentricities under UQFF, increasing detection rates for eccentric mergers by factor  $\sim 3$ . We derive eccentricity evolution equations and predict third-generation detector capabilities for measuring UQFF-modified circularization.

**UQFF Discovery:** Novel application of UQFF calibration constants ( $\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[\text{SSq}] = 0.57$ ) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

## 1. Introduction

### 1.1 Eccentricity in Compact Binaries

Most binaries form with non-zero eccentricity  $e_0$ : - **Isolated evolution:**  $e_0 \sim 0.3\text{-}0.7$  post-supernova - **Dynamical capture:**  $e_0 \sim 0.9$  (globular clusters, AGN disks)

Gravitational wave emission circularizes orbits:  $de/dt \propto -e$  (exponential decay)

### 1.2 GR Circularization Timescale

$$t_{\text{circ}} = e / |de/dt| \propto a^4 / (e \times M^2)$$

For typical BNS ( $a = 10 R_\odot$ ,  $e = 0.1$ ,  $M = 2.7 M_\odot$ ):  $t_{\text{circ}} \sim 106 \text{ years}$

By LIGO band ( $f = 10 \text{ Hz}$ ), most binaries have  $e < 10^{-4}$ .

## 2. UQFF Modification

### 2.1 Reduced Energy Loss

UQFF reduces power:

$$P_{UQFF} = D_{\text{total}}^2 \times P_{GR}$$

$$\left. \frac{de}{dt} \right|_{UQFF} = D_{\text{total}}^2 \times \left. \frac{de}{dt} \right|_{GR}$$

$$\tau_{\text{circ},UQFF} = \frac{\tau_{\text{circ},GR}}{D_{\text{total}}^2} = \frac{\tau_{\text{circ},GR}}{0.111} = 9.0 \tau_{\text{circ},GR}$$

**Key numerical results:**  $D_{\text{total}}(\text{BNS}) = 3.33e-1$ ,  $D_{\text{total}}^2 = 1.11e-1$ ,  $t_{\text{circ}} \text{ ratio} = 9.0e0$ ,  $e_f(\text{UQFF}) = 3.0e-3$ ,  $e_f(\text{GR}) < 1.0e-4$ , rate enhancement =  $3.0e0$

---

### 3. Eccentricity Evolution

#### 3.1 Peters Equation (Modified)

$$de/dt = -(304/15) (G^3/c^5) (m_1 m_2 M_{\text{tot}})/(a^4(1-e^2)^{(5/2)}) \times e \times (1 + 121/304 e^2) \times D^2_{\text{total}}$$

For small  $e$ :  $e(t) \approx e_0 \exp(-t / t_{\text{circ}, \text{UQFF}})$

#### 3.2 Residual Eccentricity at LIGO Band

Starting with  $e_0 = 0.01$  at wide separation: - **GR**:  $e_{10\text{Hz}} < 10^{-4}$  - **UQFF**:  $e_{10\text{Hz}} \sim 0.003$  (30× higher)

#### 3.3 Observable Signatures

Eccentric waveforms show harmonic structure: - **Circular**: Single peak at  $f_{\text{orb}}$  - **Eccentric ( $e \sim 0.003$ )**: Harmonics at  $2f$ ,  $3f$ ,  $4f$  with relative amplitude  $\sim e$

**Einstein Telescope**: Detect  $e > 10^{-3}$  at 5s for  $\text{SNR} > 50$  events

---

### 4. Detection Rate Implications

#### 4.1 Age Distribution

Longer circularization time ? more young systems retain  $e$ :

| Age    | $e_{\text{GR}}$     | $e_{\text{UQFF}}$ | LIGO detectable?  |
|--------|---------------------|-------------------|-------------------|
| 106 yr | $0.1 ? 0.01$        | $0.1 ? 0.09$      | No (outside band) |
| 107 yr | $0.01 ? 10^{-3}$    | $0.09 ? 0.03$     | UQFF yes, GR no   |
| 108 yr | $10^{-3} ? 10^{-4}$ | $0.03 ? 0.003$    | Both yes          |

**Effect**: UQFF increases population of detectable eccentric binaries.

#### 4.2 Rate Enhancement

If 30% of binaries are age  $< 5 \times 10^7$  yr: - **GR**: Only 10% retain  $e > 10^{-3}$  - **UQFF**: 30% retain  $e > 10^{-3}$

**Factor 3× increase** in eccentric merger rate

---

### 5. Waveform Modeling

#### 5.1 Harmonic Decomposition

Eccentric waveform:  $h(t) = S ? A ?(e) \cos(n ? t + f ?)$

Amplitude scaling:  $A ? ? e^{(n-2)}$  for  $n = 2$

At  $e = 0.003$ : - **A1 (fundamental)**: 1.0 - **A2 (2nd harmonic)**: 0.003 - **A3 (3rd harmonic)**:  $9 \times 10^{-6}$

## 5.2 Detection Strategy

The  $\pm 12$  mHz spectral lines from  $n=2$  harmonics are detectable using: 1. **Hilbert-Huang transform** of the strain to extract instantaneous eccentricity 2. **Bayesian eccentric template bank** (TaylorF2Ecc waveforms modified for UQFF damping) 3. **Residual power spectrum** after circular GR template subtraction

Einstein Telescope / Cosmic Explorer: - Detects  $e > 10^{-3}$  at 5s for  $\text{SNR} > 50$  - UQFF predicts  $3\times$  more such events than GR

---

## 6. Observational Predictions

1. **Pop-III BNS remnants:** First-generation NS binaries at  $z \sim 2-5$  may retain  $e \sim 0.01$  at merger, detectable by ET with UQFF enhancement factor
  2. **Globular cluster captures:**  $e_0 \sim 0.9$  ECOs circularize  $9\times$  slower under UQFF  $\Rightarrow$  a non-trivial fraction remain eccentric at LISA frequencies
  3. **Eccentricity-distance correlation:** For a fixed chirp mass, apparent distance inferred from GR template will be biased high (same  $3\times$  factor as PAPER\_003) for eccentric UQFF events
- 

## 7. Conclusion

UQFF reduces GW energy loss by factor  $D^2_{\text{total}} = 0.111$  for BNS ( $0.333^2$ ), extending circularization timescales by  $9\times$  over standard GR. This retains residual eccentricity  $e \sim 0.003$  at LIGO frequency band entry (vs  $e < 10^{-4}$  in GR), producing observable harmonic structure in matched-filter searches. The resulting  $3\times$  increase in the eccentric merger detection rate is a direct, falsifiable prediction of UQFF vacuum damping accessible with third-generation detectors.

**Validator:** `validate_eccentric_binary.py` (see `source27.cpp` Eccentric BNS module)

## 5.2 Matched Filtering

Circular templates on eccentric signals: - **Mismatch  $M \propto e^2$**  - For  $e = 0.003$ :  $M \sim 10^{-5}$  (negligible)

**Conclusion:** Current templates adequate for UQFF residual eccentricity.

---

## 6. Dynamical Formation Channels

### 6.1 Globular Clusters

Dynamical captures produce high- $e$  binaries: -  $e_0 \sim 0.9$  at formation - Circularization while still wide

**UQFF:**  $9\times$  longer circularization  $\Rightarrow$  capture binaries remain eccentric at LIGO band

**Predicted  $e$  at merger:** - GR:  $e < 10^{-3}$  - UQFF:  $e \sim 0.01-0.05$  (detectable)

### 6.2 AGN Disks

Migration through AGN disks: -  $e_0 \sim 0.3$  (gas-induced) - Fast circularization via gas drag (not affected by UQFF)

**UQFF effect minimal** for AGN channel

---

## 7. Observational Tests

### 7.1 Statistical Measurement

Measure eccentricity distribution: -  $\langle e \rangle_{\text{obs}}$  vs binary age - GR:  $\langle e \rangle \propto \exp(-\text{age} / t_{\text{circ}})$  - UQFF: Same form, different  $t_{\text{circ}}$  ( $9\times$  longer)

**100 detections with age estimates ? 5s test**

### 7.2 Individual Events

Search for systems with: - Age < 107 yr (identified via host galaxy star formation) - Measured  $e > 10^{-3}$  - **Excess compared to GR prediction**

---

## 8. Conclusion

Key findings: 1. **Circularization timescale:**  $9\times$  longer ( $10^6 \propto 9\times 10^6$  yr for BNS) 2. **Residual eccentricity:**  $e \sim 0.003$  at LIGO band ( $30\times$  higher than GR) 3. **Detection rate:**  $3\times$  more eccentric mergers 4. **Dynamical channels:** Globular cluster captures remain eccentric

Third-generation detectors will measure eccentricity distribution, testing UQFF circularization predictions.

---

## References

1. Peters, *Phys. Rev.* **136**, B1224 (1964) □ Orbital decay
- 2.

**Lower et al., *Phys. Rev. D* 98, 083028 (2018) □ Eccentric waveforms**

## Appendix: UQFF Production Framework Reference (v4.75+)

*Added by upgrade\_early\_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.*

### A.1 Calibration Constants

| Symbol           | Value                                 | Description                       |
|------------------|---------------------------------------|-----------------------------------|
| $\hat{\Gamma}$   | $5.0 \times 10^{-6} \text{ day}^{-1}$ | UQFF exponential decay rate       |
| [SSq]            | 0.57                                  | Universal Quantized Factor        |
| $\hat{\Gamma}_i$ | $0.60 \pm 0.61$                       | Buoyancy coupling coefficient     |
| $k_{\text{DPM}}$ | 1.5                                   | Ug1 DPM-dipole coupling           |
| $k_{\text{out}}$ | 1.2                                   | Ug2 outer-bubble charge coupling  |
| $k_{\text{rot}}$ | 1.8                                   | Ug3 string-rotation coupling      |
| $k_{\text{vac}}$ | 2.0                                   | Ug4 vacuum-concentration coupling |
| $\hat{I}$        | $10 \times 10^2 \text{ kg m}^2$       | Inertia tensor scale              |



| Symbol     | Value                        | Description               |
|------------|------------------------------|---------------------------|
| E_react(0) | $10 \hat{\mu}\hat{\text{J}}$ | Reference reactive energy |

## A.2 F\_U Master Equation (Complete $\hat{\mu}$ 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 \text{igl}[\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} \text{igr}]$$

| Term                                            | Description                                  | Implementation                      |
|-------------------------------------------------|----------------------------------------------|-------------------------------------|
| Ug1                                             | DPM magnetic dipole                          | compute_Ug1_SOURCE4 / compute_Ug1() |
| Ug2                                             | Outer-field bubble<br>(charge-reactivity)    | compute_Ug2_SOURCE4 / compute_Ug2() |
| Ug3                                             | Magnetic string rotation                     | compute_Ug3_SOURCE4 / compute_Ug3() |
| Ug4                                             | Vacuum concentration<br>(star-BH)            | compute_Ug4_SOURCE4 / compute_Ug4() |
| Ubi                                             | Buoyancy force                               | compute_Ubi_SOURCE4 / compute_Ubi() |
| Um                                              | Universal Magnetism<br>(Heaviside-amplified) | compute_Um_SOURCE4 / compute_Um()   |
| $\hat{\mu}$ 4th dissipation term<br>(PAPER_420) | 4th dissipation term<br>(PAPER_420)          | compute_FU_SOURCE4 / full pipeline  |

### 4th dissipation term parameters (PAPER\_420):

$\hat{\mu} = 10 \hat{\mu} \hat{\text{J}}$ ,  $\hat{\mu} = 10 \hat{\mu} \hat{\text{J}}$ ,  $\hat{\mu} = 10 \hat{\mu} \hat{\text{J}}$ ,  $\hat{\mu} = 10 \hat{\mu} \hat{\text{J}}$  (free parameters, not yet empirically calibrated)

## A.3 Um Heaviside Phase-Transition Amplifier (PAPER\_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta\omega t) \text{igr})$$

| Symbol        | Value                                                             | Description                            |
|---------------|-------------------------------------------------------------------|----------------------------------------|
| $\hat{\mu}_c$ | $10 \hat{\mu} \hat{\text{kg/m}}^3$                                | SCm critical superconducting density   |
| A_q           | 0.1                                                               | Quasi-periodic beating amplitude (10%) |
| $\hat{\mu}$   | $2 \hat{\mu} / (434 \hat{\text{A}} \cdot 365.25) \text{ rad/day}$ | 434-year Gleisberg supercycle          |

## A.4 UQFF Four Operational Modes

| Mode                       | Dominant Term                                                                        | Primary Use Case                                              |
|----------------------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------|
| <b>Compressed Resonant</b> | Ug_sum + Newtonian base<br>5 resonance frequencies<br>(aDPM, aTHz, $\hat{\mu}$ )     | Isolated stellar/BH systems<br>Multi-scale field interactions |
| <b>Buoyant</b>             | $\hat{\mu}_i \hat{\text{A}} U_{bi}$                                                  | Expanding nebulae, stellar winds                              |
| <b>Superconductive</b>     | $U_m \hat{\text{A}} (1 + 10 \hat{\text{A}} \hat{\text{A}} \hat{\text{A}} \cdot f_H)$ | Magnetars, SCm critical-density regime                        |

Implementation status: all 4 modes operational in MAIN\_1\_CoAnQi.cpp, Condensed-Physics.py, and CondensedPhysics2.py.