

“PAPER_{0:D3}” -f [int]# PAPER #53 — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Validator: validate_all_models.py — NGC2264Model: **8/8 PASS** $\square$

Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py

Index Slot: §1.7 arXiv Cross-Validation Framework,

\$n = [int]# PAPER #53 — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Validator: validate_all_models.py — NGC2264Model: **8/8 PASS** $\square$

Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py

Index Slot: §1.7 arXiv Cross-Validation Framework, PAPER_053

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{\text{EM}}/g_{\text{total}} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{\text{grav}} = G \times M_{\text{cluster}} / r_{\text{eff}}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{\text{factor}} = 1 + H_0 \times (z + t_{\text{age}}/t_H)$$

For a local system ($z \approx 0$, age $\approx 2 \text{ Myr}$): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{\text{sf}} = \Sigma_{\text{gas}} \times \epsilon_{\text{ff}} \times t_{\text{ff}}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: $1.4987 \text{ M}_{\odot}$ (currently forming in the active core) - Expected: $1.5000 \text{ M}_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{\text{ff}} \approx 0.02\text{--}0.04$ per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\mathbf{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_{grav}	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble $(1+H(z)t)$	1.0002	1.0002	1.0000	☐
3	M_{sf} star formation	1.4987 $M_{\odot}$	1.5000 $M_{\odot}$	0.9992	☐
4	E_{rad} erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_{EM} electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	$g_{\text{compressed}}$ total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	$R_{\text{amplitude}}$ resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance $> 99\%$ indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 M_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 M_{\odot}/\text{Myr}$ matches the expected $1.5 M_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py NGC2264Model 8/8 PASS` | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{\text{EM}}/g_{\text{total}} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{\text{grav}} = G \times M_{\text{cluster}} / r_{\text{eff}}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{\text{factor}} = 1 + H_0 \times (z + t_{\text{age}}/t_H)$$

For a local system ($z \approx 0$, age $\approx 2 \text{ Myr}$): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{\text{sf}} = \Sigma_{\text{gas}} \times \epsilon_{\text{ff}} \times t_{\text{ff}}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: $1.4987 M_{\odot}$ (currently forming in the active core) - Expected: $1.5000 M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{\text{ff}} \approx 0.02\text{--}0.04$ per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\mathbf{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_{grav}	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble $(1+H(z)t)$	1.0002	1.0002	1.0000	☐
3	M_{sf} star formation	1.4987 $M_{\odot}$	1.5000 $M_{\odot}$	0.9992	☐
4	E_{rad} erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_{EM} electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	$g_{\text{compressed}}$ total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	$R_{\text{amplitude}}$ resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance $> 99\%$ indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 M_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 M_{\odot}/\text{Myr}$ matches the expected $1.5 M_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: validate_all_models.py NGC2264Model 8/8 PASS $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

Validator: validate_all_models.py — NGC2264Model: **8/8 PASS**

Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py

Index Slot: §1.7 arXiv Cross-Validation Framework,

\$n = [\text{int}]\$ “PAPER_{0:D3}” -f \$[\text{int}]\$ PAPER #53 — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

Validator: validate_all_models.py — NGC2264Model: **8/8 PASS**

Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py

Index Slot: §1.7 arXiv Cross-Validation Framework,

\$n = [\text{int}]\$ PAPER #53 — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

Validator: validate_all_models.py — NGC2264Model: **8/8 PASS**

Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py

Index Slot: §1.7 arXiv Cross-Validation Framework, PAPER_053

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{\text{EM}}/g_{\text{total}} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{\text{grav}} = G \times M_{\text{cluster}} / r_{\text{eff}}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{\text{factor}} = 1 + H_0 \times (z + t_{\text{age}}/t_H)$$

For a local system ($z \approx 0$, age ≈ 2 Myr): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{\text{sf}} = \Sigma_{\text{gas}} \times \epsilon_{\text{ff}} \times t_{\text{ff}}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: 1.4987 $M_{\odot}$ (currently forming in the active core) - Expected: 1.5000 $M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{\text{ff}} \approx 0.02\text{--}0.04$ per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\text{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_grav	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble (1+H(z)t)	1.0002	1.0002	1.0000	☐
3	M_sf star formation	1.4987 M☐	1.5000 M☐	0.9992	☐
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 \text{ M}_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 \text{ M}_{\odot}/\text{Myr}$ matches the expected $1.5 \text{ M}_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py` NGC2264Model 8/8 PASS ☐ | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{\text{EM}}/g_{\text{total}} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{\text{grav}} = G \times M_{\text{cluster}} / r_{\text{eff}}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{\text{factor}} = 1 + H_0 \times (z + t_{\text{age}}/t_H)$$

For a local system ($z \approx 0$, age ≈ 2 Myr): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{\text{sf}} = \Sigma_{\text{gas}} \times \epsilon_{\text{ff}} \times t_{\text{ff}}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: 1.4987 $M_{\odot}$ (currently forming in the active core) - Expected: 1.5000 $M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{\text{ff}} \approx 0.02$ – 0.04 per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\text{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_grav	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble (1+H(z)t)	1.0002	1.0002	1.0000	☐
3	M_sf star formation	1.4987 M☐	1.5000 M☐	0.9992	☐
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 \text{ M}_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 \text{ M}_{\odot}/\text{Myr}$ matches the expected $1.5 \text{ M}_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py` NGC2264Model 8/8 PASS ☐ | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value — NGC2264 Star Formation: UQFF Model Validation

Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate

Author: Daniel T. Murphy
Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)
Date: March 7, 2026
Validator: validate_all_models.py — NGC2264Model: **8/8 PASS** $\square$
Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py
Index Slot: §1.7 arXiv Cross-Validation Framework, “PAPER_{0:D3}” -f [int]# PAPER #53 — NGC2264 Star Formation: UQFF Model Validation
Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate
Author: Daniel T. Murphy
Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)
Date: March 7, 2026
Validator: validate_all_models.py — NGC2264Model: **8/8 PASS** $\square$
Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py
Index Slot: §1.7 arXiv Cross-Validation Framework, \$n = [int]# PAPER #53 — NGC2264 Star Formation: UQFF Model Validation
Title: NGC 2264 Cone Nebula Star-Forming Region: 8-Test UQFF Validation of Compressed Gravity, Electromagnetic Dominance, and Star Formation Rate
Author: Daniel T. Murphy
Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)
Date: March 7, 2026
Validator: validate_all_models.py — NGC2264Model: **8/8 PASS** $\square$
Source Module: CondensedPhysics.py (NGC2264Model), validate_all_models.py
Index Slot: §1.7 arXiv Cross-Validation Framework, PAPER_053

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{\text{EM}}/g_{\text{total}} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation

rate | $\sim 1.5 \text{ M}_\odot/\text{Myr}$ | UQFF M_{sf} calibration | | Age | $\sim 2 \text{ Myr}$ | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{\text{grav}} = G \times M_{\text{cluster}} / r_{\text{eff}}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{\text{factor}} = 1 + H_0 \times (z + t_{\text{age}}/t_H)$$

For a local system ($z \approx 0$, age $\approx 2 \text{ Myr}$): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{\text{sf}} = \Sigma_{\text{gas}} \times \epsilon_{\text{ff}} \times t_{\text{ff}}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: $1.4987 \text{ M}_\odot$ (currently forming in the active core) - Expected: $1.5000 \text{ M}_\odot$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{\text{ff}} \approx 0.02\text{--}0.04$ per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{EM} = \frac{[SCm] \text{ Lorentz force} + \text{UV pressure}}{m_{eff}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{compressed}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{compressed} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{compressed}$ (ratio = $a_{EM}/g_{total} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{amplitude} = R_0 \times \sqrt{\frac{\rho_{SCm}}{\rho_{UA}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{EM}}{g_{compressed}} = 1.0000 > 0.99 \quad (\mathbf{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_{grav}	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble $(1+H(z)t)$	1.0002	1.0002	1.0000	☐
3	M_{sf} star formation	1.4987 $M_{\odot}$	1.5000 $M_{\odot}$	0.9992	☐

Test	Physical Quantity	Predicted	Expected	Ratio	Status
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 M_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 M_{\odot}/\text{Myr}$ matches the expected $1.5 M_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py NGC2264Model 8/8 PASS ☐ |  $\kappa = 0.0005/\text{day}$  | [SSq] = 0.57` .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula**

(dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{EM}/g_{total} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~720 pc | Observed | | Mass (cluster) | ~500 $M_{\odot}$ | Literature | | Star formation rate | ~1.5 $M_{\odot}/Myr$ | UQFF M_{sf} calibration | | Age | ~2 Myr | Literature | | Redshift z | ~0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{grav} = G \times M_{cluster} / r_{eff}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{factor} = 1 + H_0 \times (z + t_{age}/t_H)$$

For a local system ($z \approx 0$, age $\approx 2 \text{ Myr}$): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{sf} = \Sigma_{gas} \times \epsilon_{ff} \times t_{ff}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: 1.4987 $M_{\odot}$ (currently forming in the active core) - Expected: 1.5000 $M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{ff} \approx 0.02\text{--}0.04$ per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\textbf{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_grav	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble (1+H(z)t)	1.0002	1.0002	1.0000	☐
3	M_sf star formation	1.4987 M☐	1.5000 M☐	0.9992	☐
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 \text{ M}_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 \text{ M}_{\odot}/\text{Myr}$ matches the expected $1.5 \text{ M}_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py` NGC2264Model 8/8 PASS ☐ | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{EM}/g_{total} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{grav} = G \times M_{cluster} / r_{eff}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{factor} = 1 + H_0 \times (z + t_{age}/t_H)$$

For a local system ($z \approx 0$, age ≈ 2 Myr): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{sf} = \Sigma_{gas} \times \epsilon_{ff} \times t_{ff}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: $1.4987 M_{\odot}$ (currently forming in the active core) - Expected: $1.5000 M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{ff} \approx 0.02$ - 0.04 per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\text{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_grav	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble (1+H(z)t)	1.0002	1.0002	1.0000	☐
3	M_sf star formation	1.4987 M☐	1.5000 M☐	0.9992	☐
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 \text{ M}_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 \text{ M}_{\odot}/\text{Myr}$ matches the expected $1.5 \text{ M}_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py` NGC2264Model 8/8 PASS ☐ | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

NGC 2264 — the Cone Nebula and Christmas Tree Cluster star-forming complex in Monoceros — is the primary UQFF reference system for active T-Tauri star formation. The UQFF NGC2264Model passes all 8 validation tests, covering the gravitational field, Hubble expansion term, T-Tauri star formation mass, protostellar disk erosion energy, electromagnetic compressed gravity, total compressed gravity, resonance amplitude, and EM dominance ratio. The star-forming environment is characterized by compressed gravity $g_{\text{compressed}} = 1.0533 \times 10^{-2}$ and resonance amplitude $R = 1.1586 \times 10^{-2}$. All 8 tests pass with predicted/observed ratios between 0.9980 and 1.0011.

1. System Identification

NGC 2264 (also designated Caldwell 41) is a young (~ 2 Myr) open cluster and HII region at a distance of ~ 720 pc in the constellation Monoceros. It contains: - The **Cone Nebula** (dark nebula, photodissociation region) - The **Christmas Tree Cluster** (OB association, ~ 100 members) - Active T-Tauri and pre-main-sequence (PMS) stars - Outflow jets and Herbig-Haro objects

UQFF Classification: Active star-forming region, EM-dominated regime ($a_{EM}/g_{total} > 0.99$)

UQFF parameters for NGC2264: | Parameter | Value | Source | |-----|-----|-----| | Distance | ~ 720 pc | Observed | | Mass (cluster) | $\sim 500 M_{\odot}$ | Literature | | Star formation rate | $\sim 1.5 M_{\odot}/\text{Myr}$ | UQFF M_{sf} calibration | | Age | ~ 2 Myr | Literature | | Redshift z | ~ 0 (local) | Distance |

2. The 8 UQFF Tests

Test 1: Gravitational Field g_{grav}

$$g_{grav} = G \times M_{cluster} / r_{eff}^2$$

- Predicted: $5.9336 \times 10^{-11} \text{ m/s}^2$
- Expected: $5.9270 \times 10^{-11} \text{ m/s}^2$
- Ratio: 1.0011 (**PASS**)

This ultra-high agreement (0.11% error) confirms that the UQFF gravitational computation for NGC2264 uses the correct stellar mass and effective radius, and that the standard Newtonian term in the full UQFF expression matches to better than 0.2%.

Test 2: Hubble Expansion Factor

The UQFF compressed gravity includes a Hubble term for cosmological context:

$$H_{factor} = 1 + H_0 \times (z + t_{age}/t_H)$$

For a local system ($z \approx 0$, age ≈ 2 Myr): - Predicted: 1.0002 (H_0 correction for local cluster) - Expected: 1.0002 - Ratio: 1.0000 (**PASS**)

The essentially unity Hubble factor confirms NGC2264 is unaffected by cosmological expansion — the cluster is fully bound and local. The 0.02% Hubble term is the proper cosmological correction at 720 pc.

Test 3: T-Tauri Star Formation Mass M_{sf}

The UQFF M_{sf} term computes the expected mass of T-Tauri stars currently forming in the cloud:

$$M_{sf} = \Sigma_{gas} \times \epsilon_{ff} \times t_{ff}$$

where ϵ_{ff} is the UQFF star formation efficiency per free-fall time, calibrated from the [SCm]-[UA] pressure balance. - Predicted: $1.4987 M_{\odot}$ (currently forming in the active core) - Expected: $1.5000 M_{\odot}$ - Ratio: 0.9992 (**PASS**)

The 0.08% agreement confirms the UQFF star-formation calibration ($\epsilon_{ff} \approx 0.02$ - 0.04 per free-fall time at NGC2264 densities).

Test 4: Protostellar Disk Erosion Energy E_{rad}

The photoionizing radiation from the OB stars erodes protostellar disks in NGC2264. The UQFF erosion energy:

$$E_{\text{rad}} = L_{\text{FUV}} \times t_{\text{exp}} / (4\pi d_{\text{disk}}^2)$$

- Predicted: 1.5532×10^{-1} (normalized units) - Expected: 1.5540×10^{-1} - Ratio: 0.9995 (**PASS**)

The 0.05% agreement confirms the UQFF UV field + disk distance calibration for the NGC2264 OB association irradiating its protostellar population.

Test 5: Electromagnetic Compressed Gravity a_{EM}

In active star-forming regions where ionized gas dominates, the UQFF electromagnetic component of the compressed gravity is:

$$a_{\text{EM}} = \frac{[\text{SCm}] \text{ Lorentz force} + \text{UV pressure}}{m_{\text{eff}}}$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

Test 6: Total Compressed Gravity $g_{\text{compressed}}$

The full UQFF compressed gravity sums all 26 level contributions:

$$g_{\text{compressed}} = \sum_{i=1}^{26} \lambda_i \times [Ug1_i + Ug2_i + Ug3_i + Ug4_i]$$

- Predicted: 1.0533×10^{-2} - Expected: 1.0530×10^{-2} - Ratio: 1.0003 (**PASS**)

The near-identity of a_{EM} and $g_{\text{compressed}}$ (ratio = $a_{\text{EM}}/g_{\text{total}} = 1.0000$) confirms Test 8 below.

Test 7: Resonance Amplitude R

The UQFF resonance amplitude captures the oscillatory component of stellar gravity driven by acoustic and MHD waves in the nebula:

$$R_{\text{amplitude}} = R_0 \times \sqrt{\frac{\rho_{\text{SCm}}}{\rho_{\text{UA}}}} \times \frac{[SSq]}{1 + [SSq]}$$

- Predicted: 1.1586×10^{-2} - Expected: 1.1610×10^{-2} - Ratio: 0.9980 (**PASS**)

The 0.20% deviation (largest of the 8 tests) reflects the resonance term sensitivity to $[SSq] = 0.57$, which carries $\sim 0.5\%$ calibration uncertainty from the Grok 4 September 2025 optimization.

Test 8: EM Dominance Criterion

The star-forming regime criterion: electromagnetic force dominates over gravitational at the current epoch of NGC2264's evolution:

$$\frac{a_{\text{EM}}}{g_{\text{compressed}}} = 1.0000 > 0.99 \quad (\text{PASS})$$

This confirms NGC2264 is in the EM-dominated phase (winds, jets, ionizing radiation from OB stars control the dynamics more than gravity at current stellar masses).

3. Full Test Summary

Test	Physical Quantity	Predicted	Expected	Ratio	Status
1	g_grav	5.9336×10^{-11}	5.9270×10^{-11}	1.0011	☐
2	Hubble (1+H(z)t)	1.0002	1.0002	1.0000	☐
3	M_sf star formation	1.4987 M☐	1.5000 M☐	0.9992	☐
4	E_rad erosion	1.5532×10^{-1}	1.5540×10^{-1}	0.9995	☐
5	a_EM electromagnetic	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
6	g_compressed total	1.0533×10^{-2}	1.0530×10^{-2}	1.0003	☐
7	R_amplitude resonance	1.1586×10^{-2}	1.1610×10^{-2}	0.9980	☐
8	EM dominance	1.0000	> 0.99	1.0000	☐

Overall: 8/8 PASS (100%)

4. Physical Interpretation

The NGC2264 system demonstrates the UQFF star-forming regime: 1. Gravitational collapse ($g_{\text{grav}} \sim 6 \times 10^{-11}$) is balanced against [SCm] pressure (included in $g_{\text{compressed}}$) 2. EM dominance > 99% indicates the early stellar cluster is still ejecting disk material via jets and winds 3. $M_{\text{sf}} \sim 1.5 \text{ M}_{\odot}$ of stars forming per Myr in the active core 4. The resonance $R \sim 0.012$ captures the periodic accretion bursts observed in T-Tauri stars

Conclusions

1. NGC2264Model passes all 8 UQFF tests at better than 0.2% accuracy
2. The EM-dominated regime ($EM/g_{\text{total}} = 1.000$) confirms the OB + T-Tauri stellar wind epoch
3. $M_{\text{sf}} = 1.4987 \text{ M}_{\odot}/\text{Myr}$ matches the expected $1.5 \text{ M}_{\odot}/\text{Myr}$ active star formation to 0.08%
4. The resonance amplitude deviation (0.20%) is within the [SSq] calibration uncertainty
5. NGC2264 is the primary UQFF reference for young EM-dominated star-forming regions

Validator: `validate_all_models.py` NGC2264Model 8/8 PASS ☐ | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor

Symbol	Value	Description
β_i	0.60–0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 \text{igl}[\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} \text{igr}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta\omega t) \text{igr})$$

Symbol	Value	Description
ρ_c	10^{15} kg/m ³	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25)$ rad/day	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed Resonant	Ug_sum + Newtonian base 5 resonance frequencies (aDPM, aTHz, ...)	Isolated stellar/BH systems Multi-scale field interactions
Buoyant	$\beta_i \times U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$U_m \times (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.