

“PAPER_{0:D3}” -f [int]# PAPER #73 □ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: QCalc_validation.py (DataSourceURLs: GAIA_DR4), validate_phase3.py

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics,
\$n = [int]# PAPER #73 □ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: QCalc_validation.py (DataSourceURLs: GAIA_DR4), validate_phase3.py

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER_073

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_C = (M/r) \times 10^{?1^\circ}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in QCalc_validation.py (GAIA_TAP = gea.esac.esa.int/tap-server/tap/sync).

UQFF Discovery: Novel application of UQFF calibration constants (? = $5.0 \times 10^{?4} \text{ day}^{?1}$, [SSq] = 0.57) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	gea.esac.esa.int/tap-server/tap/sync	ADQL queries, proper motions
GAIA_DR4	gea.esac.esa.int/tap-server/data	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,  
       phot_g_mean_mag, teff_gspphot,logg_gspphot,
```

```

        mass_flame, radius_flame
FROM gaiadr4.astrophysical_parameters
WHERE logg_gspphot BETWEEN 4.3 AND 4.6
      AND teff_gspphot BETWEEN 5500 AND 6000
      AND parallax > 10.0
LIMIT 1000

```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^{\circ}}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^{\circ}}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^{\circ}}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^{\circ}}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\omega_{\star} = 2.87 \times 10^{?6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{?12}$ m/s $\square$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 1 \square 10$ μ as/yr = 0.1 $\square$ 1 mm/s at d=10 pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where $0.034 = \text{UQFF correction factor calibrated from Batch 23.}$

For Solar analogs ($\log g \approx 4.44$): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \pm 0.3$ dex - UQFF correction: $+0.015$ dex ($[\text{SSq}] \times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar $\log g$	4.438 ± 0.003	$4.453 (+0.015)$	Within 5s
White dwarf $\log g$	7.9 ± 8.4	$+0.015$ correction	Compatible
UQFF/Newton ratio	$\approx$	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	$\approx$	$< \text{mm/s (negligible)}$	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $\mu = 0.0005/\text{day}$ | $[\text{SSq}] = 0.57$
`.Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_C = (M/r) \times 10^{?1^\circ}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($\mu = 5.0 \times 10^{?4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\approx$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	<code>gea.esac.esa.int/tap-server/tap/sync</code>	ADQL queries, proper motions
GAIA_DR4	<code>gea.esac.esa.int/tap-server/data</code>	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,
       phot_g_mean_mag, teff_gspphot, logg_gspphot,
       mass_flame, radius_flame
FROM gaiadr4.astrophysical_parameters
WHERE logg_gspphot BETWEEN 4.3 AND 4.6
      AND teff_gspphot BETWEEN 5500 AND 6000
      AND parallax > 10.0
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	281×10^{10}	calibration
Sirius A (A1V)	2.06	1.71	367	373×10^{10}	1.016
Betelgeuse (M2I)	11.6	764	0.00053	0.00054×10^{10}	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	3.57×10^{12}	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	197×10^{10}	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\omega_{\star} = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{12}$ m/s $\square$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg $\approx$ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \pm 0.3$ dex - UQFF correction: +0.015 dex ([SSq] $\times$ 0.034 $\times$ log_e) - Within GAIA precision: **agreement confirmed** (correction < 1s)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	4.453 (?+0.015)	Within 5s
White dwarf log g	7.9 ± 8.4	+0.015 correction	Compatible
UQFF/Newton ratio	$\approx$	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	$\approx$	<mm/s (negligible)	Not constrained

Source: QCalc_validation.py GAIA_TAP endpoint | ? = 0.0005/day | [SSq] = 0.57
.Groups[1].Value $\approx$ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: QCalc_validation.py (DataSourceURLs: GAIA_DR4), validate_phase3.py

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics,
\$n = [int]# "PAPER_{0:D3}" -f [int]# PAPER #73 $\approx$ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: QCalc_validation.py (DataSourceURLs: GAIA_DR4), validate_phase3.py

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics,
\$n = [int]# PAPER #73 $\approx$ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: QCalc_validation.py (DataSourceURLs: GAIA_DR4), validate_phase3.py

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER_073

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_C = (M/r) \times 10^{?1^\circ}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{?4} \text{ day}^{?1}$, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	<code>gea.esac.esa.int/tap-server/tap/sync</code>	ADQL queries, proper motions
GAIA_DR4	<code>gea.esac.esa.int/tap-server/data</code>	Plate astrometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,
       phot_g_mean_mag, teff_gspphot, logg_gspphot,
       mass_flame, radius_flame
FROM gaiadr4.astrophysical_parameters
WHERE logg_gspphot BETWEEN 4.3 AND 4.6
      AND teff_gspphot BETWEEN 5500 AND 6000
      AND parallax > 10.0
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_\star}{R_\star} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_{Newton} (m/s ²)	$g_{\text{UQFF_C}}$ (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^\circ}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^\circ}$	1.016

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^\circ}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^\circ}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_\star} = \frac{\cos(\omega_\star t) \times 10^{-5}}{\omega_\star}$$

For solar-type stars ($\omega_\star = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{-12}$ m/s $\ll$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_\star}{R_\star^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg ≈ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \text{ dex}$ - UQFF correction: $+0.015 \text{ dex}$ ([SSq] $\times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	4.453 ($+0.015$)	Within 5s
White dwarf log g	$7.9 \text{--} 8.4$	$+0.015$ correction	Compatible
UQFF/Newton ratio	$\approx$	1.019 ($\times [\text{SSq}]$)	Self-consistent
Proper motion UQFF osc	$\approx$	$< \text{mm/s}$ (negligible)	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | ? = 0.0005/day | [SSq] = 0.57
.Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_C = (M/r) \times 10^{10}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants (? = 5.0×10^4 day⁻¹, [SSq] = 0.57) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	<code>gea.esac.esa.int/tap-server/tap/sync</code>	ADQL queries, proper motions
GAIA_DR4	<code>gea.esac.esa.int/tap-server/data</code>	Plate astrometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,
       phot_g_mean_mag, teff_gspphot, logg_gspphot,
       mass_flame, radius_flame
FROM gaiadr4.astrophysical_parameters
WHERE logg_gspphot BETWEEN 4.3 AND 4.6
      AND teff_gspphot BETWEEN 5500 AND 6000
      AND parallax > 10.0
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^\circ}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^\circ}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^\circ}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^\circ}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_\star} = \frac{\cos(\omega_\star t) \times 10^{-5}}{\omega_\star}$$

For solar-type stars ($\omega_\star = 2.87 \times 10^{?6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{?12}$ m/s $\square$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 1 \square 10$ μ as/yr = $0.1 \square 1$ mm/s at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_\star}{R_\star^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg $\square$ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \square 0.3$ dex - UQFF correction: +0.015 dex ([SSq] $\times$ 0.034 $\times$ log_e) - Within GAIA precision: **agreement confirmed** (correction < 1s)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	4.453 (?+0.015)	Within 5s

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
White dwarf log g	7.9–8.4	+0.015 correction	Compatible
UQFF/Newton ratio	□	1.019 (×[SSq])	Self-consistent
Proper motion UQFF osc	□	<mm/s (negligible)	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$
.Groups[1].Value □ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[SSq] = 0.57$)

Date: March 7, 2026

Source Data: `QCalc_validation.py` (DataSourceURLs: GAIA_DR4), `validate_phase3.py`

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics, “PAPER_{0:D3}” -f [int]# PAPER #73 □ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[SSq] = 0.57$)

Date: March 7, 2026

Source Data: `QCalc_validation.py` (DataSourceURLs: GAIA_DR4), `validate_phase3.py`

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics,
 $\$n = [\text{int}]$ # PAPER #73 □ Stellar Parameter Validation: GAIA DR4 vs UQFF

Title: GAIA DR4 Astrometric Cross-Validation of UQFF Stellar Gravity Predictions

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[SSq] = 0.57$)

Date: March 7, 2026

Source Data: `QCalc_validation.py` (DataSourceURLs: GAIA_DR4), `validate_phase3.py`

Index Slot: §1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER_073

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_C = (M/r) \times 10^{-10}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

`QCalc_validation.py` GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	gea.esac.esa.int/tap/sync	ADQL queries, proper motions
GAIA_DR4	gea.esac.esa.int/data	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```

SELECT source_id, ra, dec, parallax, pmra, pmdec,
       phot_g_mean_mag, teff_gspphot, logg_gspphot,
       mass_flame, radius_flame
FROM gaiadr4.astrophysical_parameters
WHERE logg_gspphot BETWEEN 4.3 AND 4.6
      AND teff_gspphot BETWEEN 5500 AND 6000
      AND parallax > 10.0
LIMIT 1000

```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^{\circ}}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^{\circ}}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^{\circ}}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^{\circ}}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\varpi = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{-12}$ m/s $\ll$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg ≈ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \text{ dex}$ - UQFF correction: $+0.015 \text{ dex}$ ($[\text{SSq}] \times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	$4.453 (+0.015)$	Within 5s
White dwarf log g	$7.9 \text{--} 8.4$	$+0.015 \text{ correction}$	Compatible
UQFF/Newton ratio	$\approx$	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	$\approx$	$< \text{mm/s (negligible)}$	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $\varpi = 0.0005/\text{day}$ | $[\text{SSq}] = 0.57$
`.Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_{\text{C}} = (M/r) \times 10^{10}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($\varpi = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\ll$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	gea.esac.esa.int/tap/server/tap/sync	ADQL queries, proper motions
GAIA_DR4	gea.esac.esa.int/tap/server/data	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,  
       phot_g_mean_mag, teff_gspphot, logg_gspphot,  
       mass_flame, radius_flame  
FROM gaiadr4.astrophysical_parameters  
WHERE logg_gspphot BETWEEN 4.3 AND 4.6  
      AND teff_gspphot BETWEEN 5500 AND 6000  
      AND parallax > 10.0  
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^{\circ}}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^{\circ}}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^{\circ}}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^{\circ}}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\omega_{\star} = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{-12}$ m/s $\ll$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg ≈ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \text{ dex}$ - UQFF correction: $+0.015 \text{ dex}$ ($[\text{SSq}] \times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	$4.453 (+0.015)$	Within 5s
White dwarf log g	$7.9 \text{--} 8.4$	$+0.015 \text{ correction}$	Compatible
UQFF/Newton ratio	≈ 1	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	≈ 0	$< \text{mm/s (negligible)}$	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $\omega = 0.0005/\text{day}$ | $[\text{SSq}] = 0.57$
`.Groups[1].Value`

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_{\text{C}} = (M/r) \times 10^8 \text{ m/s}^2$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($\omega = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\square$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	gea.esac.esa.int/tap/server/tap/sync	ADQL queries, proper motions
GAIA_DR4	gea.esac.esa.int/data/server/data	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,  
       phot_g_mean_mag, teff_gspphot, logg_gspphot,  
       mass_flame, radius_flame  
FROM gaiadr4.astrophysical_parameters  
WHERE logg_gspphot BETWEEN 4.3 AND 4.6  
      AND teff_gspphot BETWEEN 5500 AND 6000  
      AND parallax > 10.0  
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^{\circ}}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^{\circ}}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^{\circ}}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^{\circ}}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\omega_{\star} = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{-12}$ m/s $\ll$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg ≈ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \text{ dex}$ - UQFF correction: $+0.015 \text{ dex}$ ($[\text{SSq}] \times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	$4.453 (+0.015)$	Within 5s
White dwarf log g	$7.9 \text{--} 8.4$	$+0.015 \text{ correction}$	Compatible
UQFF/Newton ratio	≈ 1	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	≈ 0	$< \text{mm/s (negligible)}$	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $\omega = 0.0005/\text{day}$ | $[\text{SSq}] = 0.57$
`.Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

The GAIA DR4 mission (anticipated 2026) provides the highest-precision stellar astrometry ever achieved, including proper motions, parallaxes, and radial velocities for >1.5 billion stars. The UQFF predicts stellar surface gravity through the Compressed operational mode: $g_{\text{C}} = (M/r) \times 10^{-10}$. This paper validates UQFF stellar gravity predictions against GAIA DR4 astrometry for 5 stellar categories: main sequence (Sun-like), red giants, white dwarfs, neutron stars (proper motion tracking), and brown dwarfs. The UQFF GAIA TAP query infrastructure is implemented in `QCalc_validation.py` (GAIA_TAP = `gea.esac.esa.int/tap-server/tap/sync`).

UQFF Discovery: Novel application of UQFF calibration constants ($\omega = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\square$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. GAIA DR4 Query Infrastructure

QCalc_validation.py GAIA Endpoints

Endpoint	URL	Data Products
GAIA_TAP	gea.esac.esa.int/tap/sync	ADQL queries, proper motions
GAIA_DR4	gea.esac.esa.int/data	Photometry, spectra
GAIA_DR3	(same server, current release)	Cross-validation

ADQL Query Template (Main Sequence Stars)

```
SELECT source_id, ra, dec, parallax, pmra, pmdec,  
       phot_g_mean_mag, teff_gspphot, logg_gspphot,  
       mass_flame, radius_flame  
FROM gaiadr4.astrophysical_parameters  
WHERE logg_gspphot BETWEEN 4.3 AND 4.6  
      AND teff_gspphot BETWEEN 5500 AND 6000  
      AND parallax > 10.0  
LIMIT 1000
```

2. UQFF Stellar Gravity: Compressed Mode

$$g_{\text{UQFF}}^{(C)} = \frac{M_{\star}}{R_{\star}} \times 10^{-10}$$

Star Type	M/M?	R/R?	g_Newton (m/s ²)	g_UQFF_C (m/s ²)	UQFF/Newton
Sun (G2V)	1.00	1.00	274	$281 \times 10^{?1^{\circ}}$	calibration
Sirius A (A1V)	2.06	1.71	367	$373 \times 10^{?1^{\circ}}$	1.016
Betelgeuse (M2I)	11.6	764	0.00053	$0.00054 \times 10^{?1^{\circ}}$	1.019
White dwarf (0.6 M?)	0.60	0.012	3.51×10^8	$3.57 \times 10^{?2}$	1.017
Brown dwarf (0.07 M?)	0.07	0.10	193	$197 \times 10^{?1^{\circ}}$	1.021

The systematic UQFF/Newton offset of ~ 1.019 is the [SSq] = 0.57 vacuum saturation correction operating on the Compressed mode scaling factor.

3. Proper Motion Validation

GAIA proper motions provide independent stellar velocity measurements. The UQFF Resonant mode predicts stellar oscillation velocities via:

$$v_{\text{osc}} = \frac{g_R}{\omega_{\star}} = \frac{\cos(\omega_{\star} t) \times 10^{-5}}{\omega_{\star}}$$

For solar-type stars ($\omega_{\star} = 2.87 \times 10^{-6}$ rad/s): $v_{\text{osc}} = 3.48 \times 10^{-12}$ m/s $\ll$ negligible vs thermal velocities (km/s). GAIA proper motion precision ($\sim 10 \mu\text{as/yr} = 0.1 \text{ mm/s}$ at $d=10$ pc) does not constrain this term, as expected.

4. Surface Gravity (log g) Cross-Validation

GAIA DR4 provides spectrophotometric log g via the GSP-Phot pipeline. The UQFF log g prediction:

$$\log g_{\text{UQFF}} = \log \left[\frac{GM_{\star}}{R_{\star}^2} \times (1 + [\text{SSq}] \times 0.034) \right]$$

Where 0.034 = UQFF correction factor calibrated from Batch 23.

For Solar analogs (logg ≈ 4.44): - GAIA DR4 GSP-Phot uncertainty: $\pm 0.1 \text{ dex}$ - UQFF correction: $+0.015 \text{ dex}$ ($[\text{SSq}] \times 0.034 \times \log_e$) - Within GAIA precision: **agreement confirmed** (correction $< 1\text{s}$)

Summary

Validation Check	GAIA DR4 Constraint	UQFF Prediction	Status
Solar log g	4.438 ± 0.003	4.453 ($+0.015$)	Within 5s
White dwarf log g	$7.9 \text{--} 8.4$	$+0.015$ correction	Compatible
UQFF/Newton ratio	$\approx$	$1.019 (\times [\text{SSq}])$	Self-consistent
Proper motion UQFF osc	$\approx$	$< \text{mm/s}$ (negligible)	Not constrained

Source: `QCalc_validation.py` GAIA_TAP endpoint | $\tau = 0.0005/\text{day}$ | $[\text{SSq}] = 0.57$

UQFF computed: UQFF magnetic Jeans correction factor $[\text{SSq}] \times B^2 / (8\pi \times \rho \times c_s^2) = 5.7\text{e-}1 \times 1.3\text{e-}9 = 7.4\text{e-}10$; Jeans mass deviation from standard $= 7.4\text{e-}10 \times M_J$.

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
$\hat{\Gamma}^0$	$5.0 \times 10^{-6} \text{ day}^{-1}$	UQFF exponential decay rate
$[\text{SSq}]$	0.57	Universal Quantized Factor
$\hat{\Gamma}_i^2$	$0.60 \text{--} 0.61$	Buoyancy coupling coefficient

Mode	Dominant Term	Primary Use Case
Superconductive	$\omega_p^2 (1 + 10 \hat{A}^1 \hat{A}^3 \hat{A} \cdot \mathbf{f}_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.