

PAPER_374 — J1610+1811 Relativistic Quasar Jet UQFF-NS Coupling

Star Magic UQFF Whitepaper Series

Author: Daniel T. Murphy | **Session 101** | **Source:** grok_share_11254865.txt (lines 5100-5200)

Distinct from PAPER_360 (FU/Bi at z=6.5 — same object, different physics)

Abstract

$$F_{U,Bi} = \kappa \cdot \frac{\rho_{SCm}}{\rho_{UA}} \cdot (U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_m + U_{bi})$$

This paper presents the coupling of UQFF resonance gravity (from the 12-term MUGE model, PAPER_371) into a Navier-Stokes quasar jet simulation constrained by the relativistic parameters of the high-redshift quasar J1610+1811. While PAPER_360 computed FU and Bi properties for J1610+1811 at z=6.5, this paper addresses the same object at z=3.122 with a physically motivated relativistic jet velocity $v_{SCm} = 0.99c$ derived from observed jet power and luminosity. This represents the first NS-UQFF coupling simulation driven by actual high-redshift quasar observational data.

1. Observational Data (J1610+1811)

Parameter	Symbol	Value	Source
Redshift	z	3.122	Optical/spectroscopic
Jet power	P_{jet}	$\sim 4 \times 10^{45}$ W	X-ray observation
Total luminosity	L	$\sim 2 \times 10^{46}$ W	Multi-band
Derived jet velocity	v_{SCm}	$0.99 \cdot c = 2.97 \times 10^8$ m/s	P_{jet}/L constraint
Lorentz factor	γ	$1/\sqrt{1-0.99^2} \approx 7.09$	Special relativity

Derivation of v_{SCm} : The jet velocity is constrained such that the relativistic kinetic-power ratio $P_{jet}/L \approx 0.2$ is consistent with $v_{SCm}/c \approx 0.99$ for a relativistic jet.

Note on z: PAPER_360 used z=6.5 (the FU/Bi high-z quasar paper); PAPER_374 uses z=3.122 which appears in the standalone C++ code section of grok_share_11254865.txt (lines 5100+) where the `simulate_quasar_jet` function was annotated with these observational values. These represent two distinct epochs or sourcing conventions in the Grok thread.

2. Coupling Algorithm

The UQFF-NS coupling proceeds as follows:

1. Instantiate SGR1745-2900 proxy SMBH → MUGESystem sagA (SgrA* as quasar host)
2. Compute $g_{UQFF} = \text{compute_resonance_MUGE}(\text{sagA}, \text{ResonanceParams}\{\})$

```

→ master 12-term MUGE acceleration (PAPER_371)
3. Instantiate FluidSolver (N=32 grid, visc=0.0001, dt=0.1)
   (Jos Stam Stable Fluids method, PAPER_369)
4. Set jet_force = v_Scm_rel / 10.0 = 2.97×107 m/s2
5. For step = 1..10:
   a. Inject jet force into central column:
      for i ∈ [N/4, 3N/4]: v[i, N/2] += jet_force
   b. Add UQFF body force uniformly:
      for all cells: v[i,j] += g_UQFF / 1e30
   c. Execute NS step: diffuse → advect → project
6. Compute mean |v| = (1/N2) Σi,j √(uij2 + vij2)
7. Print ASCII velocity field:
   '#' → |v| > 1.0    '+' → |v| > 0.5
   '.' → |v| > 0.1    ' ' → |v| ≤ 0.1

```

3. Physical Interpretation

The coupling of $g_UQFF / 1e30$ as a body force in the NS grid models the effect of the vacuum-energy-mediated gravitational field from a SMBH (SgrA* proxy, $M=8.155 \times 10^{36}$ kg) on the fluid dynamics of a relativistic jet. The factor 1e30 normalises the UQFF acceleration (which spans $\sim 10^{-9}$ to 10^{39} m/s² across the 12 terms) to the NS grid scale (dimensionless fluid velocity units of order 1).

The relativistic velocity $v_SCm = 0.99c$ appears in the jet forcing term and in the Lorentz correction derived in PAPER_375.

4. Distinction from PAPER_360 and PAPER_369

Feature	PAPER_360	PAPER_369	PAPER_374
Object	J1610+1811	Generic SgrA*	J1610+1811
z	6.5	—	3.122
Physics	FU, Bi calculations	NS Stable Fluids	NS + UQFF resonance force
v _{jet}	—	1e8 m/s (generic)	0.99c = 2.97e8 m/s (relativistic)
UQFF coupling	None	Velocity only	Full 12-term MUGE body force

5. Implementation

C++: STAR_MAGIC_09SEPT_UQFF_MODULE.cpp, namespace J1610QuasarJet - Constants: z_redshift=3.122, P_jet=4e45, L_luminosity=2e46, v_Scm_rel=0.99c - Function: simulate_relativistic_quasar_jet(os, NS_steps=10)

Python: CondensedPhysics4.py, class J1610RelativisticQuasarJetUQFFNSCalculator (CP4 #22)
