

PAPER_744: M16 Eagle Nebula MUGE — Star Formation and Radiation Erosion

Author: Daniel T. Murphy

Framework: Universal Quantum Field Superconductive Framework (UQFF)

Session: 180 continuation | v5.38

Date: 2025

CP4 Class: #328 — M16EagleNebulaRadiationMUGECalculator

Abstract

The M16 Eagle Nebula (NGC 6611) is the iconic site of the “Pillars of Creation” — dense molecular cloud columns being sculpted by intense photoionization from massive OB stars. This paper derives the MUGE for M16 incorporating two new environmental terms: $M_{sf}(t)$, the time-dependent star formation rate modulating the total mass, and E_{rad} , the radiation erosion term that removes mass from pillar structures over time. These terms together capture the dynamic photo-evaporative environment that distinguishes star-forming nebulae from passive systems.

1. Introduction

M16 is an H II region and open cluster complex at 2 kpc distance, containing the giant OB association NGC 6611. The “Pillars of Creation” (Hubble 1995 image) are three columns of molecular hydrogen being photoevaporated by UV radiation from hot stars. The pillars host ongoing star formation (YSOs, protostars) while simultaneously losing mass to radiation erosion, creating a self-regulating gravitational environment.

Key parameters: - Distance = 2 kpc - $M_{cluster}$ (visible) = $5 \times 10^3 M_{\odot}$ - M_{cloud} (total) $\approx 8 \times 10^4 M_{\odot}$ - SFR $\approx 10^{-3} M_{\odot}/yr$ - $T_{ionized} \approx 10^4 K$ - UV flux = 10^7 Habings (ionizing radiation) - $B \approx 10^{-4} T$ (magnetic field in cloud)

2. M16 Eagle Nebula MUGE

$$\begin{aligned} g_{M16}(r,t) = & (G \cdot M(t)) / r^2 \cdot (1 + H(z) \cdot t) \cdot (1 - B/B_{crit}) \cdot (1 + M_{sf}(t)) \\ & + (U_{g1} + U_{g2} + U_{g3} + U_{g4}) \\ & + U_i \\ & + (\Lambda \cdot c^2 / 3) \\ & + (\hbar / \sqrt{(\Delta x \cdot \Delta p)}) \cdot \int (\psi \cdot H \cdot \psi \, dV) \cdot (2\pi / t_{Hubble}) \\ & + \rho_{gas} \cdot V \cdot g \\ & - E_{rad} \quad \quad \quad [radiation erosion -- NEW] \\ & + (M_{vis} + M_{DM}) \cdot (\delta \rho / \rho + 3 \cdot G \cdot M / r^3) \end{aligned}$$

3. $M_{sf}(t)$ — Time-Dependent Star Formation Rate Modulator

Star formation modifies the effective gravitational mass available:

$$M_{sf}(t) = SFR \cdot t / M_0$$

SFR = star formation rate ($M_{\odot}/yr$)

t = time elapsed (yr)
M_0 = initial cloud mass (M_⊙)

For M16:

M_sf(t) = (10⁻³ M_⊙/yr) · t / (8×10⁴ M_⊙)
M_sf(1 Myr) ≈ 0.0125 (1.25% mass converted)

This term enters multiplicatively in the gravitational term:

g_grav · (1 + M_sf(t)) ≈ g_grav · (1 + 0.0125) at t = 1 Myr

As star formation proceeds, the effective mass increases and local gravity strengthens, triggering further collapse — a positive feedback loop.

4. E_rad — Radiation Erosion Term

UV photons from OB stars photoevaporate the pillar surfaces, removing mass at:

$\dot{m}_{\text{evap}} = \Phi_{\text{UV}} \cdot m_{\text{H}} / (\alpha_{\text{B}} \cdot n_{\text{H}})$

$\Phi_{\text{UV}} = 10^7$ Habings = UV photon flux (photons/m²/s)
m_H = hydrogen mass = 1.67×10⁻²⁷ kg
 $\alpha_{\text{B}} = 2.6 \times 10^{-13}$ cm³/s (case B recombination)
n_H = 10³ cm⁻³ (column density)

The gravitational equivalent (effective deceleration from mass loss):

$E_{\text{rad}} = G \cdot \dot{m}_{\text{evap}} \cdot t / (r^2 \cdot M_{\text{cloud}})$

$E_{\text{rad}}(1 \text{ Myr}, r=0.5 \text{ pc}) \approx 3 \times 10^{-12} \text{ m/s}^2$ (opposing gravity)

This erosion term opposes collapse, creating the dynamic equilibrium observed in pillar lifetimes (~few Myr).

5. UQFF Gravity Components

U_g1: magnetic dipole from cloud B-field threading

$\mu_{\text{dipole}} = \text{charge density} \times \text{pillar area} \times \omega_{\text{rotation}}$
≈ 10⁻⁴⁵ A·m² (weak for molecular cloud)

U_g2: aether-superconductive field in ionized HII region

B_super = $\mu_0 \cdot H_{\text{aether_HII}} \approx 5 \text{ T}$ (elevated near ionization front)
U_g2 ≈ 10⁷ J/m³

U_g3: external gravity from cluster mass distribution

$U_{\text{g3}} = G \cdot M_{\text{NGC6611}} / r_{\text{cluster}}^2$

U_g4: galactic tidal field at 2 kpc from center

$U_{\text{g4}} \approx 2.5 \times 10^{-20} \text{ J/m}^3$

6. Equilibrium Analysis

The pillars maintain their structure while:

$$g_{\text{grav}} \cdot (1 + M_{\text{sf}}) > E_{\text{rad}} \quad [\text{net infall}]$$

When E_{rad} overcomes $(1 + M_{\text{sf}})$:

$$E_{\text{rad}} / g_{\text{grav}} > (1 + M_{\text{sf}}) \quad [\text{pillar evaporated}]$$

Current M16 pillar state: - $g_{\text{grav}} \approx 2 \times 10^{-11} \text{ m/s}^2$ - $E_{\text{rad}} \approx 3 \times 10^{-12} \text{ m/s}^2$ - $(1 + M_{\text{sf}}) \approx 1.0125$ - Net: pillars are infalling (star formation wins at present epoch)

7. Temporal Evolution

$$M(t) = M_0 \cdot (1 + M_{\text{sf}}(t)) - \dot{m}_{\text{evap}} \cdot t$$

At $t = 3 \text{ Myr}$ (estimated pillar lifetime):

$$M(3 \text{ Myr}) = M_0 \cdot (1.037) - \Delta m_{\text{evap}} \approx 0.85 M_0$$

The pillars are expected to be fully ionized within $\sim 5 \text{ Myr}$, making M16 a transient feature of galactic star formation history.

8. Comparison to Pillars of Creation (Companion UQFF Module)

M16 shares the Pillars of Creation geometry with the pre-existing CP4 module. The key distinction: - **Pillars of Creation** module: static geometry, UV photoionization - **This paper (M16)**: full temporal MUGE with $M_{\text{sf}}(t) + E_{\text{rad}}$ dynamic evolution

9. Conclusion

The M16 Eagle Nebula MUGE introduces two novel UQFF terms: $M_{\text{sf}}(t)$ captures positive-feedback star formation mass growth, while E_{rad} captures the opposing radiation erosion. Together they define the pillar equilibrium condition and predict pillar lifetimes consistent with current observations ($\sim 3\text{--}5 \text{ Myr}$). This framework generalizes to all photo-evaporating star-forming regions in the spiral arm environment.

Copyright - Daniel T. Murphy, daniel.murphy00@gmail.com. UQFF Framework. PAPER_744, CP4 class #328. Session 180 continuation v5.38.