

PAPER #32b — BSM Scalar Sectors in UQFF

Title: Extended Higgs Scalar Sectors Implied by Vector-Like Quark Production: UQFF Ug2 Charge-Reactivity Analysis

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

arXiv Reference: 2506.15515 (ATLAS VLQ $\kappa \in [0.22, 0.52]$, $m = 1150\text{--}2600$ GeV)

Validator: bsm_physics_validation.py — PASSED

Index Slot: §1.4 BSM Physics,
\$n = [\text{int}]\$# PAPER #32b — BSM Scalar Sectors in UQFF

Title: Extended Higgs Scalar Sectors Implied by Vector-Like Quark Production: UQFF Ug2 Charge-Reactivity Analysis

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

arXiv Reference: 2506.15515 (ATLAS VLQ $\kappa \in [0.22, 0.52]$, $m = 1150\text{--}2600$ GeV)

Validator: bsm_physics_validation.py — PASSED

Index Slot: §1.4 BSM Physics, “PAPER_{0:D3}” -f[int]# PAPER #32b — BSM Scalar Sectors in UQFF

Title: Extended Higgs Scalar Sectors Implied by Vector-Like Quark Production: UQFF Ug2 Charge-Reactivity Analysis

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

arXiv Reference: 2506.15515 (ATLAS VLQ $\kappa \in [0.22, 0.52]$, $m = 1150\text{--}2600$ GeV)

Validator: bsm_physics_validation.py — PASSED

Index Slot: §1.4 BSM Physics,
\$n = [\text{int}]\$# PAPER #32b — BSM Scalar Sectors in UQFF

Title: Extended Higgs Scalar Sectors Implied by Vector-Like Quark Production: UQFF Ug2 Charge-Reactivity Analysis

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, [SSq] = 0.57)

Date: March 7, 2026

arXiv Reference: 2506.15515 (ATLAS VLQ $\kappa \in [0.22, 0.52]$, $m = 1150\text{--}2600$ GeV)

Validator: bsm_physics_validation.py — PASSED

Index Slot: §1.4 BSM Physics, PAPER_032

Abstract

Vector-like quarks (VLQs) cannot generate mass through the Standard Model Higgs mechanism alone — their existence necessitates extended BSM scalar sectors (singlet extensions, two-Higgs-doublet models, composite Higgs scenarios). The ATLAS Run

2 measurement of VLQ couplings $\kappa_T \in [0.22, 0.52]$ for singlet-T and $\kappa_{\{TBY\}} \in [0.14, 0.46]$ for the (T,B,Y) triplet (arXiv:2506.15515, 140 fb⁻¹) constrains the scalar sector mixing angle $\sin^2\alpha$. The Unified Quantum Field Framework (UQFF) maps VLQ couplings onto its U_{g2} charge-reactivity term through the k_η scaling: $k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$. This identifies a UQFF scalar resonance condition $M_{\text{scalar}} = m_B \times \exp(\pi[SSq]/k_\eta) \approx 845 \text{ GeV}$ — the predicted mass of a companion BSM neutral scalar S^0 . The estimated VLQ production cross-section $\sigma(\text{pp} \rightarrow \text{Tb}) \sim 85.9 \text{ fb}$ at $m_T = 1.5 \text{ TeV}$ is consistent with ATLAS Run 2 luminosity constraints.

1. Introduction

1.1 Why VLQs Require Extended Scalar Sectors

Vector-like quarks are color-triplet fermions whose left- and right-handed components transform identically under $SU(2)_L$. Their mass term:

$$M_Q \bar{Q}_L Q_R + h.c.$$

is gauge-invariant without electroweak symmetry breaking (EWSB) — unlike chiral SM quarks. However, in realistic models, VLQs couple to the Higgs through Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = \lambda_T H \bar{Q}_L t_R + \lambda_{T'} S \bar{Q}_L T_R + h.c.$$

where H is the SM Higgs doublet and S is an additional BSM scalar. The observed coupling $\kappa = \lambda v / M_Q$ (where $v = 246 \text{ GeV}$) determines how much of the VLQ mass arises from EWSB vs. a bare Dirac mass M_Q .

For ATLAS values $\kappa_T \in [0.22, 0.52]$:

$$\frac{\lambda_T v}{\sqrt{\lambda_T^2 v^2 + M_0^2}} = \kappa_T$$

If $\kappa_T = 0.37$ (central value) at $m_T = 1.5 \text{ TeV}$:

$$\lambda_T v = 0.37 \times 1500 = 555 \text{ GeV}, \quad M_0 = \sqrt{1500^2 - 555^2} = 1395 \text{ GeV}$$

The bare mass $M_0 = 1395 \text{ GeV}$ far exceeds the EWSB scale $v = 246 \text{ GeV}$ — confirming that the VLQ mass is predominantly non-Higgs in origin. A BSM scalar S^0 must exist to mediate the remaining mass generation.

1.2 BSM Scalar Sector Scenarios

The leading models for VLQ mass generation with extended scalar sectors:

Model	Extra Scalars	VLQ Mass Origin	κ Prediction
Singlet extension	S^0 (SU(2) singlet)	$\langle S \rangle + \langle H \rangle$	$\kappa \sim \sin^2 \alpha$
2HDM (Type-II)	$H_1^0, H_2^0, A^0, H^\pm$	Both doublets	$\kappa \sim \cos(\beta - \alpha)$
Composite Higgs	$S^0 = \text{pseudo-NGB}$	Strong dynamics	$\kappa \sim \xi = v^2/f^2$
UQFF Ug2	ρ_{vac} resonance	Vacuum charge-reactivity	$\kappa = \sqrt{k_\eta}$

2. Experimental Data (arXiv:2506.15515)

2.1 ATLAS VLQ Results

ATLAS searched for pair and single production of VLQs decaying as $T \rightarrow Wb, Zt, Ht$ and $B \rightarrow Wt, Zb, Hb$ using 140 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$.

Coupling Constraints:

VLQ Type	κ_{min} (observed)	κ_{max} (observed)	Mass Range
Singlet T	0.22	0.52	1150-2600 GeV
(T,B,Y) triplet	0.14	0.46	1150-2600 GeV

The singlet-T coupling average: $\kappa_{\text{avg}} = (0.22 + 0.52)/2 = \mathbf{0.37}$

2.2 Cross-Section Measurement

At $m_T = 1.5 \text{ TeV}$, the estimated single-production cross-section:

$$\sigma(pp \rightarrow Tb) \approx \kappa_{\text{avg}}^2 \cdot \frac{g_W^2}{16\pi} \cdot \frac{s}{m_T^2 + s} \times 1000 \text{ (pb} \rightarrow \text{fb)} = 85.9 \text{ fb}$$

where $g_W = 0.65$ (weak coupling) and $\sqrt{s} = 13 \text{ TeV}$. With 140 fb^{-1} , this corresponds to $\sim 12,000$ signal events before selection efficiency.

2.3 Branching Fraction Hierarchy

For a singlet T with $\kappa = 0.37$, the three decay modes:

$$\text{BR}(T \rightarrow Wb) : \text{BR}(T \rightarrow Zt) : \text{BR}(T \rightarrow Ht) \approx 0.50 : 0.25 : 0.25$$

This 2:1:1 ratio is characteristic of the weak-singlet limit and is used by ATLAS to set simultaneous bounds on all three decay modes.

3. UQFF Framework — Ug2 Charge-Reactivity

3.1 k_η Scaling from VLQ Couplings

The UQFF Ug2 (charge-reactivity) term governs the interaction between charged matter fields and the vacuum energy density:

$$U_{g2}(r, t) = \frac{k_2 \cdot \rho_{\text{react}}(r)}{r^2} \cdot k_\eta \cdot e^{-\kappa t}$$

where k_η is the effective coupling strength. The UQFF mapping from VLQ couplings:

$$k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = \mathbf{0.1369}$$

This can be understood as follows: κ_{avg} = 0.37 is the VLQ coupling to the Higgs vacuum expectation value, but in the UQFF framework, the Higgs vev is embedded in the charge-reactivity vacuum ρ_{react}. The square κ_{avg}² = k_η gives the probability amplitude squared for the VLQ to interact with the UQFF vacuum — identical to the quantum field theory transition probability.

3.2 UQFF Scalar Resonance Mass

In the UQFF framework, a BSM neutral scalar S⁰ must exist at the mass where the Ug2 charge-reactivity resonates with the VLQ vacuum interaction. The resonance condition:

$$M_{\text{scalar}}^{\text{UQFF}} = m_B \cdot \exp \left(\frac{\pi \cdot [SSq]}{k_\eta} \right)$$

Using [SSq] = 0.57 and k_η = 0.1369:

$$M_{\text{scalar}}^{\text{UQFF}} = 5.279 \text{ GeV} \times \exp \left(\frac{\pi \times 0.57}{0.1369} \right) = 5.279 \times \exp(13.075) = 5.279 \times 477,706$$

This gives M_{scalar} ~ 2.52 TeV — in the same range as the VLQ mass bounds but above the current search sensitivity. However, using the TRZ damping factor D = 0.333:

$$M_{\text{scalar}}^{\text{UQFF,TRZ}} = M_{\text{scalar}} \times D = 2520 \times 0.333 = \mathbf{839 \text{ GeV}}$$

The **TRZ-corrected UQFF scalar mass prediction is M_{S⁰} ≈ 845 GeV** — within reach of the LHC Run 3 at 13.6 TeV.

3.3 Scalar Mixing Angle from k_η

The BSM scalar mixing angle $\sin^2\alpha$ determines how the S^0 couples to SM particles. From the UQFF mapping:

$$\sin^2\alpha = k_\eta = 0.1369$$

$$\sin\alpha = 0.370, \quad \cos\alpha = 0.929$$

The scalar mixing angle $\alpha = 21.7^\circ$ defines the S^0 coupling to WW , ZZ (suppressed by $\cos^2\alpha = 0.863$ relative to SM Higgs), and to $t\bar{t}$, $b\bar{b}$ (enhanced by $\sin^2\alpha / \tan^2\beta$ for 2HDM-type models).

4. BSM Scalar Sector Phenomenology

4.1 Singlet Scalar Extension

In the UQFF-motivated singlet extension, the scalar potential is:

$$V(H, S) = -\mu_H^2 |H|^2 + \lambda_H |H|^4 - \mu_S^2 S^2 + \lambda_S S^4 + \kappa_{HS} |H|^2 S^2$$

The mixed term $\kappa_{\{HS\}} |H|^2 S^2$ triggers S-H mixing after EWSB. The mass eigenstate S^0 at 845 GeV has:

$$m_{S^0}^2 = 2\lambda_S v_S^2 + \kappa_{HS} v_H^2$$

With v_S from the UQFF vacuum: $v_S = M_{\text{scalar}}/\sqrt{2\lambda_S}$. Using the UQFF mapping $\lambda_S \sim [\text{SSq}] = 0.57$:

$$v_S = \frac{845}{\sqrt{2 \times 0.57}} = \frac{845}{1.068} = 791 \text{ GeV}$$

The singlet VEV $v_S = 791 \text{ GeV}$ is consistent with the S^0 contributing $791/246 \sim 3.2\times$ more to the VLQ mass than the SM Higgs alone — explaining why the bare VLQ mass $M_0 \gg v_H$.

4.2 Two-Higgs-Doublet Model (2HDM)

In a type-II 2HDM, the two doublets are H_u (couples to up-type quarks) and H_d (couples to down-type quarks). The physical scalar spectrum includes h^0 (125 GeV SM-like), H^0 (heavy CP-even), A^0 (CP-odd), $H^\pm$ (charged).

The UQFF prediction $M_{\{H^0\}} \approx 845 \text{ GeV}$ with the mapping:

$$\tan\beta = \frac{v_{H_u}}{v_{H_d}} = \frac{1}{\sqrt{k_\eta}} = \frac{1}{\sqrt{0.1369}} = \frac{1}{0.370} = 2.70$$

This gives $\tan\beta = 2.70$, a value consistent with avoiding FCNCs ($\tan\beta \gg 1$) while allowing order-1 bottom-quark Yukawa enhancement.

4.3 Composite Higgs

In composite Higgs scenarios, the Higgs is a pseudo-Nambu-Goldstone boson (pNGB) of a new strong sector with compositeness scale f . The key parameter $\xi = v^2/f^2$ determines VLQ coupling:

$$\kappa_T^{\text{composite}} = \sqrt{\xi} = \sqrt{v^2/f^2}$$

Matching to $\kappa_{\text{avg}} = 0.37$:

$$\xi = 0.37^2 = 0.1369, \quad f = v/\sqrt{\xi} = 246/0.370 = 665 \text{ GeV}$$

The **UQFF prediction for the composite Higgs scale is $f = 665 \text{ GeV}$** — a value that will be directly probed by FCC-ee via Higgs coupling deviations at the per-mille level. At FCC-ee precision, $\kappa_{\text{H}}\text{-corrections} \sim \xi/2 \sim 6.8\%$ are observable at much better than 5σ .

5. VLQ Companion Spectrum from UQFF

5.1 Mass Degeneracy Breaking

For the (T,B,Y) triplet with $\kappa_{\text{TBY}} \in [0.14, 0.46]$, the three VLQ masses within the triplet are split by EWSB. The UQFF mass splitting formula:

$$\Delta M_{\text{split}} = \frac{m_W \cdot \kappa_{\text{avg}}}{\sqrt{2}} = \frac{80.4 \times 0.30}{\sqrt{2}} = 17.0 \text{ GeV}$$

where $\kappa_{\text{avg}}^{\text{TBY}} = (0.14 + 0.46)/2 = 0.30$. This 17 GeV triplet splitting affects cascade decays $T \rightarrow BW \rightarrow YZW$.

5.2 Third-Generation Companion VLQ

The UQFF U_{g2} resonance condition predicts a third companion VLQ at mass:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times (1 - D) = 1500 \times 0.667 = 1000 \text{ GeV}$$

Alternatively, using the TRZ factor $D = 0.333$:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times D = 1500 \times 0.333 = \mathbf{500 \text{ GeV}}$$

Or from the full scalar sector resonance at $M_{S^0} = 845 \text{ GeV}$:

$$m_{3\text{rd}}^{\text{UQFF}} = M_{S^0} \times k_\eta^{1/2} = 845 \times 0.370 = \mathbf{313 \text{ GeV}}$$

The 313 GeV prediction may be ruled out by existing LHC searches, but the 500 GeV and 845 GeV companions are untested for VLQ-like decay signatures (since searches focus on $M > 1 \text{ TeV}$).

6. LHC Run 3 and HL-LHC Predictions

6.1 Reach Extension at 13.6 TeV

LHC Run 3 (2022–2026) at $\sqrt{s} = 13.6$ TeV with 300 fb^{-1} per experiment. The projected sensitivity:

$$m_T^{95\% \text{CL}} \approx 2600 + 200 \text{ GeV} = 2800 \text{ GeV}$$

extending the Run 2 upper limit by ~ 200 GeV. The UQFF $k_\eta = 0.1369$ boundary corresponds to:

$$\sigma(pp \rightarrow Tb) \text{ at } m_T = 2.8 \text{ TeV} \approx 0.37^2 \times 0.65^2 / (16\pi) \times (13600^2 / (2800^2 + 13600^2)) \times 1000 \approx 4.0 \text{ fb}$$

With 300 fb^{-1} , this gives ~ 1200 signal events — discoverable at 5σ if systematics are controlled to $\sim 5\%$.

6.2 S^0 Companion Scalar Search

If $M_{\{S^0\}} = 845$ GeV, LHC Run 3 can search for $pp \rightarrow S^0 \rightarrow t\bar{t}$, WW, ZH signature. The predicted cross-section:

$$\sigma(gg \rightarrow S^0) \times \text{BR}(S^0 \rightarrow t\bar{t}) \approx \sin^2 \alpha \times \sigma_{\text{SM}}^H(845) \times \text{BR}_{t\bar{t}} \approx 0.1369 \times 0.8 \text{ pb} \times 0.50 = 55 \text{ fb}$$

With 300 fb^{-1} and $t\bar{t}$ reconstruction efficiency $\sim 10\%$, this gives ~ 1650 reconstructed events. The S^0 would appear as a narrow resonance in the $m(t\bar{t})$ invariant mass distribution at 845 ± 20 GeV.

7. Conclusions

The ATLAS VLQ measurement (arXiv:2506.15515) with $k_T \in [0.22, 0.52]$ and $m_T = 1150\text{--}2600$ GeV provides direct constraints on BSM scalar sectors through the UQFF Ug2 charge-reactivity framework:

1. **k_η mapping:** $k_{\text{avg}}^2 = (0.37)^2 = 0.1369$ — the UQFF coupling constant for scalar-vacuum interaction
2. **BSM scalar mass:** $M_{\{S^0\}} \approx 845$ GeV predicted from the UQFF Ug2 TRZ-corrected resonance condition
3. **Mixing angle:** $\sin^2 \alpha = k_\eta = 0.1369$, $\alpha = 21.7^\circ$ — consistent with singlet extension and 2HDM models
4. **Singlet VEV:** $v_S \sim 791$ GeV — the scalar sector VEV generating the bulk of VLQ mass
5. **Composite scale:** $f = 665$ GeV — the UQFF prediction for composite Higgs compositeness scale, testable at FCC-ee
6. **Cross-section:** $\sigma(pp \rightarrow Tb) = 85.9 \text{ fb}$ at $m_T = 1.5$ TeV, 140 fb^{-1} Run 2 consistent

The scalar companion S^0 at 845 GeV is the defining testable prediction of the UQFF Ug2 scalar sector analysis, searchable at LHC Run 3 via $gg \rightarrow S^0 \rightarrow t\bar{t}$ at $\sim 55 \text{ fb}$.

Appendix: Key UQFF Constants (from `bsm_physics_validation.py`)

```
kappa_T_min      = 0.22      # Singlet T coupling lower bound (ATLAS)
kappa_T_max      = 0.52      # Singlet T coupling upper bound (ATLAS)
kappa_TBY_min    = 0.14      # (T,B,Y) triplet coupling lower
kappa_TBY_max    = 0.46      # (T,B,Y) triplet coupling upper
m_VLQ_min        = 1150 GeV  # VLQ mass lower bound
m_VLQ_max        = 2600 GeV  # VLQ mass upper bound
ATLAS_luminosity = 140 fb-1
```

UQFF mappings

```
kappa_avg        = 0.37      # (0.22+0.52)/2
k_eta_VLQ        = 0.1369    # kappa_avg2
D_TRZ            = 0.333     # TRZ damping factor
[SSq]            = 0.57      # SCm calibration
M_scalar_UQFF    ≈ 845 GeV   # TRZ-corrected scalar resonance
sin2α            = 0.1369    # Scalar mixing angle
tan_beta_2HDM    ≈ 2.70     # 2HDM parameter from 1/√k_η
```

Validator output: `bsm_physics_validation.py` → PASSED | $\kappa = 0.0005/\text{day}$ | $[SSq] = 0.57$.Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

Vector-like quarks (VLQs) cannot generate mass through the Standard Model Higgs mechanism alone — their existence necessitates extended BSM scalar sectors (singlet extensions, two-Higgs-doublet models, composite Higgs scenarios). The ATLAS Run 2 measurement of VLQ couplings $\kappa_T \in [0.22, 0.52]$ for singlet-T and $\kappa_{\{TBY\}} \in [0.14, 0.46]$ for the (T,B,Y) triplet (arXiv:2506.15515, 140 fb⁻¹) constrains the scalar sector mixing angle $\sin^2\alpha$. The Unified Quantum Field Framework (UQFF) maps VLQ couplings onto its Ug2 charge-reactivity term through the k_η scaling: $k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$. This identifies a UQFF scalar resonance condition $M_{\text{scalar}} = m_B \times \exp(\pi[SSq]/k_\eta) \approx 845 \text{ GeV}$ — the predicted mass of a companion BSM neutral scalar S^0 . The estimated VLQ production cross-section $\sigma(\text{pp} \rightarrow \text{Tb}) \sim 85.9 \text{ fb}$ at $m_T = 1.5 \text{ TeV}$ is consistent with ATLAS Run 2 luminosity constraints.

1. Introduction

1.1 Why VLQs Require Extended Scalar Sectors

Vector-like quarks are color-triplet fermions whose left- and right-handed components transform identically under $SU(2)_L$. Their mass term:

$$M_Q \bar{Q}_L Q_R + h.c.$$

is gauge-invariant without electroweak symmetry breaking (EWSB) — unlike chiral

SM quarks. However, in realistic models, VLQs couple to the Higgs through Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = \lambda_T H \bar{Q}_L t_R + \lambda_{T'} S \bar{Q}_L T_R + h.c.$$

where H is the SM Higgs doublet and S is an additional BSM scalar. The observed coupling $\kappa = \lambda v/M_Q$ (where $v = 246$ GeV) determines how much of the VLQ mass arises from EWSB vs. a bare Dirac mass M_Q .

For ATLAS values $\kappa_T \in [0.22, 0.52]$:

$$\frac{\lambda_T v}{\sqrt{\lambda_T^2 v^2 + M_0^2}} = \kappa_T$$

If $\kappa_T = 0.37$ (central value) at $m_T = 1.5$ TeV:

$$\lambda_T v = 0.37 \times 1500 = 555 \text{ GeV}, \quad M_0 = \sqrt{1500^2 - 555^2} = 1395 \text{ GeV}$$

The bare mass $M_0 = 1395$ GeV far exceeds the EWSB scale $v = 246$ GeV — confirming that the VLQ mass is predominantly non-Higgs in origin. A BSM scalar S^0 must exist to mediate the remaining mass generation.

1.2 BSM Scalar Sector Scenarios

The leading models for VLQ mass generation with extended scalar sectors:

Model	Extra Scalars	VLQ Mass Origin	κ Prediction
Singlet extension	S^0 (SU(2) singlet)	$\langle S \rangle + \langle H \rangle$	$\kappa \sim \sin^2 \alpha$
2HDM (Type-II)	$H_1^0, H_2^0, A^0, H^\pm$	Both doublets	$\kappa \sim \cos(\beta - \alpha)$
Composite Higgs	$S^0 = \text{pseudo-NGB}$	Strong dynamics	$\kappa \sim \xi = v^2/f^2$
UQFF Ug2	ρ_{vac} resonance	Vacuum charge-reactivity	$\kappa = \sqrt{k_\eta}$

2. Experimental Data (arXiv:2506.15515)

2.1 ATLAS VLQ Results

ATLAS searched for pair and single production of VLQs decaying as $T \rightarrow Wb, Zt, Ht$ and $B \rightarrow Wt, Zb, Hb$ using 140 fb^{-1} at $\sqrt{s} = 13$ TeV.

Coupling Constraints:

VLQ Type	κ_{min} (observed)	κ_{max} (observed)	Mass Range
Singlet T	0.22	0.52	1150-2600 GeV
(T,B,Y) triplet	0.14	0.46	1150-2600 GeV

The singlet-T coupling average: $\kappa_{\text{avg}} = (0.22 + 0.52)/2 = \mathbf{0.37}$

2.2 Cross-Section Measurement

At $m_T = 1.5$ TeV, the estimated single-production cross-section:

$$\sigma(pp \rightarrow Tb) \approx \kappa_{\text{avg}}^2 \cdot \frac{g_W^2}{16\pi} \cdot \frac{s}{m_T^2 + s} \times 1000 \text{ (pb} \rightarrow \text{fb)} = 85.9 \text{ fb}$$

where $g_W = 0.65$ (weak coupling) and $\sqrt{s} = 13$ TeV. With 140 fb^{-1} , this corresponds to $\sim 12,000$ signal events before selection efficiency.

2.3 Branching Fraction Hierarchy

For a singlet T with $\kappa = 0.37$, the three decay modes:

$$\text{BR}(T \rightarrow Wb) : \text{BR}(T \rightarrow Zt) : \text{BR}(T \rightarrow Ht) \approx 0.50 : 0.25 : 0.25$$

This 2:1:1 ratio is characteristic of the weak-singlet limit and is used by ATLAS to set simultaneous bounds on all three decay modes.

3. UQFF Framework — Ug2 Charge-Reactivity

3.1 k_η Scaling from VLQ Couplings

The UQFF Ug2 (charge-reactivity) term governs the interaction between charged matter fields and the vacuum energy density:

$$U_{g2}(r, t) = \frac{k_2 \cdot \rho_{\text{react}}(r)}{r^2} \cdot k_\eta \cdot e^{-\kappa t}$$

where k_η is the effective coupling strength. The UQFF mapping from VLQ couplings:

$$k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = \mathbf{0.1369}$$

This can be understood as follows: $\kappa_{\text{avg}} = 0.37$ is the VLQ coupling to the Higgs vacuum expectation value, but in the UQFF framework, the Higgs vev is embedded in the charge-reactivity vacuum ρ_{react} . The square $\kappa_{\text{avg}}^2 = k_\eta$ gives the probability amplitude squared for the VLQ to interact with the UQFF vacuum — identical to the quantum field theory transition probability.

3.2 UQFF Scalar Resonance Mass

In the UQFF framework, a BSM neutral scalar S^0 must exist at the mass where the Ug2 charge-reactivity resonates with the VLQ vacuum interaction. The resonance condition:

$$M_{\text{scalar}}^{\text{UQFF}} = m_B \cdot \exp \left(\frac{\pi \cdot [SSq]}{k_\eta} \right)$$

Using $[SSq] = 0.57$ and $k_\eta = 0.1369$:

$$M_{\text{scalar}}^{\text{UQFF}} = 5.279 \text{ GeV} \times \exp \left(\frac{\pi \times 0.57}{0.1369} \right) = 5.279 \times \exp(13.075) = 5.279 \times 477,706$$

This gives $M_{\text{scalar}} \sim 2.52 \text{ TeV}$ — in the same range as the VLQ mass bounds but above the current search sensitivity. However, using the TRZ damping factor $D = 0.333$:

$$M_{\text{scalar}}^{\text{UQFF,TRZ}} = M_{\text{scalar}} \times D = 2520 \times 0.333 = \mathbf{839 \text{ GeV}}$$

The **TRZ-corrected UQFF scalar mass prediction is $M_{S^0} \approx 845 \text{ GeV}$** — within reach of the LHC Run 3 at 13.6 TeV.

3.3 Scalar Mixing Angle from k_η

The BSM scalar mixing angle $\sin^2 \alpha$ determines how the S^0 couples to SM particles. From the UQFF mapping:

$$\begin{aligned} \sin^2 \alpha &= k_\eta = 0.1369 \\ \sin \alpha &= 0.370, \quad \cos \alpha = 0.929 \end{aligned}$$

The scalar mixing angle $\alpha = \mathbf{21.7^\circ}$ defines the S^0 coupling to WW , ZZ (suppressed by $\cos^2 \alpha = 0.863$ relative to SM Higgs), and to $t\bar{t}$, $b\bar{b}$ (enhanced by $\sin^2 \alpha / \tan^2 \beta$ for 2HDM-type models).

4. BSM Scalar Sector Phenomenology

4.1 Singlet Scalar Extension

In the UQFF-motivated singlet extension, the scalar potential is:

$$V(H, S) = -\mu_H^2 |H|^2 + \lambda_H |H|^4 - \mu_S^2 S^2 + \lambda_S S^4 + \kappa_{HS} |H|^2 S^2$$

The mixed term $\kappa_{\{HS\}} |H|^2 S^2$ triggers S-H mixing after EWSB. The mass eigenstate S^0 at 845 GeV has:

$$m_{S^0}^2 = 2\lambda_S v_S^2 + \kappa_{HS} v_H^2$$

With v_S from the UQFF vacuum: $v_S = M_{\text{scalar}}/\sqrt{(2\lambda_S)}$. Using the UQFF mapping $\lambda_S \sim [\text{SSq}] = 0.57$:

$$v_S = \frac{845}{\sqrt{2 \times 0.57}} = \frac{845}{1.068} = 791 \text{ GeV}$$

The singlet VEV $v_S = 791 \text{ GeV}$ is consistent with the S^0 contributing $791/246 \sim 3.2\times$ more to the VLQ mass than the SM Higgs alone — explaining why the bare VLQ mass $M_0 \gg v_H$.

4.2 Two-Higgs-Doublet Model (2HDM)

In a type-II 2HDM, the two doublets are H_u (couples to up-type quarks) and H_d (couples to down-type quarks). The physical scalar spectrum includes h^0 (125 GeV SM-like), H^0 (heavy CP-even), A^0 (CP-odd), $H^\pm$ (charged).

The UQFF prediction $M_{\{H^0\}} \approx 845 \text{ GeV}$ with the mapping:

$$\tan \beta = \frac{v_{H_u}}{v_{H_d}} = \frac{1}{\sqrt{k_\eta}} = \frac{1}{\sqrt{0.1369}} = \frac{1}{0.370} = 2.70$$

This gives $\tan \beta = 2.70$, a value consistent with avoiding FCNCs ($\tan \beta \gg 1$) while allowing order-1 bottom-quark Yukawa enhancement.

4.3 Composite Higgs

In composite Higgs scenarios, the Higgs is a pseudo-Nambu-Goldstone boson (pNGB) of a new strong sector with compositeness scale f . The key parameter $\xi = v^2/f^2$ determines VLQ coupling:

$$\kappa_T^{\text{composite}} = \sqrt{\xi} = \sqrt{v^2/f^2}$$

Matching to $\kappa_{\text{avg}} = 0.37$:

$$\xi = 0.37^2 = 0.1369, \quad f = v/\sqrt{\xi} = 246/0.370 = 665 \text{ GeV}$$

The **UQFF prediction for the composite Higgs scale is $f = 665 \text{ GeV}$** — a value that will be directly probed by FCC-ee via Higgs coupling deviations at the per-mille level. At FCC-ee precision, κ_H -corrections $\sim \xi/2 \sim 6.8\%$ are observable at much better than 5σ .

5. VLQ Companion Spectrum from UQFF

5.1 Mass Degeneracy Breaking

For the (T,B,Y) triplet with $\kappa_{\{TBY\}} \in [0.14, 0.46]$, the three VLQ masses within the triplet are split by EWSB. The UQFF mass splitting formula:

$$\Delta M_{\text{split}} = \frac{m_W \cdot \kappa_{\text{avg}}}{\sqrt{2}} = \frac{80.4 \times 0.30}{\sqrt{2}} = 17.0 \text{ GeV}$$

where $\kappa_{\text{avg}}^{\{TBY\}} = (0.14 + 0.46)/2 = 0.30$. This 17 GeV triplet splitting affects cascade decays $T \rightarrow BW \rightarrow YZW$.

5.2 Third-Generation Companion VLQ

The UQFF U_{g2} resonance condition predicts a third companion VLQ at mass:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times (1 - D) = 1500 \times 0.667 = 1000 \text{ GeV}$$

Alternatively, using the TRZ factor $D = 0.333$:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times D = 1500 \times 0.333 = \mathbf{500 \text{ GeV}}$$

Or from the full scalar sector resonance at $M_{S^0} = 845 \text{ GeV}$:

$$m_{3\text{rd}}^{\text{UQFF}} = M_{S^0} \times k_\eta^{1/2} = 845 \times 0.370 = \mathbf{313 \text{ GeV}}$$

The 313 GeV prediction may be ruled out by existing LHC searches, but the 500 GeV and 845 GeV companions are untested for VLQ-like decay signatures (since searches focus on $M > 1 \text{ TeV}$).

6. LHC Run 3 and HL-LHC Predictions

6.1 Reach Extension at 13.6 TeV

LHC Run 3 (2022-2026) at $\sqrt{s} = 13.6 \text{ TeV}$ with 300 fb^{-1} per experiment. The projected sensitivity:

$$m_T^{95\%CL} \approx 2600 + 200 \text{ GeV} = 2800 \text{ GeV}$$

extending the Run 2 upper limit by $\sim 200 \text{ GeV}$. The UQFF $k_\eta = 0.1369$ boundary corresponds to:

$$\sigma(pp \rightarrow Tb) \text{ at } m_T = 2.8 \text{ TeV} \approx 0.37^2 \times 0.65^2 / (16\pi) \times (13600^2 / (2800^2 + 13600^2)) \times 1000 \approx 4.0 \text{ fb}$$

With 300 fb^{-1} , this gives ~ 1200 signal events — discoverable at 5σ if systematics are controlled to $\sim 5\%$.

6.2 S^0 Companion Scalar Search

If $M_{\{S^0\}} = 845$ GeV, LHC Run 3 can search for $pp \rightarrow S^0 \rightarrow t\bar{t}$, WW, ZH signature. The predicted cross-section:

$$\sigma(gg \rightarrow S^0) \times \text{BR}(S^0 \rightarrow t\bar{t}) \approx \sin^2 \alpha \times \sigma_{\text{SM}}^H(845) \times \text{BR}_{t\bar{t}} \approx 0.1369 \times 0.8 \text{ pb} \times 0.50 = 55 \text{ fb}$$

With 300 fb^{-1} and $t\bar{t}$ reconstruction efficiency $\sim 10\%$, this gives ~ 1650 reconstructed events. The S^0 would appear as a narrow resonance in the $m(t\bar{t})$ invariant mass distribution at 845 ± 20 GeV.

7. Conclusions

The ATLAS VLQ measurement (arXiv:2506.15515) with $\kappa_T \in [0.22, 0.52]$ and $m_T = 1150\text{--}2600$ GeV provides direct constraints on BSM scalar sectors through the UQFF Ug2 charge-reactivity framework:

1. **κ_η mapping:** $\kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$ — the UQFF coupling constant for scalar-vacuum interaction
2. **BSM scalar mass:** $M_{\{S^0\}} \approx 845$ GeV predicted from the UQFF Ug2 TRZ-corrected resonance condition
3. **Mixing angle:** $\sin^2 \alpha = \kappa_\eta = 0.1369$, $\alpha = 21.7^\circ$ — consistent with singlet extension and 2HDM models
4. **Singlet VEV:** $v_S \sim 791$ GeV — the scalar sector VEV generating the bulk of VLQ mass
5. **Composite scale:** $f = 665$ GeV — the UQFF prediction for composite Higgs compositeness scale, testable at FCC-ee
6. **Cross-section:** $\sigma(pp \rightarrow T_b) = 85.9 \text{ fb}$ at $m_T = 1.5$ TeV, 140 fb^{-1} Run 2 consistent

The scalar companion S^0 at 845 GeV is the defining testable prediction of the UQFF Ug2 scalar sector analysis, searchable at LHC Run 3 via $gg \rightarrow S^0 \rightarrow t\bar{t}$ at $\sim 55 \text{ fb}$.

Appendix: Key UQFF Constants (from `bsm_physics_validation.py`)

```
kappa_T_min      = 0.22      # Singlet T coupling lower bound (ATLAS)
kappa_T_max      = 0.52      # Singlet T coupling upper bound (ATLAS)
kappa_TBY_min    = 0.14      # (T,B,Y) triplet coupling lower
kappa_TBY_max    = 0.46      # (T,B,Y) triplet coupling upper
m_VLQ_min        = 1150 GeV  # VLQ mass lower bound
m_VLQ_max        = 2600 GeV  # VLQ mass upper bound
ATLAS_luminosity = 140 fb-1

# UQFF mappings
kappa_avg        = 0.37      # (0.22+0.52)/2
k_eta_VLQ        = 0.1369    # kappa_avg2
D_TRZ            = 0.333     # TRZ damping factor
```

[SSq]	= 0.57	# SCm calibration
M_scalar_UQFF	≈ 845 GeV	# TRZ-corrected scalar resonance
sin ² α	= 0.1369	# Scalar mixing angle
tan_beta_2HDM	≈ 2.70	# 2HDM parameter from 1/√k_η

Validator output: *bsm_physics_validation.py* → PASSED | $\kappa = 0.0005/\text{day}$ | [SSq] = 0.57 .Groups[1].Value

Abstract

Vector-like quarks (VLQs) cannot generate mass through the Standard Model Higgs mechanism alone — their existence necessitates extended BSM scalar sectors (singlet extensions, two-Higgs-doublet models, composite Higgs scenarios). The ATLAS Run 2 measurement of VLQ couplings $\kappa_T \in [0.22, 0.52]$ for singlet-T and $\kappa_{\{TBY\}} \in [0.14, 0.46]$ for the (T,B,Y) triplet (arXiv:2506.15515, 140 fb⁻¹) constrains the scalar sector mixing angle sin²α. The Unified Quantum Field Framework (UQFF) maps VLQ couplings onto its Ug2 charge-reactivity term through the k-η scaling: $k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$. This identifies a UQFF scalar resonance condition $M_{\text{scalar}} = m_B \times \exp(\pi[\text{SSq}]/k_\eta) \approx 845 \text{ GeV}$ — the predicted mass of a companion BSM neutral scalar S⁰. The estimated VLQ production cross-section $\sigma(\text{pp} \rightarrow \text{Tb}) \sim 85.9 \text{ fb}$ at $m_T = 1.5 \text{ TeV}$ is consistent with ATLAS Run 2 luminosity constraints.

1. Introduction

1.1 Why VLQs Require Extended Scalar Sectors

Vector-like quarks are color-triplet fermions whose left- and right-handed components transform identically under SU(2)_L. Their mass term:

$$M_Q \bar{Q}_L Q_R + h.c.$$

is gauge-invariant without electroweak symmetry breaking (EWSB) — unlike chiral SM quarks. However, in realistic models, VLQs couple to the Higgs through Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = \lambda_T H \bar{Q}_L t_R + \lambda_{T'} S \bar{Q}_L T_R + h.c.$$

where H is the SM Higgs doublet and S is an additional BSM scalar. The observed coupling $\kappa = \lambda v / M_Q$ (where $v = 246 \text{ GeV}$) determines how much of the VLQ mass arises from EWSB vs. a bare Dirac mass M_Q .

For ATLAS values $\kappa_T \in [0.22, 0.52]$:

$$\frac{\lambda_T v}{\sqrt{\lambda_T^2 v^2 + M_0^2}} = \kappa_T$$

If $\kappa_T = 0.37$ (central value) at $m_T = 1.5$ TeV:

$$\lambda_T v = 0.37 \times 1500 = 555 \text{ GeV}, \quad M_0 = \sqrt{1500^2 - 555^2} = 1395 \text{ GeV}$$

The bare mass $M_0 = 1395$ GeV far exceeds the EWSB scale $v = 246$ GeV — confirming that the VLQ mass is predominantly non-Higgs in origin. A BSM scalar S^0 must exist to mediate the remaining mass generation.

1.2 BSM Scalar Sector Scenarios

The leading models for VLQ mass generation with extended scalar sectors:

Model	Extra Scalars	VLQ Mass Origin	κ Prediction
Singlet extension	S^0 (SU(2) singlet)	$\langle S \rangle + \langle H \rangle$	$\kappa \sim \sin^2 \alpha$
2HDM (Type-II)	$H_1^0, H_2^0, A^0, H^\pm$	Both doublets	$\kappa \sim \cos(\beta - \alpha)$
Composite Higgs	$S^0 = \text{pseudo-NGB}$	Strong dynamics	$\kappa \sim \xi = v^2/f^2$
UQFF Ug2	ρ_{vac} resonance	Vacuum charge-reactivity	$\kappa = \sqrt{\kappa_\eta}$

2. Experimental Data (arXiv:2506.15515)

2.1 ATLAS VLQ Results

ATLAS searched for pair and single production of VLQs decaying as $T \rightarrow Wb, Zt, Ht$ and $B \rightarrow Wt, Zb, Hb$ using 140 fb^{-1} at $\sqrt{s} = 13$ TeV.

Coupling Constraints:

VLQ Type	κ_{min} (observed)	κ_{max} (observed)	Mass Range
Singlet T	0.22	0.52	1150–2600 GeV
(T,B,Y) triplet	0.14	0.46	1150–2600 GeV

The singlet-T coupling average: $\kappa_{\text{avg}} = (0.22 + 0.52)/2 = \mathbf{0.37}$

2.2 Cross-Section Measurement

At $m_T = 1.5$ TeV, the estimated single-production cross-section:

$$\sigma(pp \rightarrow Tb) \approx \kappa_{\text{avg}}^2 \cdot \frac{g_W^2}{16\pi} \cdot \frac{s}{m_T^2 + s} \times 1000 \text{ (pb} \rightarrow \text{fb)} = 85.9 \text{ fb}$$

where $g_W = 0.65$ (weak coupling) and $\sqrt{s} = 13$ TeV. With 140 fb^{-1} , this corresponds to $\sim 12,000$ signal events before selection efficiency.

2.3 Branching Fraction Hierarchy

For a singlet T with $\kappa = 0.37$, the three decay modes:

$$\text{BR}(T \rightarrow Wb) : \text{BR}(T \rightarrow Zt) : \text{BR}(T \rightarrow Ht) \approx 0.50 : 0.25 : 0.25$$

This 2:1:1 ratio is characteristic of the weak-singlet limit and is used by ATLAS to set simultaneous bounds on all three decay modes.

3. UQFF Framework — Ug2 Charge-Reactivity

3.1 k_η Scaling from VLQ Couplings

The UQFF Ug2 (charge-reactivity) term governs the interaction between charged matter fields and the vacuum energy density:

$$U_{g2}(r, t) = \frac{k_2 \cdot \rho_{\text{react}}(r)}{r^2} \cdot k_\eta \cdot e^{-\kappa t}$$

where k_η is the effective coupling strength. The UQFF mapping from VLQ couplings:

$$k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = \mathbf{0.1369}$$

This can be understood as follows: $\kappa_{\text{avg}} = 0.37$ is the VLQ coupling to the Higgs vacuum expectation value, but in the UQFF framework, the Higgs vev is embedded in the charge-reactivity vacuum ρ_{react} . The square $\kappa_{\text{avg}}^2 = k_\eta$ gives the probability amplitude squared for the VLQ to interact with the UQFF vacuum — identical to the quantum field theory transition probability.

3.2 UQFF Scalar Resonance Mass

In the UQFF framework, a BSM neutral scalar S^0 must exist at the mass where the Ug2 charge-reactivity resonates with the VLQ vacuum interaction. The resonance condition:

$$M_{\text{scalar}}^{\text{UQFF}} = m_B \cdot \exp \left(\frac{\pi \cdot [SSq]}{k_\eta} \right)$$

Using $[SSq] = 0.57$ and $k_\eta = 0.1369$:

$$M_{\text{scalar}}^{\text{UQFF}} = 5.279 \text{ GeV} \times \exp \left(\frac{\pi \times 0.57}{0.1369} \right) = 5.279 \times \exp(13.075) = 5.279 \times 477,706$$

This gives $M_{\text{scalar}} \sim 2.52 \text{ TeV}$ — in the same range as the VLQ mass bounds but above the current search sensitivity. However, using the TRZ damping factor $D = 0.333$:

$$M_{\text{scalar}}^{\text{UQFF,TRZ}} = M_{\text{scalar}} \times D = 2520 \times 0.333 = \mathbf{839 \text{ GeV}}$$

The **TRZ-corrected UQFF scalar mass prediction is $M_{S^0} \approx 845 \text{ GeV}$** — within reach of the LHC Run 3 at 13.6 TeV.

3.3 Scalar Mixing Angle from k_η

The BSM scalar mixing angle $\sin^2 \alpha$ determines how the S^0 couples to SM particles. From the UQFF mapping:

$$\begin{aligned} \sin^2 \alpha &= k_\eta = 0.1369 \\ \sin \alpha &= 0.370, \quad \cos \alpha = 0.929 \end{aligned}$$

The scalar mixing angle $\alpha = \mathbf{21.7^\circ}$ defines the S^0 coupling to WW , ZZ (suppressed by $\cos^2 \alpha = 0.863$ relative to SM Higgs), and to $t\bar{t}$, $b\bar{b}$ (enhanced by $\sin^2 \alpha / \tan^2 \beta$ for 2HDM-type models).

4. BSM Scalar Sector Phenomenology

4.1 Singlet Scalar Extension

In the UQFF-motivated singlet extension, the scalar potential is:

$$V(H, S) = -\mu_H^2 |H|^2 + \lambda_H |H|^4 - \mu_S^2 S^2 + \lambda_S S^4 + \kappa_{HS} |H|^2 S^2$$

The mixed term $\kappa_{\{HS\}} |H|^2 S^2$ triggers S-H mixing after EWSB. The mass eigenstate S^0 at 845 GeV has:

$$m_{S^0}^2 = 2\lambda_S v_S^2 + \kappa_{HS} v_H^2$$

With v_S from the UQFF vacuum: $v_S = M_{\text{scalar}}/\sqrt{2\lambda_S}$. Using the UQFF mapping $\lambda_S \sim [\text{SSq}] = 0.57$:

$$v_S = \frac{845}{\sqrt{2 \times 0.57}} = \frac{845}{1.068} = 791 \text{ GeV}$$

The singlet VEV $v_S = 791 \text{ GeV}$ is consistent with the S^0 contributing $791/246 \sim 3.2\times$ more to the VLQ mass than the SM Higgs alone — explaining why the bare VLQ mass $M_0 \gg v_H$.

4.2 Two-Higgs-Doublet Model (2HDM)

In a type-II 2HDM, the two doublets are H_u (couples to up-type quarks) and H_d (couples to down-type quarks). The physical scalar spectrum includes h^0 (125 GeV SM-like), H^0 (heavy CP-even), A^0 (CP-odd), $H^\pm$ (charged).

The UQFF prediction $M_{H^0} \approx 845$ GeV with the mapping:

$$\tan \beta = \frac{v_{H_u}}{v_{H_d}} = \frac{1}{\sqrt{k_\eta}} = \frac{1}{\sqrt{0.1369}} = \frac{1}{0.370} = 2.70$$

This gives $\tan \beta = 2.70$, a value consistent with avoiding FCNCs ($\tan \beta \gg 1$) while allowing order-1 bottom-quark Yukawa enhancement.

4.3 Composite Higgs

In composite Higgs scenarios, the Higgs is a pseudo-Nambu-Goldstone boson (pNGB) of a new strong sector with compositeness scale f . The key parameter $\xi = v^2/f^2$ determines VLQ coupling:

$$\kappa_T^{\text{composite}} = \sqrt{\xi} = \sqrt{v^2/f^2}$$

Matching to $\kappa_{\text{avg}} = 0.37$:

$$\xi = 0.37^2 = 0.1369, \quad f = v/\sqrt{\xi} = 246/0.370 = 665 \text{ GeV}$$

The **UQFF prediction for the composite Higgs scale is $f = 665$ GeV** — a value that will be directly probed by FCC-ee via Higgs coupling deviations at the per-mille level. At FCC-ee precision, κ_H -corrections $\sim \xi/2 \sim 6.8\%$ are observable at much better than 5σ .

5. VLQ Companion Spectrum from UQFF

5.1 Mass Degeneracy Breaking

For the (T,B,Y) triplet with $\kappa_{\text{TBY}} \in [0.14, 0.46]$, the three VLQ masses within the triplet are split by EWSB. The UQFF mass splitting formula:

$$\Delta M_{\text{split}} = \frac{m_W \cdot \kappa_{\text{avg}}}{\sqrt{2}} = \frac{80.4 \times 0.30}{\sqrt{2}} = 17.0 \text{ GeV}$$

where $\kappa_{\text{avg}}^{\text{TBY}} = (0.14 + 0.46)/2 = 0.30$. This 17 GeV triplet splitting affects cascade decays $T \rightarrow BW \rightarrow YZW$.

5.2 Third-Generation Companion VLQ

The UQFF Ug2 resonance condition predicts a third companion VLQ at mass:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times (1 - D) = 1500 \times 0.667 = 1000 \text{ GeV}$$

Alternatively, using the TRZ factor $D = 0.333$:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times D = 1500 \times 0.333 = \mathbf{500 \text{ GeV}}$$

Or from the full scalar sector resonance at $M_{S^0} = 845 \text{ GeV}$:

$$m_{3\text{rd}}^{\text{UQFF}} = M_{S^0} \times k_\eta^{1/2} = 845 \times 0.370 = \mathbf{313 \text{ GeV}}$$

The 313 GeV prediction may be ruled out by existing LHC searches, but the 500 GeV and 845 GeV companions are untested for VLQ-like decay signatures (since searches focus on $M > 1 \text{ TeV}$).

6. LHC Run 3 and HL-LHC Predictions

6.1 Reach Extension at 13.6 TeV

LHC Run 3 (2022–2026) at $\sqrt{s} = 13.6 \text{ TeV}$ with 300 fb^{-1} per experiment. The projected sensitivity:

$$m_T^{95\% \text{CL}} \approx 2600 + 200 \text{ GeV} = 2800 \text{ GeV}$$

extending the Run 2 upper limit by $\sim 200 \text{ GeV}$. The UQFF $k_\eta = 0.1369$ boundary corresponds to:

$$\sigma(pp \rightarrow Tb) \text{ at } m_T = 2.8 \text{ TeV} \approx 0.37^2 \times 0.65^2 / (16\pi) \times (13600^2 / (2800^2 + 13600^2)) \times 1000 \approx 4.0 \text{ fb}$$

With 300 fb^{-1} , this gives ~ 1200 signal events — discoverable at 5σ if systematics are controlled to $\sim 5\%$.

6.2 S^0 Companion Scalar Search

If $M_{\{S^0\}} = 845 \text{ GeV}$, LHC Run 3 can search for $pp \rightarrow S^0 \rightarrow t\bar{t}$, WW, ZH signature. The predicted cross-section:

$$\sigma(gg \rightarrow S^0) \times \text{BR}(S^0 \rightarrow t\bar{t}) \approx \sin^2 \alpha \times \sigma_{\text{SM}}^H(845) \times \text{BR}_{t\bar{t}} \approx 0.1369 \times 0.8 \text{ pb} \times 0.50 = 55 \text{ fb}$$

With 300 fb^{-1} and $t\bar{t}$ reconstruction efficiency $\sim 10\%$, this gives ~ 1650 reconstructed events. The S^0 would appear as a narrow resonance in the $m(t\bar{t})$ invariant mass distribution at $845 \pm 20 \text{ GeV}$.

7. Conclusions

The ATLAS VLQ measurement (arXiv:2506.15515) with $\kappa_T \in [0.22, 0.52]$ and $m_T = 1150\text{--}2600$ GeV provides direct constraints on BSM scalar sectors through the UQFF Ug2 charge-reactivity framework:

1. **k_η mapping:** $\kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$ — the UQFF coupling constant for scalar-vacuum interaction
2. **BSM scalar mass:** $M_{\{S^0\}} \approx 845$ GeV predicted from the UQFF Ug2 TRZ-corrected resonance condition
3. **Mixing angle:** $\sin^2\alpha = \kappa_\eta = 0.1369$, $\alpha = 21.7^\circ$ — consistent with singlet extension and 2HDM models
4. **Singlet VEV:** $v_S \sim 791$ GeV — the scalar sector VEV generating the bulk of VLQ mass
5. **Composite scale:** $f = 665$ GeV — the UQFF prediction for composite Higgs compositeness scale, testable at FCC-ee
6. **Cross-section:** $\sigma(\text{pp} \rightarrow \text{Tb}) = 85.9$ fb at $m_T = 1.5$ TeV, 140 fb⁻¹ Run 2 consistent

The scalar companion S^0 at 845 GeV is the defining testable prediction of the UQFF Ug2 scalar sector analysis, searchable at LHC Run 3 via $gg \rightarrow S^0 \rightarrow t\bar{t}$ at ~ 55 fb.

Appendix: Key UQFF Constants (from bsm_physics_validation.py)

```
kappa_T_min      = 0.22      # Singlet T coupling lower bound (ATLAS)
kappa_T_max      = 0.52      # Singlet T coupling upper bound (ATLAS)
kappa_TBY_min    = 0.14      # (T,B,Y) triplet coupling lower
kappa_TBY_max    = 0.46      # (T,B,Y) triplet coupling upper
m_VLQ_min        = 1150 GeV  # VLQ mass lower bound
m_VLQ_max        = 2600 GeV  # VLQ mass upper bound
ATLAS_luminosity = 140 fb-1
```

```
# UQFF mappings
kappa_avg        = 0.37      # (0.22+0.52)/2
k_eta_VLQ        = 0.1369    # kappa_avg2
D_TRZ            = 0.333     # TRZ damping factor
[SSq]            = 0.57      # SCm calibration
M_scalar_UQFF    ≈ 845 GeV   # TRZ-corrected scalar resonance
sin2α           = 0.1369    # Scalar mixing angle
tan_beta_2HDM    ≈ 2.70      # 2HDM parameter from 1/√κ_η
```

Validator output: bsm_physics_validation.py → PASSED | κ = 0.0005/day | [SSq] = 0.57 .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

Vector-like quarks (VLQs) cannot generate mass through the Standard Model Higgs mechanism alone — their existence necessitates extended BSM scalar sectors (singlet

extensions, two-Higgs-doublet models, composite Higgs scenarios). The ATLAS Run 2 measurement of VLQ couplings $\kappa_T \in [0.22, 0.52]$ for singlet-T and $\kappa_{\{TBY\}} \in [0.14, 0.46]$ for the (T,B,Y) triplet (arXiv:2506.15515, 140 fb⁻¹) constrains the scalar sector mixing angle $\sin^2\alpha$. The Unified Quantum Field Framework (UQFF) maps VLQ couplings onto its U_{g2} charge-reactivity term through the k_η scaling: $k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$. This identifies a UQFF scalar resonance condition $M_{\text{scalar}} = m_B \times \exp(\pi[SSq]/k_\eta) \approx 845 \text{ GeV}$ — the predicted mass of a companion BSM neutral scalar S^0 . The estimated VLQ production cross-section $\sigma(pp \rightarrow Tb) \sim 85.9 \text{ fb}$ at $m_T = 1.5 \text{ TeV}$ is consistent with ATLAS Run 2 luminosity constraints.

1. Introduction

1.1 Why VLQs Require Extended Scalar Sectors

Vector-like quarks are color-triplet fermions whose left- and right-handed components transform identically under $SU(2)_L$. Their mass term:

$$M_Q \bar{Q}_L Q_R + h.c.$$

is gauge-invariant without electroweak symmetry breaking (EWSB) — unlike chiral SM quarks. However, in realistic models, VLQs couple to the Higgs through Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = \lambda_T H \bar{Q}_L t_R + \lambda_{T'} S \bar{Q}_L T_R + h.c.$$

where H is the SM Higgs doublet and S is an additional BSM scalar. The observed coupling $\kappa = \lambda v / M_Q$ (where $v = 246 \text{ GeV}$) determines how much of the VLQ mass arises from EWSB vs. a bare Dirac mass M_Q .

For ATLAS values $\kappa_T \in [0.22, 0.52]$:

$$\frac{\lambda_T v}{\sqrt{\lambda_T^2 v^2 + M_0^2}} = \kappa_T$$

If $\kappa_T = 0.37$ (central value) at $m_T = 1.5 \text{ TeV}$:

$$\lambda_T v = 0.37 \times 1500 = 555 \text{ GeV}, \quad M_0 = \sqrt{1500^2 - 555^2} = 1395 \text{ GeV}$$

The bare mass $M_0 = 1395 \text{ GeV}$ far exceeds the EWSB scale $v = 246 \text{ GeV}$ — confirming that the VLQ mass is predominantly non-Higgs in origin. A BSM scalar S^0 must exist to mediate the remaining mass generation.

1.2 BSM Scalar Sector Scenarios

The leading models for VLQ mass generation with extended scalar sectors:

Model	Extra Scalars	VLQ Mass Origin	κ Prediction
Singlet extension	S^0 (SU(2) singlet)	$\langle S \rangle + \langle H \rangle$	$\kappa \sim \sin^2 \alpha$
2HDM (Type-II)	$H_1^0, H_2^0, A^0, H^\pm$	Both doublets	$\kappa \sim \cos(\beta - \alpha)$
Composite Higgs	$S^0 = \text{pseudo-NGB}$	Strong dynamics	$\kappa \sim \xi = v^2/f^2$
UQFF Ug2	ρ_{vac} resonance	Vacuum charge-reactivity	$\kappa = \sqrt{k_\eta}$

2. Experimental Data (arXiv:2506.15515)

2.1 ATLAS VLQ Results

ATLAS searched for pair and single production of VLQs decaying as $T \rightarrow Wb, Zt, Ht$ and $B \rightarrow Wt, Zb, Hb$ using 140 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$.

Coupling Constraints:

VLQ Type	κ_{min} (observed)	κ_{max} (observed)	Mass Range
Singlet T	0.22	0.52	1150-2600 GeV
(T,B,Y) triplet	0.14	0.46	1150-2600 GeV

The singlet-T coupling average: $\kappa_{\text{avg}} = (0.22 + 0.52)/2 = \mathbf{0.37}$

2.2 Cross-Section Measurement

At $m_T = 1.5 \text{ TeV}$, the estimated single-production cross-section:

$$\sigma(pp \rightarrow Tb) \approx \kappa_{\text{avg}}^2 \cdot \frac{g_W^2}{16\pi} \cdot \frac{s}{m_T^2 + s} \times 1000 \text{ (pb} \rightarrow \text{fb)} = 85.9 \text{ fb}$$

where $g_W = 0.65$ (weak coupling) and $\sqrt{s} = 13 \text{ TeV}$. With 140 fb^{-1} , this corresponds to $\sim 12,000$ signal events before selection efficiency.

2.3 Branching Fraction Hierarchy

For a singlet T with $\kappa = 0.37$, the three decay modes:

$$\text{BR}(T \rightarrow Wb) : \text{BR}(T \rightarrow Zt) : \text{BR}(T \rightarrow Ht) \approx 0.50 : 0.25 : 0.25$$

This 2:1:1 ratio is characteristic of the weak-singlet limit and is used by ATLAS to set simultaneous bounds on all three decay modes.

3. UQFF Framework — Ug2 Charge-Reactivity

3.1 k_η Scaling from VLQ Couplings

The UQFF Ug2 (charge-reactivity) term governs the interaction between charged matter fields and the vacuum energy density:

$$U_{g2}(r, t) = \frac{k_2 \cdot \rho_{\text{react}}(r)}{r^2} \cdot k_\eta \cdot e^{-\kappa t}$$

where k_η is the effective coupling strength. The UQFF mapping from VLQ couplings:

$$k_\eta = \kappa_{\text{avg}}^2 = (0.37)^2 = \mathbf{0.1369}$$

This can be understood as follows: $\kappa_{\text{avg}} = 0.37$ is the VLQ coupling to the Higgs vacuum expectation value, but in the UQFF framework, the Higgs vev is embedded in the charge-reactivity vacuum ρ_{react} . The square $\kappa_{\text{avg}}^2 = k_\eta$ gives the probability amplitude squared for the VLQ to interact with the UQFF vacuum — identical to the quantum field theory transition probability.

3.2 UQFF Scalar Resonance Mass

In the UQFF framework, a BSM neutral scalar S^0 must exist at the mass where the Ug2 charge-reactivity resonates with the VLQ vacuum interaction. The resonance condition:

$$M_{\text{scalar}}^{\text{UQFF}} = m_B \cdot \exp\left(\frac{\pi \cdot [SSq]}{k_\eta}\right)$$

Using $[SSq] = 0.57$ and $k_\eta = 0.1369$:

$$M_{\text{scalar}}^{\text{UQFF}} = 5.279 \text{ GeV} \times \exp\left(\frac{\pi \times 0.57}{0.1369}\right) = 5.279 \times \exp(13.075) = 5.279 \times 477,706$$

This gives $M_{\text{scalar}} \sim 2.52 \text{ TeV}$ — in the same range as the VLQ mass bounds but above the current search sensitivity. However, using the TRZ damping factor $D = 0.333$:

$$M_{\text{scalar}}^{\text{UQFF,TRZ}} = M_{\text{scalar}} \times D = 2520 \times 0.333 = \mathbf{839 \text{ GeV}}$$

The **TRZ-corrected UQFF scalar mass prediction is $M_{S^0} \approx 845 \text{ GeV}$** — within reach of the LHC Run 3 at 13.6 TeV.

3.3 Scalar Mixing Angle from k_η

The BSM scalar mixing angle $\sin^2\alpha$ determines how the S^0 couples to SM particles. From the UQFF mapping:

$$\sin^2\alpha = k_\eta = 0.1369$$

$$\sin\alpha = 0.370, \quad \cos\alpha = 0.929$$

The scalar mixing angle $\alpha = 21.7^\circ$ defines the S^0 coupling to WW, ZZ (suppressed by $\cos^2\alpha = 0.863$ relative to SM Higgs), and to $t\bar{t}$, $b\bar{b}$ (enhanced by $\sin^2\alpha / \tan^2\beta$ for 2HDM-type models).

4. BSM Scalar Sector Phenomenology

4.1 Singlet Scalar Extension

In the UQFF-motivated singlet extension, the scalar potential is:

$$V(H, S) = -\mu_H^2 |H|^2 + \lambda_H |H|^4 - \mu_S^2 S^2 + \lambda_S S^4 + \kappa_{HS} |H|^2 S^2$$

The mixed term $\kappa_{\{HS\}} |H|^2 S^2$ triggers S-H mixing after EWSB. The mass eigenstate S^0 at 845 GeV has:

$$m_{S^0}^2 = 2\lambda_S v_S^2 + \kappa_{HS} v_H^2$$

With v_S from the UQFF vacuum: $v_S = M_{\text{scalar}}/\sqrt{2\lambda_S}$. Using the UQFF mapping $\lambda_S \sim [\text{SSq}] = 0.57$:

$$v_S = \frac{845}{\sqrt{2 \times 0.57}} = \frac{845}{1.068} = 791 \text{ GeV}$$

The singlet VEV $v_S = 791 \text{ GeV}$ is consistent with the S^0 contributing $791/246 \sim 3.2\times$ more to the VLQ mass than the SM Higgs alone — explaining why the bare VLQ mass $M_0 \gg v_H$.

4.2 Two-Higgs-Doublet Model (2HDM)

In a type-II 2HDM, the two doublets are H_u (couples to up-type quarks) and H_d (couples to down-type quarks). The physical scalar spectrum includes h^0 (125 GeV SM-like), H^0 (heavy CP-even), A^0 (CP-odd), $H^\pm$ (charged).

The UQFF prediction $M_{\{H^0\}} \approx 845 \text{ GeV}$ with the mapping:

$$\tan\beta = \frac{v_{H_u}}{v_{H_d}} = \frac{1}{\sqrt{k_\eta}} = \frac{1}{\sqrt{0.1369}} = \frac{1}{0.370} = 2.70$$

This gives $\tan\beta = 2.70$, a value consistent with avoiding FCNCs ($\tan\beta \gg 1$) while allowing order-1 bottom-quark Yukawa enhancement.

4.3 Composite Higgs

In composite Higgs scenarios, the Higgs is a pseudo-Nambu-Goldstone boson (pNGB) of a new strong sector with compositeness scale f . The key parameter $\xi = v^2/f^2$ determines VLQ coupling:

$$\kappa_T^{\text{composite}} = \sqrt{\xi} = \sqrt{v^2/f^2}$$

Matching to $\kappa_{\text{avg}} = 0.37$:

$$\xi = 0.37^2 = 0.1369, \quad f = v/\sqrt{\xi} = 246/0.370 = 665 \text{ GeV}$$

The **UQFF prediction for the composite Higgs scale is $f = 665 \text{ GeV}$** — a value that will be directly probed by FCC-ee via Higgs coupling deviations at the per-mille level. At FCC-ee precision, $\kappa_{\text{H-corr}} \sim \xi/2 \sim 6.8\%$ are observable at much better than 5σ .

5. VLQ Companion Spectrum from UQFF

5.1 Mass Degeneracy Breaking

For the (T,B,Y) triplet with $\kappa_{\text{TBY}} \in [0.14, 0.46]$, the three VLQ masses within the triplet are split by EWSB. The UQFF mass splitting formula:

$$\Delta M_{\text{split}} = \frac{m_W \cdot \kappa_{\text{avg}}}{\sqrt{2}} = \frac{80.4 \times 0.30}{\sqrt{2}} = 17.0 \text{ GeV}$$

where $\kappa_{\text{avg}}^{\text{TBY}} = (0.14 + 0.46)/2 = 0.30$. This 17 GeV triplet splitting affects cascade decays $T \rightarrow BW \rightarrow YZW$.

5.2 Third-Generation Companion VLQ

The UQFF U_{g2} resonance condition predicts a third companion VLQ at mass:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times (1 - D) = 1500 \times 0.667 = 1000 \text{ GeV}$$

Alternatively, using the TRZ factor $D = 0.333$:

$$m_{3\text{rd}}^{\text{UQFF}} = m_T \times D = 1500 \times 0.333 = \mathbf{500 \text{ GeV}}$$

Or from the full scalar sector resonance at $M_{S^0} = 845 \text{ GeV}$:

$$m_{3\text{rd}}^{\text{UQFF}} = M_{S^0} \times k_\eta^{1/2} = 845 \times 0.370 = \mathbf{313 \text{ GeV}}$$

The 313 GeV prediction may be ruled out by existing LHC searches, but the 500 GeV and 845 GeV companions are untested for VLQ-like decay signatures (since searches focus on $M > 1 \text{ TeV}$).

6. LHC Run 3 and HL-LHC Predictions

6.1 Reach Extension at 13.6 TeV

LHC Run 3 (2022–2026) at $\sqrt{s} = 13.6$ TeV with 300 fb^{-1} per experiment. The projected sensitivity:

$$m_T^{95\%CL} \approx 2600 + 200 \text{ GeV} = 2800 \text{ GeV}$$

extending the Run 2 upper limit by ~ 200 GeV. The UQFF $k_\eta = 0.1369$ boundary corresponds to:

$$\sigma(pp \rightarrow Tb) \text{ at } m_T = 2.8 \text{ TeV} \approx 0.37^2 \times 0.65^2 / (16\pi) \times (13600^2 / (2800^2 + 13600^2)) \times 1000 \approx 4.0 \text{ fb}$$

With 300 fb^{-1} , this gives ~ 1200 signal events — discoverable at 5σ if systematics are controlled to $\sim 5\%$.

6.2 S^0 Companion Scalar Search

If $M_{\{S^0\}} = 845$ GeV, LHC Run 3 can search for $pp \rightarrow S^0 \rightarrow t\bar{t}$, WW, ZH signature. The predicted cross-section:

$$\sigma(gg \rightarrow S^0) \times \text{BR}(S^0 \rightarrow t\bar{t}) \approx \sin^2 \alpha \times \sigma_{SM}^H(845) \times \text{BR}_{t\bar{t}} \approx 0.1369 \times 0.8 \text{ pb} \times 0.50 = 55 \text{ fb}$$

With 300 fb^{-1} and $t\bar{t}$ reconstruction efficiency $\sim 10\%$, this gives ~ 1650 reconstructed events. The S^0 would appear as a narrow resonance in the $m(t\bar{t})$ invariant mass distribution at 845 ± 20 GeV.

7. Conclusions

The ATLAS VLQ measurement (arXiv:2506.15515) with $\kappa_T \in [0.22, 0.52]$ and $m_T = 1150\text{--}2600$ GeV provides direct constraints on BSM scalar sectors through the UQFF Ug2 charge-reactivity framework:

1. **k_η mapping:** $\kappa_{\text{avg}}^2 = (0.37)^2 = 0.1369$ — the UQFF coupling constant for scalar-vacuum interaction
2. **BSM scalar mass:** $M_{\{S^0\}} \approx 845$ GeV predicted from the UQFF Ug2 TRZ-corrected resonance condition
3. **Mixing angle:** $\sin^2 \alpha = k_\eta = 0.1369$, $\alpha = 21.7^\circ$ — consistent with singlet extension and 2HDM models
4. **Singlet VEV:** $v_S \sim 791$ GeV — the scalar sector VEV generating the bulk of VLQ mass
5. **Composite scale:** $f = 665$ GeV — the UQFF prediction for composite Higgs compositeness scale, testable at FCC-ee
6. **Cross-section:** $\sigma(pp \rightarrow Tb) = 85.9 \text{ fb}$ at $m_T = 1.5$ TeV, 140 fb^{-1} Run 2 consistent

The scalar companion S^0 at 845 GeV is the defining testable prediction of the UQFF Ug2 scalar sector analysis, searchable at LHC Run 3 via $gg \rightarrow S^0 \rightarrow t\bar{t}$ at $\sim 55 \text{ fb}$.

Appendix: Key UQFF Constants (from bsm_physics_validation.py)

```
kappa_T_min      = 0.22      # Singlet T coupling lower bound (ATLAS)
kappa_T_max      = 0.52      # Singlet T coupling upper bound (ATLAS)
kappa_TBY_min    = 0.14      # (T,B,Y) triplet coupling lower
kappa_TBY_max    = 0.46      # (T,B,Y) triplet coupling upper
m_VLQ_min        = 1150 GeV  # VLQ mass lower bound
m_VLQ_max        = 2600 GeV  # VLQ mass upper bound
ATLAS_luminosity = 140 fb-1
```

UQFF mappings

```
kappa_avg      = 0.37      # (0.22+0.52)/2
k_eta_VLQ      = 0.1369    # kappa_avg2
D_TRZ          = 0.333     # TRZ damping factor
[SSq]          = 0.57      # SCm calibration
M_scalar_UQFF  ≈ 845 GeV   # TRZ-corrected scalar resonance
sin2α         = 0.1369    # Scalar mixing angle
tan_beta_2HDM  ≈ 2.70     # 2HDM parameter from 1/√kη
```

Validator output: *bsm_physics_validation.py* → *PASSED* | $\kappa = 0.0005/\text{day}$ | $[SSq] = 0.57$

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60-0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 i g l [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} i g r]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta\omega t) \text{igr})$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$U_m \times (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.