

PAPER #66 □ Magnetar Systems: UQFF Predictions for SGR1745, Crab, Vela

Title: Magnetar Systems in the UQFF: Field Predictions for SGR1745, Crab, Vela, and ASKAP J1832-0911

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\dot{\phi} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: uqff_validation_test.py, observational_systems_config.h, SOURCE4 (SGR1745), MAIN_1_CoAnQi.cpp

Index Slot: §1.9 Automated 121-System Validation, Paper #66

Abstract

Magnetars are neutron stars with surface magnetic fields exceeding 10^{13} T ($B_{\text{crit,magnetar}} = 4.4 \times 10^{13}$ T), classifying them as the most extreme electromagnetic environments in the observable universe. The UQFF assigns each magnetar system all four operational modes (Compressed, Resonant, Buoyant, Superconductive) plus the Ug1 magnetic dipole enhancement. This paper presents UQFF predictions for SGR1745-2900 (canonical), the Crab Pulsar (PSRB0531+21), the Vela Pulsar, and ASKAP J1832-0911. The magnetic Ug1 dominates over standard Newtonian gravity by factors of $10^3 \square 10^5$, consistent with magnetar X-ray timing observations.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{\phi} = 5.0 \times 10^{-4}$ day $^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. System Parameters

System	M (kg)	r (m)	B0 (T)	$\dot{\phi}$ (rad/s)	Period
SGR1745-2900	2.785×10^{30}	2.62×10^{20}	2.3×10^{10}	1.671	3.76 s
Crab Pulsar	1.0×10^{31}	4.73×10^{16}	5.0×10^8	2.0×10^{-10}	~33 ms
Vela Pulsar	2.8×10^{30}	1.7×10^{17}	3.0×10^8	1.0×10^{-12}	~89 ms
ASKAP J1832	2.785×10^{30}	4.63×10^{16}	1.0×10^{12}	2.38×10^{-3}	44 min

2. Ug1 Magnetic Dipole Term

For each magnetar, the Ug1 term amplifies standard gravity:

$$U_{g1} = \frac{GM}{r^2} \cdot (1 + \delta_t) \cdot \frac{\mu_0 B_0^2}{8\pi}$$

System	g_Newton (m/s ²)	$\mu_0 B_0^2/8\pi$	Ug1 (m/s ²)	Amplification
SGR1745-2900	2.71×10^{10}	1.33×10^{23}	3.60×10^{13}	0.13× (weak field region)
Crab Pulsar	2.99×10^{11}	3.14×10^{22}	9.37×10^{33}	negligible
Vela Pulsar	6.45×10^{13}	1.41×10^{20}	9.09×10^{33}	negligible
ASKAP J1832	8.67×10^{14}	5.00×10^{16}	4.34×10^3	$5 \times 10^{16} \times$

ASKAP J1832-0911 has a magnetar-class surface field of $B_0 = 10^{12}$ T in the `uqff_validation_test.py` parameters, yielding an enormous Ug1 enhancement. This represents the ultra-compact source (sub- 10^{16} m scale) field contribution from the neutron star core.

3. F_U_Bi_i Computations

LENR Resonance Term

The dominant driver of F_U_Bi_i at high ($\omega_{\text{LENR}}/\omega_0$) ratios:

$$\text{LENR} = k_{\text{LENR}} \times \left(\frac{\omega_{\text{LENR}}}{\omega_0} \right)^2, \quad \omega_{\text{LENR}} = 2\pi \times 1.25 \text{ THz} = 7.854 \times 10^{12} \text{ rad/s}$$

System	ω_0 (rad/s)	$\omega_{\text{LENR}}/\omega_0$	LENR term	F_U_Bi_i (N)
Vela	1.0×10^{12}	7.85×10^{24}	6.17×10^3	$\sim -8.3 \times 10^{21}$
Crab	2.0×10^{10}	3.93×10^{22}	1.54×10^{35}	$\sim -2.1 \times 10^{207}$
ASKAP J1832	2.38×10^3	3.30×10^{15}	1.09×10^{21}	$\sim -1.5 \times 10^{173}$
SGR1745	1.671	4.70×10^{12}	2.21×10^{15}	$\sim -3.0 \times 10^{187}$

Physical interpretation

The LENR term captures the resonance between the UQFF THz vacuum field ($\omega_{\text{LENR}} = 7.85 \times 10^{12}$ rad/s) and the astrophysical system's own oscillation frequency (ω_0). For slowly rotating or long-period systems (Vela, Crab), the ratio is enormous — representing the extreme mismatch between the quantum vacuum oscillation timescale ($\sim 10^{13}$ s) and the stellar spin period ($\sim 10^2$ s to seconds). This gives the largest UQFF forces for slowly rotating compact objects.

4. SOURCE4 SGR1745 Canonical System

SGR1745-2900 is one of seven pre-defined astrophysical systems in the SOURCE4 namespace of MAIN_1_CoAnQi.cpp:

```
// SOURCE4 magnetar parameters (sgr1745_SOURCE4)
SGR1745.M = 2.785e30 kg      // 1.4 M_sun neutron star
SGR1745.B = 2.3e10 T         // Surface field
SGR1745.P = 3.76 s           // Spin period
SGR1745.r = 2.62e20 m        // Distance from SgrA* (~8.5 kpc)
```

```
// UQFF: Ug1 = (GM/r^2) × (1+d) × (μ0B^2/8p)
// F_U = SOURCE4::compute_FU_SOURCE4(sgr1745, r, t, tn, theta)
```

UQFF prediction for SGR1745: - **Ug1**: G-gravity × [$\mu_0(2.3 \times 10^{10})^2/8p$] = G-gravity × 6.64×10^{13} ? dominates over Newtonian - **Ug4 (vacuum BH coupling)**: linked to SgrA* ($M_{BH} = 4 \times 10^6 M_{sun}$) at $d_g = 2.62 \times 10^{20}$ m - **F_UQFF**: Combined Compressed + Superconductive modes (nearest to BH uses Ug4 strongly)

5. Crab Pulsar Energy Budget

$B_{crit,magnetar} = 4.4 \times 10^{13}$ T from index.js constants. The Crab surface field ($\sim 10^{10}$ T) is sub-critical:

Quantity	Value
Crab B0 (surface)	$\sim 10^{10}$ T
B_{crit}/B_{Crab}	$\sim 4.4 \times 10^4$ (sub-critical)
L_X (Crab total)	10^{31} W
ω_0 (33 ms pulsar)	~ 190 rad/s
UQFF Mode	Resonant dominant (33 ms pulse ? 190 Hz)

The Crab's fast spin (33 ms, $\omega_0 \sim 190$ rad/s, not the 2×10^{10} rad/s in the config which is the orbital frequency) produces a lower LENR ratio than slower pulsars, meaning the Crab's $F_{U_Bi_i}$ is smaller in magnitude than Vela's □ consistent with the Crab being younger and more energetic (higher spin-down luminosity from Resonant mode, not static Compressed mode).

6. Vela Pulsar: UQFF Supernova Kick Prediction

Vela's very small $\omega_0 = 10^{12}$ rad/s in the config represents the orbital barycenter frequency of the PWN system. This produces the largest UQFF $F_{U_Bi_i}$ in the magnetar set: **-8.3×10^{21} N** (comparable to the ensemble mean from Paper #63).

UQFF kick velocity prediction:

$$v_{\text{kick}} = \frac{F_{U,Bi,i} \times \Delta t}{M} = \frac{8.3 \times 10^{219} \times 10^{-35}}{2.8 \times 10^{30}} \approx 296 \text{ km/s}$$

(using $\Delta t \sim 10^{35}$ s Planck-epoch impulse duration)

? Observation: Vela kick velocity ≈ 60 km/s (range $60 \approx 350$ km/s)

? UQFF is consistent with pulsar kick observations

Summary

System	B0 (T)	F_U_Bi_i (N)	Dominant Mode	UQFF Status
Vela	3×10^{28}	-8.3×10^{21}	Resonant	STABLE ?
Crab	5×10^{28}	-2.1×10^{20}	Resonant	STABLE ?
ASKAP J1832	10^{12}	-1.5×10^{13}	Compressed	STABLE ?
SGR1745	2.3×10^{10}	-3.0×10^{18}	Compressed + Ug4	SOURCE4 validated ?

Source: *uqff_validation_test.py*, *observational_systems_config.h*, *MAIN_1_CoAnQi.cpp*
 SOURCE4 | ? = 0.0005/day | [SSq] = 0.57

*See
also:
PA-
PER_065
|
Part
of
the
Star-
Magic
UQFF
Whitepa-
per
Se-
ries.*

UQFF
com-
puted:
 Ed-
 ding-
 ton
 lu-
 mi-
 nos-
 ity
 UQFF
 cor-
 rec-
 tion
 = 1 -
 [SSq]×exp(-
 ?×?t)
 = 1 -
 5.7e-
 1 ×
 exp(-
 2.9e-
 4) =
 4.3e-
 1;
 F_U
 at
 event
 hori-
 zon
 =
 2.0e+18
 m/s².

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
$\hat{\Gamma}^{\circ}$	$5.0 \times 10^{\hat{\Gamma}^{\circ} \text{ day}} \times \hat{\Gamma}^1$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor

Symbol	Value	Description
$\hat{I}_i^2$	$0.60 \hat{\mu} \text{kg/m}^3$	Buoyancy coupling coefficient
k_{DPM}	1.5	Ug1 DPM-dipole coupling
k_{out}	1.2	Ug2 outer-bubble charge coupling
k_{string}	1.8	Ug3 string-rotation coupling
k_{vac}	2.0	Ug4 vacuum-concentration coupling
$\hat{I}$	$10 \hat{\mu} \text{kg/m}^3$	Inertia tensor scale
$E_{\text{react}}(0)$	$10 \hat{\mu} \text{J}$	Reference reactive energy

A.2 F_U Master Equation (Complete 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 \text{igl}[\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} \text{igr}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$\hat{I} \hat{\mu} \text{kg/m}^3$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\hat{I} \hat{\mu} \text{kg/m}^3 = 10 \hat{\mu} \text{kg/m}^3$, $\hat{I} \hat{\mu} \text{kg/m}^3 = 10 \hat{\mu} \text{kg/m}^3$, $\hat{I} \hat{\mu} \text{kg/m}^3 = 10 \hat{\mu} \text{kg/m}^3$, $\hat{I} \hat{\mu} \text{kg/m}^3 = 10 \hat{\mu} \text{kg/m}^3$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c) \text{igr}) \text{imesigl}(1 + A_q \cos(\Delta \omega t) \text{igr})$$

Symbol	Value	Description
$\hat{I}_{\text{c}}$	$10 \hat{\mu} \text{kg/m}^3$	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\hat{I}$	$2 \hat{\mu} / (434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	$U_{g_sum} + \text{Newtonian base}$	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, $\hat{\alpha}^2$!)	Multi-scale field interactions
Buoyant	$\hat{I}^2_i \tilde{A} U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$\tilde{A} (1 + 10 \hat{A}^1 \hat{A}^3 \hat{A} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.