

“PAPER_{0:D3}” -f [int]# PAPER #83 □ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation,
\$n = [int]# PAPER #83 □ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation, PAPER_083

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $\delta_c \sim 0.45$. The UQFF modifies the critical density threshold δ_{c_UQFF} through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15}\text{--}10^{17}$ g may constitute part of dark matter □ the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants (? = 5.0×10^4 day?¹, [SSq] = 0.57) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR δ_c

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}} \right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24}$? negligible.

UQFF dc = 0.45 (unchanged from GR) $\square$ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82 k_B T_{\text{UQFF}} = 2.82 k_B \times 0.99 T_H = 0.99 E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$f_{\text{PBH}}^{\text{UQFF}} = f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4}$$

$$= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965$$

UQFF reduces f_{PBH} by 3.5% □ within the 10□20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` *primordial BH tests* | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$.Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $dc \sim 0.45$. The UQFF modifies the critical density threshold dc_{UQFF} through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15}\text{--}10^{17}$ g may constitute part of dark matter □ the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR dc

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}}\right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24} \approx$ negligible.

UQFF dc = 0.45 (unchanged from GR) $\square$ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82k_B T_{\text{UQFF}} = 2.82k_B \times 0.99T_H = 0.99E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$\begin{aligned} f_{\text{PBH}}^{\text{UQFF}} &= f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4} \\ &= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965 \end{aligned}$$

UQFF reduces f_{PBH} by 3.5% $\square$ within the 10 $\square$ 20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` *primordial BH tests* | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$ `.Groups[1].Value` $\square$ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[SSq] = 0.57$)

Date: March 7, 2026

Source Data: `validate_hawking_temperature.py` (primordial BH test, Test 2), `validate_phase3.py`

Index Slot: §1.11 Black Hole Physics & Hawking Radiation,

$\$n = [\text{int}] \# \text{“PAPER_}\{0:\text{D}3\}” -f [\text{int}] \# \text{PAPER \#83} \square$ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation,
\$n = [\text{int}]\$ PAPER #83 □ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation, PAPER_083

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $\delta_c \sim 0.45$. The UQFF modifies the critical density threshold δ_c through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_{\text{H}}$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15} \text{--} 10^{17} \text{ g}$ may constitute part of dark matter □ the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR δ_c

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}} \right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24}$? negligible.

UQFF dc = 0.45 (unchanged from GR) □ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) □ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82 k_B T_{\text{UQFF}} = 2.82 k_B \times 0.99 T_H = 0.99 E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$f_{\text{PBH}}^{\text{UQFF}} = f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4}$$

$$= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965$$

UQFF reduces f_{PBH} by 3.5% within the 10–20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` primordial BH tests | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$. Groups[1].Value “PAPER_{0:D3}” -f \$n

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $dc \sim 0.45$. The UQFF modifies the critical density threshold dc_{UQFF} through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15}\text{--}10^{17}$ g may constitute part of dark matter – the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4}$ day $^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR dc

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}}\right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24} \approx$ negligible.

UQFF dc = 0.45 (unchanged from GR) $\square$ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82k_B T_{\text{UQFF}} = 2.82k_B \times 0.99T_H = 0.99E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$\begin{aligned} f_{\text{PBH}}^{\text{UQFF}} &= f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4} \\ &= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965 \end{aligned}$$

UQFF reduces f_{PBH} by 3.5% $\square$ within the 10 $\square$ 20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` *primordial BH tests* | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$ `.Groups[1].Value` $\square$ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[SSq] = 0.57$)

Date: March 7, 2026

Source Data: `validate_hawking_temperature.py` (primordial BH test, Test 2), `validate_phase3.py`

Index Slot: §1.11 Black Hole Physics & Hawking Radiation, “PAPER_{0:D3}” -f [int]# PAPER #83 $\square$ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation,

\$n = [\text{int}]\$ PAPER #83 □ Primordial Black Hole Mass Distribution via UQFF

Title: Primordial Black Hole Mass Distribution: UQFF Corrections to Formation Thresholds and Present-Day Density

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($? = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: validate_hawking_temperature.py (primordial BH test, Test 2), validate_phase3.py

Index Slot: §1.11 Black Hole Physics & Hawking Radiation, PAPER_083

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $\delta_c \sim 0.45$. The UQFF modifies the critical density threshold δ_c through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_{\text{H}}$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15} \text{--} 10^{17} \text{ g}$ may constitute part of dark matter □ the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis □ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR δ_c

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}} \right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24}$? negligible.

UQFF dc = 0.45 (unchanged from GR) □ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) □ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82 k_B T_{\text{UQFF}} = 2.82 k_B \times 0.99 T_H = 0.99 E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$f_{\text{PBH}}^{\text{UQFF}} = f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4}$$

$$= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965$$

UQFF reduces f_{PBH} by 3.5% within the 10–20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` primordial BH tests | $\dot{\gamma} = 0.0005/\text{day}$ | $[SSq] = 0.57$. Groups[1].Value “PAPER_{0:D3}” -f \$n

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $dc \sim 0.45$. The UQFF modifies the critical density threshold dc_{UQFF} through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15}\text{--}10^{17}$ g may constitute part of dark matter – the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{\gamma} = 5.0 \times 10^{-4}$ day $^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR dc

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}}\right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{24} \approx$ negligible.

UQFF dc = 0.45 (unchanged from GR) $\square$ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82k_B T_{\text{UQFF}} = 2.82k_B \times 0.99T_H = 0.99E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15}\text{--}10^{17}$ g, avoiding microlensing and CMB):

$$\begin{aligned} f_{\text{PBH}}^{\text{UQFF}} &= f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4} \\ &= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965 \end{aligned}$$

UQFF reduces f_{PBH} by 3.5% within the 10–20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` primordial BH tests | $? = 0.0005/\text{day}$ | $[SSq] = 0.57$.Groups[1].Value

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $dc \sim 0.45$. The UQFF modifies the critical density threshold dc_{UQFF} through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15}\text{--}10^{17}$ g may constitute part of dark matter if the UQFF

predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($\delta_c = 5.0 \times 10^{-4}$ day⁻¹, [SSq] = 0.57) uniquely enabling this analysis $\square$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR δ_c

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}} \right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{-24}$ $\approx$ negligible.

UQFF $\delta_c = 0.45$ (unchanged from GR) $\square$ the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2\Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82 k_B T_{\text{UQFF}} = 2.82 k_B \times 0.99 T_H = 0.99 E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15} \square 10^{17}$ g, avoiding microlensing and CMB):

$$\begin{aligned} f_{\text{PBH}}^{\text{UQFF}} &= f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4} \\ &= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965 \end{aligned}$$

UQFF reduces f_{PBH} by 3.5% $\square$ within the 10 $\square$ 20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	5.70×10^{11} kg	5.73×10^{11} kg	+0.5%
Peak emission energy	E_{peak}	$0.99 E_{\text{peak}}$	-1%
f_{PBH} (dark matter)	f	$0.965f$	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` primordial BH tests | $? = 0.0005/\text{day}$ | `[SSq] = 0.57 .Groups[1].Value "PAPER_{0:D3}" -f $n`

Abstract

Primordial black holes (PBHs) form during radiation domination from density perturbations exceeding $\delta_c \sim 0.45$. The UQFF modifies the critical density threshold δ_c through the Buoyant vacuum pressure correction and extends PBH survival above the GR threshold mass $M_{\text{threshold}}$ by 3.5%. The evaporation temperature of surviving PBHs is $T_{\text{UQFF}} = 0.99 \times T_H$, slightly reducing their current gamma-ray flux. PBHs with $M \sim 10^{15} \text{--} 10^{17} \text{ g}$ may constitute part of dark matter – the UQFF predicts the dark matter fraction f_{PBH} is unaffected (the 3.5% mass threshold shift does not move PBHs into or out of the observationally allowed window).

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4 \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis – establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. PBH Formation Threshold

Standard GR δ_c

$$\delta_c = 0.45 \text{ (Harrison-Zel'dovich radiation era)}$$

UQFF Correction

The UQFF Buoyant mode adds vacuum pressure during the radiation era:

$$P_{\text{UQFF}} = P_{\text{radiation}} + P_{\text{vacuum}} = \frac{\rho_{\text{rad}} c^2}{3} + \rho_{\text{vac}} c^2 \times [UA]$$

The fractional correction:

$$\delta_c^{\text{UQFF}} = \delta_c \times \left(1 - \frac{P_{\text{vacuum}}}{P_{\text{rad}}} \right) = 0.45 \times (1 - [UA] \times z_{\text{formation}}^{-4})$$

At $z_{\text{formation}} = 106$ (typical PBH epoch): $P_{\text{vacuum}}/P_{\text{rad}} = 0.0001 \times 10^{-24}$? negligible.

UQFF $\delta_c = 0.45$ (unchanged from GR) – the formation threshold is not perturbed.

2. Survival Mass Threshold

Standard GR

$$M_{\text{threshold}}^{\text{GR}} = 5.70 \times 10^{11} \text{ kg} \quad (t_{\text{evap}} = t_{\text{universe}} = 4.35 \times 10^{17} \text{ s})$$

UQFF Threshold

$$M_{\text{threshold}}^{\text{UQFF}} = M_{\text{threshold}}^{\text{GR}} \times (T_{\text{UQFF}}/T_H)^{-4/3} = 5.70 \times 10^{11} \times 0.99^{-4/3} = 5.73 \times 10^{11} \text{ kg}$$

UQFF threshold: $5.73 \times 10^{11} \text{ kg}$ (vs GR $5.70 \times 10^{11} \text{ kg}$) $\square$ 0.5% increase.

3. Present-Day PBH Gamma-Ray Flux

For PBHs near threshold ($M \sim M_{\text{threshold}}$), the Hawking radiation contributes to the diffuse gamma-ray background:

$$\frac{d^2 \Phi_\gamma}{dE d\Omega} = \frac{1}{4\pi} \int_0^\infty n_{\text{PBH}}(M) \frac{d^2 N_\gamma}{dE dt}(M, T_{\text{UQFF}}) dM$$

The UQFF $T_{\text{UQFF}} = 0.99 T_H$ reduces peak gamma-ray energy by 1%:

$$E_{\text{peak}}^{\text{UQFF}} = 2.82 k_B T_{\text{UQFF}} = 2.82 k_B \times 0.99 T_H = 0.99 E_{\text{peak}}^{\text{GR}}$$

Effect on observational bounds: UQFF shifts the PBH mass-density observational exclusion window by 0.5% in M . This does not conflict with any Fermi-LAT, INTEGRAL, or CMB spectral distortion constraint.

4. PBH dark matter fraction f_{PBH}

For M_{PBH} in the asteroid belt window ($10^{15} \square 10^{17} \text{ g}$, avoiding microlensing and CMB):

$$\begin{aligned} f_{\text{PBH}}^{\text{UQFF}} &= f_{\text{PBH}}^{\text{GR}} \times \frac{M_{\text{threshold}}^{\text{UQFF}}}{M_{\text{threshold}}^{\text{GR}}} \times \frac{T_{\text{UQFF}}^4}{T_H^4} \\ &= f_{\text{PBH}} \times 1.005 \times 0.96 = f_{\text{PBH}} \times 0.965 \end{aligned}$$

UQFF reduces f_{PBH} by 3.5% $\square$ within the 10 $\square$ 20% uncertainty on current PBH dark matter constraints.

Summary

PBH Property	Standard GR	UQFF	Change
Formation threshold dc	0.45	0.45 (unchanged)	$< 10^{-24}$
Survival mass threshold	$5.70 \times 10^{11} \text{ kg}$	$5.73 \times 10^{11} \text{ kg}$	+0.5%

PBH Property	Standard GR	UQFF	Change
Peak emission energy	E_peak	0.99 E_peak	-1%
f_PBH (dark matter)	f	0.965f	-3.5%
Fermi-LAT constraints	Not violated	Not violated	Compatible

Source: `validate_hawking_temperature.py` primordial BH tests | $\tau = 0.0005/\text{day}$ | $[SSq] = 0.57$

UQFF computed: Eddington luminosity UQFF correction = $1 - [SSq] \times \exp(-\tau \times t) = 1 - 5.7e-1 \times \exp(-2.9e-4) = 4.3e-1$; F_U at event horizon = $2.0e+18 \text{ m/s}^2$.

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
$\hat{\Gamma}^0$	$5.0 \hat{\Delta} \times 10 \hat{\Delta} \times \hat{\Delta} \times \text{day} \times \hat{\Delta}^1$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
$\hat{\Gamma}^2_i$	$0.60 \hat{\Delta} \times 0.61$	Buoyancy coupling coefficient
$k \hat{\Delta} \times \hat{\Delta}$	1.5	Ug1 DPM-dipole coupling
$k \hat{\Delta} \times \hat{\Delta}$	1.2	Ug2 outer-bubble charge coupling
$k \hat{\Delta} \times \hat{\Delta}$	1.8	Ug3 string-rotation coupling
$k \hat{\Delta} \times \hat{\Delta}$	2.0	Ug4 vacuum-concentration coupling
$\hat{I} \cdot$	$10 \hat{\Delta} \times \hat{\Delta}^2 \hat{\Delta}^2$	Inertia tensor scale
$E_{\text{react}}(0)$	$10 \hat{\Delta} \times \hat{\Delta} \times \hat{\Delta} \times J$	Reference reactive energy

A.2 F_U Master Equation (Complete $\hat{\Delta} \times \hat{\Delta}$ 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 i g l [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} i g r]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>

Term	Description	Implementation
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
4th dissipation term (PAPER_420)	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\hat{I}_{\text{sc}} = 10 \hat{\mu} \text{ kg/m}^3$, $\hat{I}_{\text{sc}} = 10 \hat{\mu} \text{ kg/m}^3$, $\hat{I}_{\text{sc}} = 10 \hat{\mu} \text{ kg/m}^3$, $\hat{I}_{\text{sc}} = 10 \hat{\mu} \text{ kg/m}^3$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c)igr) \text{imesigl}(1 + A_q \cos(\Delta\omega t)igr)$$

Symbol	Value	Description
$\hat{I}_{\text{sc}}$	$10 \hat{\mu} \text{ kg/m}^3$	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\hat{I}_{\text{sc}}$	$2 \hat{\mu} / (434 \hat{\cdot} 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed Resonant	Ug_sum + Newtonian base 5 resonance frequencies (aDPM, aTHz, $\hat{\mu}$)	Isolated stellar/BH systems Multi-scale field interactions
Buoyant Superconductive	$\hat{I}_{\text{sc}}^2 \hat{\mu} \text{ Ubi}$ $\hat{I}_{\text{sc}} (1 + 10 \hat{\mu} \hat{\cdot} \hat{\mu} \hat{\cdot} f_H)$	Expanding nebulae, stellar winds Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.