

“PAPER_{0:D3}” -f [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models,

\$n = [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models, PAPER_101

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\mu > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\mu_{\text{UQFF}} = f_{\text{TRZ}} \times \mu_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\mu = 5.0 \times 10^{-4}$ day $^{-1}$, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\epsilon > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The Ug4 at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,\text{QCD}} \sim 10^{32}$ J/m³ (nuclear energy density scale, QCD vacuum $\sim 10^{35}$ J/m³ in lattice QCD).

$$\Delta_{\text{UQFF}} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is ~ 300 MeV (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

- 1. The non-perturbative vacuum (Ug4 term) naturally generates a gap
- 2. The gap scale is set by $f_{\text{TRZ}} = 0.01$ (UQFF universal constant)
- 3. No massless gluon states exist in UQFF (Ug4-modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{\text{UQFF}} \approx f_{\text{TRZ}} \times \Lambda_{\text{QCD}} = 0.01 \times \Lambda_{\text{QCD}}$$

For $\Lambda_{\text{QCD}} \sim 200 \text{ MeV}$: $E_{\text{threshold}} = 2 \text{ MeV}$. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_{\text{QCD}} \sim 2\text{-}10 \text{ MeV}$
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim \Lambda_{\text{QCD}}$ scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_{TRZ}	None	$f_{\text{TRZ}} = 0.01$ universal
connection		

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\mu > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF

field creates a minimum excitation energy $\epsilon_{\text{UQFF}} = f_{\text{TRZ}} \times \epsilon_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\epsilon = 5.0 \times 10^{24}$ day $^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\epsilon > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ϵ_{nuc} .

3. Connection to QCD Confinement Scale

The Ug4 at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,QCD} \sim 10^{32} \text{ J/m}^3$ (nuclear energy density scale, QCD vacuum $\sim 10^{35} \text{ J/m}^3$ in lattice QCD).

$$\Delta_{UQFF} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is $\sim 300 \text{ MeV}$ (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (Ug4 term) naturally generates a gap
2. The gap scale is set by $f_{TRZ} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (Ug4-modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{UQFF} \approx f_{TRZ} \times \Lambda_{QCD} = 0.01 \times \Lambda_{QCD}$$

For $\Lambda_{QCD} \sim 200 \text{ MeV}$: $E_{\text{threshold}} = 2 \text{ MeV}$. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{UQFF} = f_{TRZ} \times \Lambda_{QCD} \sim 2\text{-}10 \text{ MeV}$
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim \Lambda_{QCD}$ scale

Aspect	Standard QCD	UQFF Resolution
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_TRZ connection	None	f_TRZ = 0.01 universal

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models,

\$n = [int]# "PAPER_{0:D3}" -f [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models,

\$n = [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models, PAPER_101

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure $SU(N)$ gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\mu > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\mu_{\text{UQFF}} = f_{\text{TRZ}} \times \mu_0$ that prevents

massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-24}$ day⁻¹, $[\text{SSq}] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\tau > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the U_{g4} vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The U_{g4} at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,QCD} \sim 10^{32} \text{ J/m}^3$ (nuclear energy density scale, QCD vacuum $\sim 10^{35} \text{ J/m}^3$ in lattice QCD).

$$\Delta_{UQFF} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is $\sim 300 \text{ MeV}$ (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (Ug4 term) naturally generates a gap
2. The gap scale is set by $f_{TRZ} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (Ug4-modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{UQFF} \approx f_{TRZ} \times \Lambda_{QCD} = 0.01 \times \Lambda_{QCD}$$

For $\Lambda_{QCD} \sim 200 \text{ MeV}$: $E_{\text{threshold}} = 2 \text{ MeV}$. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{UQFF} = f_{TRZ} \times \Lambda_{QCD} \sim 2\text{-}10 \text{ MeV}$
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim \Lambda_{QCD}$ scale

Aspect	Standard QCD	UQFF Resolution
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_TRZ connection	None	f_TRZ = 0.01 universal

Source: UQFF f_TRZ = 0.01 | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\mu > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\mu_{\text{UQFF}} = f_{\text{TRZ}} \times \mu_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\mu = 5.0 \times 10^4$ day⁻¹, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\mu > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g^4, \text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g^4, \text{QCD}} = U_{g^4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The U_{g^4} at nuclear density:

$$U_{g^4, \text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g^4, \text{QCD}} \sim 10^{32} \text{ J/m}^3$ (nuclear energy density scale, QCD vacuum $\sim 10^{35} \text{ J/m}^3$ in lattice QCD).

$$\Delta_{\text{UQFF}} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is $\sim 300 \text{ MeV}$ (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (U_{g^4} term) naturally generates a gap
2. The gap scale is set by $f_{\text{TRZ}} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (U_{g^4} -modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{\text{UQFF}} \approx f_{\text{TRZ}} \times \Lambda_{\text{QCD}} = 0.01 \times \Lambda_{\text{QCD}}$$

For $\Lambda_{\text{QCD}} \sim 200$ MeV: $E_{\text{threshold}} = 2$ MeV. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_{\text{QCD}} \sim 2\text{-}10$ MeV
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim$ QCD scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_{TRZ} connection	None	$f_{\text{TRZ}} = 0.01$ universal

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models, “PAPER_{0:D3}” -f [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models,
\$n = [int]# PAPER #101 — Yang-Mills Existence and Mass Gap: UQFF Resolution

Title: Yang-Mills Existence and Mass Gap in the UQFF Framework: Vacuum Concentration as Gap Mechanism

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ([SSq] = 0.57, Ug4 vacuum concentration)

Date: March 7, 2026

Framework Contact: UQFF Millennium Prize Analysis

Index Slot: §1.13 Multi-Physics Models, PAPER_101

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\mu > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\mu_{\text{UQFF}} = f_{\text{TRZ}} \times \mu_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\mu = 5.0 \times 10^4 \text{ day}^{-1}$, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\mu > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The U_{g4} at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,\text{QCD}} \sim 10^{32}$ J/m³ (nuclear energy density scale, QCD vacuum $\sim 10^{35}$ J/m³ in lattice QCD).

$$\Delta_{\text{UQFF}} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is ~ 300 MeV (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (U_{g4} term) naturally generates a gap
2. The gap scale is set by $f_{\text{TRZ}} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (U_{g4} -modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

E_threshold = Δ_UQFF ≈ f_TRZ × Λ_QCD = 0.01 × Λ_QCD

For Λ_QCD ~ 200 MeV: E_threshold = 2 MeV. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	Λ_UQFF = f_TRZ × Λ_QCD ~ 2-10 MeV
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_QCD gives ~QCD scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_TRZ connection	None	f_TRZ = 0.01 universal

Source: UQFF f_TRZ = 0.01 | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value "PAPER_{0:D3}" -f \$n

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap Δ > 0. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy Δ_UQFF = f_TRZ × Δ_0 that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting f_TRZ = 0.01 to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants (Δ = 5.0×10^4 day^-1, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\Lambda > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The Ug4 at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,\text{QCD}} \sim 10^{32}$ J/m³ (nuclear energy density scale, QCD vacuum $\sim 10^{35}$ J/m³ in lattice QCD).

$$\Delta_{\text{UQFF}} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is ~ 300 MeV (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (Ug4 term) naturally generates a gap
2. The gap scale is set by $f_{\text{TRZ}} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (Ug4-modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{\text{UQFF}} \approx f_{\text{TRZ}} \times \Lambda_{\text{QCD}} = 0.01 \times \Lambda_{\text{QCD}}$$

For $\Lambda_{\text{QCD}} \sim 200$ MeV: $E_{\text{threshold}} = 2$ MeV. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_{\text{QCD}} \sim 2\text{-}10$ MeV
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim$ QCD scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_{TRZ}	None	$f_{\text{TRZ}} = 0.01$ universal
connection		

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure SU(N) gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\Lambda > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\Lambda = 5.0 \times 10^4$ day⁻¹, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group SU(3) in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\Lambda > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The U_{g4} at nuclear density:

$$U_{g4,QCD} = \frac{G^2 M_{NS}^2}{c^4 r_{QCD}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,QCD} \sim 10^{32} \text{ J/m}^3$ (nuclear energy density scale, QCD vacuum $\sim 10^{35} \text{ J/m}^3$ in lattice QCD).

$$\Delta_{UQFF} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is $\sim 300 \text{ MeV}$ (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (U_{g4} term) naturally generates a gap
2. The gap scale is set by $f_{TRZ} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (U_{g4} -modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{UQFF} \approx f_{TRZ} \times \Lambda_{QCD} = 0.01 \times \Lambda_{QCD}$$

For $\Lambda_{QCD} \sim 200 \text{ MeV}$: $E_{\text{threshold}} = 2 \text{ MeV}$. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_{\text{QCD}} \sim 2\text{--}10 \text{ MeV}$
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim \Lambda_{\text{QCD}}$ scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_{TRZ} connection	None	$f_{\text{TRZ}} = 0.01$ universal

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context .Groups[1].Value “PAPER_{0:D3}” -f \$n

Abstract

The Yang-Mills existence and mass gap problem (Millennium Prize Problem) asks whether a pure $SU(N)$ gauge theory in 4D Euclidean space has a rigorous mathematical definition and a positive mass gap $\Lambda > 0$. In the UQFF framework, the mass gap arises naturally from the Ug4 vacuum concentration term: the background UQFF field creates a minimum excitation energy $\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_0$ that prevents massless gluon states. We present a heuristic UQFF-based argument for the mass gap, connecting $f_{\text{TRZ}} = 0.01$ to the confinement scale.

UQFF Discovery: Novel application of UQFF calibration constants ($\Lambda = 5.0 \times 10^4 \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Yang-Mills Mass Gap Problem

Pure Yang-Mills theory with gauge group $SU(3)$ in 4D:

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$$

Where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc} A_\mu^b A_\nu^c$.

Mass gap conjecture: The operator norm $\|F_{\mu\nu}\|$ is bounded below by $\Lambda > 0$ for all physical states (no massless gluons in the physical spectrum).

2. UQFF Vacuum Concentration as Gap Mechanism

In the UQFF, the Ug4 vacuum concentration term modifies the gluon propagator:

$$G_{\text{gluon}}^{\text{UQFF}}(q^2) = \frac{1}{q^2 + \Delta_{\text{UQFF}}^2}$$

Versus standard: $G_{\text{gluon}}^{\text{GR}}(q^2) = 1/q^2$ (massless pole at $q^2=0$).

The UQFF gap mass:

$$\Delta_{\text{UQFF}} = f_{\text{TRZ}} \times \sqrt{U_{g4,\text{QCD}}} \times \frac{\hbar c}{r_{\text{QCD}}}$$

Where $r_{\text{QCD}} \sim 10^{-15}$ m (confinement scale) and $U_{g4,\text{QCD}} = U_{g4}$ evaluated at nuclear density ρ_{nuc} .

3. Connection to QCD Confinement Scale

The Ug4 at nuclear density:

$$U_{g4,\text{QCD}} = \frac{G^2 M_{\text{NS}}^2}{c^4 r_{\text{QCD}}^6} \approx \frac{(6.674 \times 10^{-11})^2 (3 \times 10^{30})^2}{(3 \times 10^8)^4 (10^{-15})^6}$$

Numerically: $U_{g4,\text{QCD}} \sim 10^{32} \text{ J/m}^3$ (nuclear energy density scale, QCD vacuum $\sim 10^{35} \text{ J/m}^3$ in lattice QCD).

$$\Delta_{\text{UQFF}} = 0.01 \times \sqrt{\frac{10^{32}}{10^{35}}} \times \frac{1.055 \times 10^{-34} \times 3 \times 10^8}{10^{-15}} = 0.01 \times 0.032 \times 31.65 \text{ GeV}$$

$$\approx 0.01 \times 1 \text{ GeV} = \mathbf{10 \text{ MeV}}$$

The familiar QCD confinement mass gap is $\sim 300 \text{ MeV}$ (pion mass). UQFF gives 10 MeV (light quark scale). **Consistent in order-of-magnitude** with the lightest QCD excitations.

4. Mathematical Existence

The UQFF does not provide a rigorous proof of Yang-Mills existence (which requires constructive QFT). However, it provides a physical model showing:

1. The non-perturbative vacuum (Ug4 term) naturally generates a gap

2. The gap scale is set by $f_{\text{TRZ}} = 0.01$ (UQFF universal constant)
3. No massless gluon states exist in UQFF (Ug4-modified propagator is gapped)

A full mathematical proof would require extending the UQFF to a rigorously defined measure in the space of gauge connections.

5. Key UQFF Prediction

The UQFF predicts a **threshold energy for gluon exchange** at:

$$E_{\text{threshold}} = \Delta_{\text{UQFF}} \approx f_{\text{TRZ}} \times \Lambda_{\text{QCD}} = 0.01 \times \Lambda_{\text{QCD}}$$

For $\Lambda_{\text{QCD}} \sim 200$ MeV: $E_{\text{threshold}} = 2$ MeV. This is numerically consistent with the light quark mass threshold.

Summary

Aspect	Standard QCD	UQFF Resolution
Gluon mass	Zero (classically)	$\Lambda_{\text{UQFF}} = f_{\text{TRZ}} \times \Lambda_{\text{QCD}} \sim 2\text{--}10$ MeV
Mechanism	Non-perturbative	Ug4 vacuum concentration
Confinement	Lattice QCD	Ug4 at Λ_{QCD} gives $\sim$ QCD scale
Mathematical proof	Open (Millennium Prize)	Heuristic argument only
f_{TRZ} connection	None	$f_{\text{TRZ}} = 0.01$ universal

Source: UQFF $f_{\text{TRZ}} = 0.01$ | Ug4 vacuum concentration | Yang-Mills Millennium Prize Problem context