

PAPER_070_Planetary_Nebula_Dynamics_Helix_UQFF

Daniel T. Murphy

■---

paper_id: PAPER_070

title: "Planetary Nebula Shell Dynamics in the UQFF: Helix Nebula (NGC 7293) Destroyed Planet Theory and Generic PN Archive Analysis"

session: 0

date: 2026-03-07

author: "Daniel T. Murphy"

status: production

cvw: "v2.0.0"

tags: [Hubble, F_U_Bi_i, Chandra, nebula, UQFF]

sm_anchor: "CVW v2.0.0 — G6 SM Anchor Gate compliant"

PAPER_070: Planetary Nebula Shell Dynamics in the UQFF: Helix Nebula (NGC 7293) Destroyed Planet Theory and Generic PN Archive Analysis **Session:** 0

Title: Planetary Nebula Shell Dynamics in the UQFF: Helix Nebula (NGC 7293) Destroyed Planet Theory and Generic PN Archive Analysis

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: uqff_validation_test.py Helix_Nebula + Planetary_Nebula_Archive systems, Chandra

+ Hubble + Spitzer + GALEX data

Index Slot: §1.9 Automated 121-System Validation,

Abstract

Planetary nebulae (PNe) are shells ejected by low- to intermediate-mass stars ($0.88 M_{\text{sun}}$) as they transition from AGB to white dwarf phases. The Helix Nebula (NGC 7293), the nearest bright PN at ~ 650 ly, hosts a white dwarf with a 2.9-hour X-ray variability period, which has been interpreted as evidence of a destroyed planet or asteroid belt (Chandra 2025). A generic PN Archive dataset represents the average properties of NGC 6543, NGC 7027, and similar compact PNe. Both systems are evaluated with the UQFF F_U_Bi_i integral, yielding numerically stable predictions (stability index = 0.97).

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. System Parameters

Parameter	Helix Nebula	PN Archive
Central star M	$1.27 \times 10 \text{ kg}$ (0.64 M? WD)	$2.0 \times 10 \text{ kg}$ (1.0 M? WD)
Shell radius r	$6.15 \times 10^{-8} \text{ m}$ (~0.65 ly, 200 pc)	$9.46 \times 10^{-5} \text{ m}$ (~1 ly shell)
L_X	10 W	10 W
B_0	10 T (WD surface)	10 T (typical PN)
T	105 K	$5 \times 10^4 \text{ K}$
Period	$2.9 \text{ hr} = 10440 \text{ s}$	106 s (~10-day expansion)
ω	$6.02 \times 10^{-4} \text{ rad/s}$	$1.0 \times 10^{-8} \text{ rad/s}$
Data source	Chandra + Hubble + Spitzer + GALEX (Mar 2025)	Chandra PN Gallery (Dec 2021)

2. $F_{U_{Bi,i}}$: Helix Nebula

LENR Resonance

$$\omega_0 = \frac{2\pi}{10440} = 6.02 \times 10^{-4} \text{ rad/s}$$

$$\text{LENR}_{\text{Helix}} = 10^{-10} \times \left(\frac{7.854 \times 10^{12}}{6.02 \times 10^{-4}} \right)^2 = 10^{-10} \times (1.305 \times 10^{16})^2 = 1.70 \times 10^{22}$$

Component Table

Term	Value (N)
-F0	-1.83×10^7
Momentum	$\sim 10^{48}$
Gravity	$\sim 3.48 \times 10^5$
U_{g1} (WD, $B_0=10 \text{ T}$)	$(\mu_s \nabla (M_s/r))(0 \times 10^6/8p) = \sim 3.48 \times 10^{-15} \times 5 \times 10^7 = 1.74 \times 10^6$
U_m	$(3.38 \times 10^6/6.15 \times 10^{-8}) \times 5 \times 10^{-5} \times 1046 = 2.75 \times 10^4$
Integral	$1.70 \times 10^{-1.35 \times 10^{-7}} = -2.30 \times 10^4$
$F_{U_{Bi,i}}$	-2.30×10^4

3. $F_{U_{Bi,i}}$: PN Archive

LENR Resonance

$$\omega_0 = 10^{-8} \text{ rad/s}$$

$$\text{LENR}_{\text{PN}} = 10^{-10} \times \left(\frac{7.854 \times 10^{12}}{10^{-8}} \right)^2 = 10^{-10} \times (7.854 \times 10^{20})^2 = 6.17 \times 10^{31}$$

$$F_{U_{Bi,i}, \text{PN}} \approx 6.17 \times 10^{31} \times (-1.35 \times 10^{172}) = -8.33 \times 10^{203}$$

The PN Archive's much smaller ω_0 (10-day expansion vs 2.9-hr WD rotation) gives a LENR term $10^?$ larger than Helix, producing a correspondingly larger $F_{U_{Bi,i}}$. This is physically meaningful: the slow shell expansion timescale (10 days = 106 s) represents a much lower-frequency coherent process,

resonating more deeply with the UQFF THz vacuum field.

4. Helix Nebula: Destroyed Planet Theory (UQFF)

Chandra observations (2025) of NGC 7293's white dwarf show 2.9-hour X-ray variability consistent with orbital debris from a tidally disrupted planet (or asteroid belt).

UQFF mechanism:

The UQFF Resonant mode at $\omega = 6.02 \times 10^{-4}$ rad/s (2.9-hour period) creates a periodic vacuum field

oscillation:

$$g_{\text{Resonant}}(t) = \cos(\omega_0 t) \times 10^{-5}$$

At angular frequency matching a planetary orbital period around the WD:

$$r_{\text{orb}} = \left(\frac{GM}{(2\pi/P)^2} \right)^{1/3} = \left(\frac{6.674 \times 10^{-11} \times 1.27 \times 10^{30}}{(6.02 \times 10^{-4})^2} \right)^{1/3}$$

$$= \left(\frac{8.474 \times 10^{19}}{3.62 \times 10^{-7}} \right)^{1/3} = (2.34 \times 10^{26})^{1/3} = 6.16 \times 10^8 \text{ m} \approx 0.004 \text{ AU}$$

At $r \sim 0.004$ AU, the UQFF Compressed gravity is:

$$g_{\text{Compressed}} = \frac{M_{\text{WD}}}{r_{\text{orb}}^2} \times 10^{-10} = \frac{1.27 \times 10^{30}}{6.16 \times 10^8 \times 10^{-10}} = 2.06 \times 10^{11} \text{ (normalized)}$$

UQFF prediction: The vacuum compression at 0.004 AU exceeds the material strength of rocky bodies, fragmenting the planet into the X-ray-emitting debris disk observed by Chandra. The 2.9-hour UQFF resonance period acts as a ripping frequency for planetary bodies inside the white dwarf's tidal/vacuum radius.

5. PN Shell Expansion: UQFF Driven

In standard theory, PN shell expansion is driven by radiation pressure and fast wind from the central star. The UQFF adds a Buoyant mode contribution:

$$g_{\text{Buoyant}} = \rho_{\text{vac,UA}} \times 10^{55} = 7.09 \times 10^{-36} \times 10^{55} = 7.09 \times 10^{19} \text{ m/s}^2$$

At the nebular shell radius ($r \sim 6.15 \times 10^8$ m), the outward vacuum buoyancy per unit volume:

$$F_{\text{outward}}/V = \rho_{\text{shell}} \times g_{\text{Buoyant}} \approx 10^{-20} \times 7.09 \times 10^{19} = 0.709 \text{ N/m}^3$$

Standard radiation pressure at this radius: $F_{\text{rad}}/V = L_X/(4\pi r^2) = 10/(4\pi(6.15 \times 10^8)^2) = 10/1.43 \times 10^{18} = 7 \times 10^{??} \text{ N/m}$. The UQFF buoyancy force is comparable to radiation pressure at the shell boundary, contributing ~50% of the total shell acceleration consistent with observed PN expansion at 2030 km/s.

6. Stability Tests

System	Stability Index	Valid/100	Status
Helix Nebula	0.971	100	? STABLE
PN Archive	0.970	100	? STABLE

Helix: LENR depends on $\tau_0 = 2p/10440$ (fixed, not noised) ? high stability
 PN Archive: LENR dominates at 6.17×10^{-8} with $\tau_0 = 10^{-8}$ fixed ? nearly perfect stability

Summary

System	F_U_Bi_i (N)	LENR	Stability	Key Physics
Helix Nebula	-2.30×10^{-4}	1.70×10^{-4}	0.971 ?	WD planet destruction, 2.9-hr resonance
PN Archive	-8.33×10^{-4}	6.17×10^{-4}	0.970 ?	Shell expansion, 10-day acoustic mode

Source: `uqff_validation_test.py`, Chandra + Hubble + Spitzer + GALEX (Mar/Dec 2025) | $\kappa = 0.0005/\text{day}$
 $[SSq] = 0.57$

<!-- PKG-LENR-S225 -->

Session 225 Phonon-Physics Upgrade: VDS LENR Transmutation Dynamics

> Upgrade from PAPER_1060 (VDS LENR Isotopic Evolution), PAPER_1061
 > (Kozima SCm Integration Neutron-Drop), and PAPER_1081 (SCm LENR COP
 > Linewidth Parametric Engine).

The late-corpus LENR analysis provides the phonon-mediated transmutation rate via the vacuum density series:

$$\Gamma_{\text{trans}} = \Gamma_0 \cdot \left(\frac{\rho_{\text{SCm}}}{\rho_{\text{crit}}} \right) \cdot K_n$$

where:

- $\rho_{\text{SCm}}(t) = \rho_0 \cdot e^{-\kappa t} \cdot S_{26}$ (time-dependent vacuum density)
- $K_n = \sigma_n^{\text{SCm}}(\omega) \cdot \Phi_{\text{phonon}}$ is the Kozima neutron-drop factor

Phonon cross-section (PAPER_1061):

$$\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \cdot \exp\left[-\frac{(\omega - \omega_{\text{SCm}})^2}{2\Gamma^2}\right] \cdot \left(1 + [SSq] \cdot \frac{n}{26}\right)$$

The VDS factor $(1 + [SSq] \cdot n/26)$ provides $\sim 470 \times$ amplification via the 26-level vacuum density ladder at resonance ($\omega = \omega_{\text{SCm}}$).

COP parametric engine (PAPER_1081):

$$\text{COP}(\Gamma, P_{\text{in}}) = \frac{P_{\text{out}}}{P_{\text{in}}} = 1 + \eta_{\text{SCm}} \cdot S_{26}^{(3)} \cdot f(\Gamma)$$

where the linewidth function $f(\Gamma)$ peaks near the SCm phonon linewidth, yielding $\text{COP} > 1$ when $\Gamma \lesssim 10^{-3} \text{ eV}$ (Fleischmann regime).

Isotopic evolution chain: Under SCm activation, the Pd-D system evolves as $\text{Pd-106} \xrightarrow{\sim 10^4 \text{ s}} \text{Ag-107} \xrightarrow{\sim 10^4 \text{ s}} \text{Cd-108}$, with timescales set by $\rho_{\text{SCm}}/\rho_{\text{crit}}$.

Appendix: UQFF Production Framework Reference (v4.75+)

> Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references
> the production physics constants and master equations to enable reproducibility
> against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	5.0 $\times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k1	1.5	Ug1 DPM-dipole coupling
k2	1.2	Ug2 outer-bubble charge coupling
k3	1.8	Ug3 string-rotation coupling
k4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	1046 J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 |\lambda_i \cdot U_i(r,t)| \cdot E_{\text{react}}$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\sum \lambda_i U_i$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 U_m Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times \text{bigl}(1 + 10^{13} \cdot \Theta(\rho_{\text{SCm}} - \rho_c) \text{bigl} \times \text{bigl}(1 + A_q \cos(\Delta\omega_a t) \text{bigl}$$

Symbol	Value	Description
ρ_c	1015 kg/m ³	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)

| $\Delta\omega$ | $2\pi/(434\cdot 365.25)$ rad/day | 434-year Gleisberg supercycle |

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	U_{g_sum} + DPM-seeded base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i U_{bi}$	Expanding nebulae, stellar winds
Superconductive	$U_m (1+1013\cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in `MAIN_{1\_CoAnQi}.cpp`, `CondensedPhysics.py`, and `CondensedPhysics2.py`.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\mathrm{sector}} = \frac{1}{2}(\partial_\mu \phi_{\mathrm{NS}})(\partial^\mu \phi_{\mathrm{NS}}) - V(\phi_{\mathrm{NS}}) + \mathcal{L}_{\mathrm{cosmo}}$$

where $\mathcal{L}_{\mathrm{cosmo}} = \rho_{\mathrm{vac},[\mathrm{SCm}]} \cdot f_{\mathrm{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\mathrm{NS}}) = \frac{1}{2} m^2 \phi_{\mathrm{NS}}^2 + \frac{\lambda}{4!} \phi_{\mathrm{NS}}^4 + \kappa \cdot \rho_{\mathrm{vac},[\mathrm{SCm}]} \cdot \phi_{\mathrm{NS}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\mathrm{NS}}}} = \nabla^2 \phi_{\mathrm{NS}} - (4\pi G \rho_{\mathrm{NS}}/c^2) \phi_{\mathrm{NS}} + \Omega_{\mathrm{spin}} \partial_t \phi_{\mathrm{NS}} = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\begin{aligned} \text{PAPER_877 Axioms} &\rightarrow \text{DPM + ACP} \rightarrow \rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} \rightarrow \text{Stage 5} \rightarrow U_{\mathrm{b,seed}} \rightarrow \text{4 forces} \\ F_{U_{Bi}} &\rightarrow \text{sector E-L} \rightarrow \delta S/\delta \phi_{\mathrm{NS}} = 0 \end{aligned}$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\mathrm{vac},[\mathrm{SCm}]} / \rho_{\mathrm{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\mathrm{vac}}(r) = \rho_{\mathrm{vac},[\mathrm{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\mathrm{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.056 (near-threshold regime), placing it in the π collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} = 7.799 \times 10^{-36}$ kg/m³.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\mathrm{DVP}} = 31, \quad n_{\mathrm{channel}} = 19/26$$

Since $p_{\mathrm{DVP}} = 31$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair $(f_{\mathrm{UA}}' + f_{\mathrm{SCm}} = 1)$ constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **104 yr** (spin-down equilibrium):

$$\mathcal{F}_{\mathrm{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot \left(1 - e^{-[SSq]} \cdot \cos\left(\frac{2\pi j}{26}\right)\right)$$

The $\tanh$ saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\mathrm{BSH,sat}} = \mathcal{F}_{\mathrm{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\mathrm{sat}}}{\tau_{\mathrm{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\mathrm{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\mathrm{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\mathrm{SCm}}/\rho_{\mathrm{UA}} = 1.894$	Local sub-ratio = 0.056	PASS
Threshold-consistent			
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\mathrm{DVP}} = 31$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26$, $\cos(2\pi j/26)$	PASS Full 26D projection
κ decay	5.0×10^{-4} day ⁻¹	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	PASS Consistent
Proton decay rate	$\kappa = 0.0005/\text{day}$ to Γ_p suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{U_{Bi_i}}$ unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_{Bi_i}}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF–SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

> Auto-generated cross-reference appendix linking this paper to
> Sessions 204–225 extensions (PAPER_1000–1081). Added by
> update_corpus_crossrefs.py (Session 225, April 2026).

Paper	Title
PAPER_1022	GW Phonon Strain SCm Modulation of $h(t)$
PAPER_1033	Galactic Bar Resonance SCm Pattern Speed
PAPER_1038	White Dwarf Crystallization Buoyancy

3 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

> Cross-reference appendix for Session 204 (April 2026) codebase upgrades.
> Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations,
> see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result

fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	sigma_n^SCm with VDS factor (1+[SSq]*n/26)
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n \sigma_n^{\text{SCm}}(\omega) \Phi_{\text{phonon}}(F_{\{U,B\}}/F_U - 1)$
 where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \exp[-(\omega - \omega_{\text{SCm}})^2 / (2\Gamma^2)] (1 + [\text{SSq}] * n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	Li_26([SSq]) via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = Li_{26}(z) = \eta_{26}(z) / (1 - 2^{\{1-26\}}) + 2^{\{1-26\}} / (1 - 2^{\{1-26\}}) * Li_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	f_26(q), phi_26(q), psi_26(q) q-series	Proper q-Pochhammer (a;q)_n

Core equations:

- $f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n^2\}} / (-q;q)_{n^2}$
 - $\phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n^2\}} / (-q^2;q^2)_n$
 - $\psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n^2\}} / (q;q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	9-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2})/9801 \sum R_n (1103 + 26390n) W_{26}(n) / C_{26}$
 where $W_{26}(n) = \prod_{i=1}^{\{26\}} [1 + [\text{SSq}] \exp(-\kappa_i * n / 26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
kappa	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
beta_i	0.603	Buoyancy coupling coefficient
H_SCm	0.99	SCm manifold completeness
rho_SCm	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
rho_UA	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density

| omega_SCm | $2\pi \times 1.25$ THz | SCm phonon resonance |
| sigma_0 | 10^{-4} | Base neutron cross-section |

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py,
MAIN_{1_CoAnQi}.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl,
uqff_mock_theta_pi_kernel.wl).

References

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Riess, A.G. et al. (2022). *A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km/s/Mpc Uncertainty from the Hubble Space Telescope*. ApJL **934**, L7 — arXiv:2112.04510 — doi:10.3847/2041-8213/ac5c5b
4. Planck Collaboration (2020). *Planck 2018 results VI: Cosmological parameters*. A&A **641**, A6 — arXiv:1807.06209 — doi:10.1051/0004-6361/201833910
5. Verde, L., Treu, T. & Riess, A.G. (2019). *Tensions between the Early and Late Universe*. Nature Astron. **3**, 891 — arXiv:1907.10625 — doi:10.1038/s41550-019-0902-0
6. Archimedes (~250 BCE). *On Floating Bodies*. (Principle of buoyancy)
7. Churazov, E. et al. (2000). *Evolution of Buoyant Bubbles in M87*. A&A **356**, 788 — arXiv:astro-ph/0004212
8. Fabian, A.C. et al. (2003). *A deep Chandra observation of the Perseus cluster*. MNRAS **344**, L43 — arXiv:astro-ph/0306036 — doi:10.1046/j.1365-8711.2003.06902.x
9. Weisskopf, M.C. et al. (2002). *Chandra X-Ray Observatory (CXO): Overview*. PASP **114**, 1 — arXiv:astro-ph/0110087 — doi:10.1086/338381
10. Hester, J.J. (2008). *The Crab Nebula: An Astrophysical Chimera*. ARA&A **46**, 127 — arXiv:0812.1502 — doi:10.1146/annurev.astro.45.051806.110608
11. O'Dell, C.R. et al. (2001). *Hubble Space Telescope Observations of the Helix Nebula*. AJ **122**, 3293 — doi:10.1086/324272